



香港城市大學
City University of Hong Kong

專業 創新 胸懷全球
Professional · Creative
For The World

CityU Scholars

Pre-collapse femoral head necrosis treated by hip abduction a computational biomechanical analysis

Li, Shaochi; Liu, Yan; Zhou, Guangquan; Zhang, Wenjuan; Wei, Shengmei; He, Jiajia;
Stephen, Liao Shaoyi; Wei, Hang

Published in:

Health Information Science and Systems

Published: 01/01/2022

Document Version:

Final Published version, also known as Publisher's PDF, Publisher's Final version or Version of Record

License:

CC BY

Publication record in CityU Scholars:

[Go to record](#)

Published version (DOI):

[10.1007/s13755-022-00175-x](https://doi.org/10.1007/s13755-022-00175-x)

Publication details:

Li, S., Liu, Y., Zhou, G., Zhang, W., Wei, S., He, J., Stephen, L. S., & Wei, H. (2022). Pre-collapse femoral head necrosis treated by hip abduction: a computational biomechanical analysis. *Health Information Science and Systems*, 10, [8]. <https://doi.org/10.1007/s13755-022-00175-x>

Citing this paper

Please note that where the full-text provided on CityU Scholars is the Post-print version (also known as Accepted Author Manuscript, Peer-reviewed or Author Final version), it may differ from the Final Published version. When citing, ensure that you check and use the publisher's definitive version for pagination and other details.

General rights

Copyright for the publications made accessible via the CityU Scholars portal is retained by the author(s) and/or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights. Users may not further distribute the material or use it for any profit-making activity or commercial gain.

Publisher permission

Permission for previously published items are in accordance with publisher's copyright policies sourced from the SHERPA RoMEO database. Links to full text versions (either Published or Post-print) are only available if corresponding publishers allow open access.

Take down policy

Contact lbscholars@cityu.edu.hk if you believe that this document breaches copyright and provide us with details. We will remove access to the work immediately and investigate your claim.

RESEARCH



Pre-collapse femoral head necrosis treated by hip abduction: a computational biomechanical analysis

Shaochi Li^{1,2}, Yan Liu³, Guangquan Zhou⁴, Wenjuan Zhang⁵, Shengmei Wei⁵, Jiajia He^{4*}, Liao Shaoyi Stephen³ and Hang Wei⁶

Abstract

Background and objective: Clinical studies indicated that femoral head collapse (FHC) occurs in 90% of patients without intervention within five years after the diagnosis of femoral head necrosis (FHN). The management of the FHN is still a great challenging task. Clinical studies indicated that hip abduction as physical therapy represents an effective hip preservation method. However, the mechanism is unclear. In this study, we use computational biomechanical technology to investigate mechanical response in FHN patients with hip abduction and establish guide protocols for FHN rehabilitation.

Materials and methods: Thirty computational models were constructed for evaluating the safety of hip abduction and comparing the biomechanical performance of hip abduction for the treatment of different necrotic classifications. The distribution of principal compressive stress (PCS) and load share ratio (LSR) were computed and used for biomechanical evaluation.

Results: Before the start of physical therapy, when the size of necrotic segment is increased and located more laterally, the damage area of PCS enlarged and LSR of subchondral cortical to trabecular bone increased. As the increase of hip abduction angle, PCS of Type B transformed into Type A, PCS of Type C1 transformed into Type B, PCS of Type C2 transformed into Type C1; Except Type C2, the LSR return to normal level.

Discussion and conclusion: Stress transfer damaged pattern correlated significantly with necrotic classification. Hip abduction motions effectively enlarge the area of PCS and recover the LSR of different structures by altering motion posture during gait. The results indicated that hip abduction may be an effective physical therapy in improving hip function and interrupt the disease pathway of FHC and THA.

Keywords: Hip abduction, Stress transfer pattern, Load share ratio, Computational biomechanics, Parametric analysis, Computational models

Introduction

Femoral head necrosis (FHN) is a relatively common disorder of the joint [1], which can damage the stress transfer path and break the load share balance of femoral head. Excessive non-physiologic mechanical changes in the hips with FHN cause subchondral cortical bone failure

and the need for total hip arthroplasty (THA) [2]. Clinical studies indicated that femoral head collapse (FHC) occurs in 90% of patients without intervention within five years after the diagnosis of FHN [3]. The management of the FHN is still a great challenging task [4]. At present, THA has been considered as a standard and effective treatment option for patients with FHC in terms of pain relief and functional improvement; Hip preserving procedures as alternative treatments were developed to maintain or reconstruct the mechanical environment of

*Correspondence: hj1382844195@163.com

⁴The First Affiliated Hospital, Guangzhou University of Chinese Medicine, Guangzhou, China

Full list of author information is available at the end of the article

the femoral head and interrupt the disease path of FHC and postpone the need of THA.


Clinical doctors and researchers share a common goal of choosing safe and effective hip preserving procedures for protecting the femoral head of the patients with pre-collapse FHN [5, 6]. These common procedures include core decompression, transtrochanteric rotation osteotomy (TRO), free and non-vascularized fibular grafting, fibular allograft. Isolated core decompression will accelerate a collapse and OA progression of the femoral head because of the lack of repaired materials and biomechanical structural support during the healing of the necrosis region [7–9]. Free vascularized fibular grafting can provide immediate structural support and vascularity, but it is often associated with serious trauma, technical difficulties and longer recovery time [10, 11]. TRO can improve the biomechanical properties in daily activity and effectively decreased the average stress by either anterior or posterior rotation [12]. However, it is a technically demanding procedure associated with high complication risks. It should be noted that these methods interrupting the disease path of FHC by altering the joint shape and/or structure were invasive treatment. They have involved restricted weight-bearing and bed rest after operations, which may result in osteoporosis, slow metabolism, muscle atrophy and poor clinical curative effect. Because of these unfavorable results, there has been considerable interest in evaluating treatment regimens that will reduce trauma and complications, improve the ability of motion in daily activity and prevent FHC.

Hip abduction as physical therapy represents an effective hip preservation method [13, 14], which can adjust stress transfer pattern of femoral head and load share ratio (LSR) of different structures by altering motion posture during gait. In theory, recovery of stress transfer function and load share relationship have beneficial disease-modifying effect on FHN. However, to our knowledge this regimen has not been previously evaluated for the treatment of FHN and it is unclear how to achieve these transformations. In this study, we use computational biomechanical technology to investigate mechanical response in FHN patients with hip abduction and establish guide protocols for FHN rehabilitation.

Materials and methods

JIC classification and research hypothesis

In 2001, the Japanese Investigation Committee (JIC) revised diagnostic criteria to clarify the definition of osteonecrosis of the femoral head [15, 16]. According to the JIC classification criteria, FHN can be classified into subtypes A, B, C1 and C2, based on the location of the lesion in the weight-bearing area (as shown in Fig. 1). Type A lesions occupy the medial one-third or less of



the weight-bearing portion, while Type B lesions occupy the medial two-thirds or less of the weight-bearing portion. Type C1 lesions occupy more than the medial two-thirds of the weight-bearing portion without extending laterally to the acetabular edge. Type C2 lesions occupy more than the medial two-thirds of the weight-bearing portion and extend laterally to the acetabular edge. We have postulated that hip abduction motion can improve the ability of motion in daily activity and prevent FHC for patients who diagnosis with pre-collapse FHN. To test this hypothesis, thirty parametric computational models were generated and the parametric definition of the models allowed study of the effect of abduction on biomechanical performance.

Parametric modeling and boundary conditions

Single-legged stance was considered a representative body position for the primary models. We used an abducent angle β ($\beta = 0^\circ, 10^\circ, 15^\circ, 20^\circ, 25^\circ, 30^\circ$) along the anteroposterior axis of femur head to investigate the stress transfer pattern. The abducent variants were depicted schematically in Fig. 2. Finally, a total of 30 different computational models were used to simulate six hip abductions with five different progresses of FHN. The applied load was performed on a rigid plate that was tied to the distal femur. Constrains were applied on pubic symphysis and sacroiliac joint. Seven muscles were modeled as axial connectors muscle forces were depicted in literature [17]: adductor longus = 560 N; adductor magnus = 600 N; gluteal maximus = 550 N; gluteal medius = 700 N; gluteal minimus = 300 N; piriformis = 500 N; tensor fascia latae = 300 N.

Results

Distributions of principal compressive stress

The major function of principal compressive trabeculae in the femoral head is loading principal compressive stress (PCS) [18–20]. The distribution of PCS is an important index for evaluating the biomechanical performance of femoral head. In this work, hip abduction configurations of femoral head allow the study of PCS.





Figure 3 showed the coronal-sagittal section of the distributions of PCT in femoral head. Figure 3β1 shown that the PCS distributed along the principal compressive trabeculae in normal condition. When the size of the lesion segment is increased and located more laterally to the acetabular edge (Fig. 3δ1 → ø1 → ý1 → δ1), the damage of PCS increasingly enlarge. Especially for Type C1 and C2, the PCS areas reduce by more than a half in both coronal and sagittal sections. After physical therapy, as shown in Fig. 3β1–β6, on the healthy condition the PCS located more laterally of the femoral head when the angle of hip abduction is increased. Figure 3δ2, ø2, ý2, δ2 → ø6, ø6, ý6, δ6 showed that the PCS area in coronal-sagittal section increased to varied extent. Especially for Type C1, the distribution of PCS only located

within the femoral head below before physical therapy. When hip abduction angle is increased, the distribution of PCS was reconstructed from the top of the femoral head to the calcar.

Efficiency of hip abduction

Average stresses are calculated from all the elements on the interesting region, which could reflect the efficacy of treatment option [12]. In this study, the average stresses of anterolateral column (S1) and principal stress transfer area (S2) were computed, respectively. Load share ratio (LSR) is an important index for evaluating the bearing-capacity of the femoral head, which is defined as the ratio of S1 to S2. Figure 4 displays the relationship between the LSR and abduction motion, and the difference among normal femoral head and four subtypes FHN. Before the start of physical therapy, there is no significant difference in LSR between the normal and necrotic femoral head with small lesion segment; when the size of necrotic segment is increased and located more laterally, LSR increase significantly. These indicate that large lesion segment (Type C1 and C2) breaks the load share relationship of different structure of femoral head. After physical therapy, LSR in Type C1 gradually transforms to normal level when hip abduction angles are more lager; while LSR in Type C2 still greater than the other subtypes and normal femoral head.

Model validation

To validate the computational model at healthy conditions, the stress transfer pattern (Fig. 3β1) vs. the morphological characteristics of the principal compressive trabeculae of cadaver bone and the bone density distributions in X-rays images were compared together [17, 18]. The simulation results are close to the previous studies results in the literature.

Discussion

Hip joint is the largest weight-bearing structure of human, whose pathophysiology progression is closely related to the biomechanics. FHN is a relatively common




Fig. 4 Load share ratios of femoral head

rupture of joint homeostasis, which causes load share unbalance and the subchondral cortical bone buckle into the mechanical strength reduced area. The ideal treatment should aim at the recovery of the biomechanical performance in necrotic femoral head and interrupt the disease path of FHC and OA. The purpose of this study was to explore the effect of hip abduction motion on the biomechanical performance of pre-collapse FHN and establish guide protocols for physical therapy.

Previous studies indicated that physical therapy regimens have been useful for a variety of hip disorder [21–30]; only a few studies have investigated the effect of hip abduction on collapse risk of FHN. Lynne et al. [31] reported that a mean three-year hip survival rate of 82% in the necrotic femoral head treated with hip core decompression and physical therapy and 86% in those treated with physical therapy alone. Cui and Yuan [13] investigated the effect of Yuan’s Plastic therapy on treatment of FHN. They found a mean thirteen-month survival rate of 95.5% in the necrotic femoral head and this regimen can let patients with FHN carry weight earlier. Sun et al. [14] investigated the mechanism of biomechanics on the treatment of avascular necrosis of the femoral head with Anti-Chaplin gait by three-dimensional finite element method. The results indicated that Anti-Chaplin gait could reduce the load on the surface of femoral head and collapse risk of femoral head. The results indicated

that the step width and abduction angle can affect stress distribution on the necrotic femur head. To our knowledge, this study is the first biomechanical study to investigate the PCS distributions and LSR in the femoral head for physical therapy of FHN.

Parametric modeling is an efficient and fast modeling strategy to manipulate the attribute of the bone in biomechanical analysis, which are suitable for creating a model family. In this study, an initial model of hip were generated based the anatomy of hip joint, and then four necrotic zone were generated by change the shape of necrotic geometry as soon as the dimension value is modified based on JIC. Six abducent angle were created as soon as the value is changed. In total, thirty computational models were generated and used to simulate a healthy and four necrotic femoral heads with six abduction angle. When $\beta=0^\circ$, our results show that as the size of the smaller necrotic segment becomes larger the likelihood of the rupture of arthrosis homeostasis increases as demonstrated by increasing LSR in subchondral bone and smaller PCS area in coronal-sagittal section. The stress transfer function in subtrondral trabeculae completely lose when the size of necrotic segment is more than the medial two thirds of the weight-bearing portion. Compared with the normal LSR, LSRs of four subtypes increase approximately 0.82%, 2.86%, 18.78% and 36.03%, respectively.

According to load sharing theory [32] that if the absence of load share, hard tissue failure and nonunion rates greatly increase. When the LSR becomes larger, the likelihood of necrotic lesion healing may reduce and the risk of deterioration will increase. Previous studies have shown that Type C1/C2 FHN need surgical intervention because the risk of a collapse in the femoral head is quite high. While the results of this study provide a detailed description of the biomechanical evidence and physical therapy regimen to protect self-head of the patient with Type C1 FHN without invasive treatment. As the angle of hip abduction increases, the areas of PCS increase in both coronal and sagittal sections. PCS of Type B could be transformed into Type A, PCS of Type C1 could be transformed into Type B. PCS of Type C2 could be transformed into Type C1. The hip abduction motions also show that LSR of Type B and C1 reduce and return to the normal level when the $\beta > 20^\circ$. While the LSR of Type C2 still increases approximately 35.96% more than the normal level, even though $\beta = 30^\circ$. The results indicated that hip abduction regimen has a good biomechanical performance for the treatment of FHN and the biomechanical performance correlated significantly with necrotic classifications. It confirmed the safety of this physical treatment regimen for the patient with a necrotic lesion without extending laterally to the acetabular edge and suggested that patients with FHN may carry weight earlier to improve biomechanical property without restricted weight-bearing and bed rest. The results also suggested that hip abduction combined with other treatment regimen on Type C2 FHN may get better clinical outcome.

Conclusion

This study found that hip abduction motion added to the effectiveness of a physical therapy regimen for patients with pre-collapse FHN. The results of this study suggest that hip abduction motions apply to patients with a necrotic lesion without extending laterally to the acetabular edge. This routine use of hip abduction motion combined with surgical treatment may get better clinical outcome in Type C2 FHN.

Author contributions

SL and JH provided the concept and edited the draft of manuscript. WZ, SW and GZ conducted the literature review and wrote the first draft of the manuscript. LSS, HW and LY edited the draft of manuscript.

Declarations

Conflict of interest

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Author details

¹Institute of Systems Engineering, Macau University of Science and Technology, Taipa, China. ²Guangzhou Administration Institute, Guangzhou, China. ³Dept of Information Systems, City University of Hong Kong, Kowloon Tong, Hong Kong. ⁴The First Affiliated Hospital, Guangzhou University of Chinese Medicine, Guangzhou, China. ⁵The First Affiliated Hospital of Kashi Prefecture, Kashi, China. ⁶School of Medical Information Engineering, Guangzhou University of Chinese Medicine, Guangzhou, China.

Received: 19 January 2022 Accepted: 12 April 2022

Published online: 14 May 2022

References

- Matej D, Srecko H, Drage D, et al. Contact stress in hips with osteonecrosis of the femoral head. *Clin Orthop Relat Res*. 2006;447:92–9.
- Lai YS, Wei HW, Cheng CK. An analysis of hip replacement data of Taiwan. Read at the northern region congress of the Asia Pacific Orthopaedic Association in conjunction with the Third Annual Meeting of the Pacific Asian Society of Minimally Invasive Spine Surgery. 2003; 5–6; Taipei, Taiwan.
- Neumayr LD, Aquilar C, Earles AN, et al. Physical therapy alone compared with core decompression and physical therapy for femoral head osteonecrosis in sickle cell disease. Results of a multicenter study at a mean of three years after treatment. *J Bone Joint Surg Am*. 2006;88(12):2573–82.
- Bose VC, Baruah BD. Resurfacing arthroplasty of the hip for avascular necrosis of the femoral head: a minimum follow-up of four years. *J Bone Jt Surg*. 2010;92-B:922–8.
- Hasegawa Y, Iwata H, Torii S, et al. Vascularized pedicle bone-grafting for nontraumatic avascular necrosis of the femoral head: a 5-to 11-year follow up. *Arch Orthop Trauma Surg*. 1997;116:251–8.
- Krachow KA, Mont MA, Maar DC. Limited femoral endoprosthesis for avascular necrosis of the femoral head. *Orthop Rev*. 1993;22:457–63.
- Lee MS, Tai CL, Senan V, et al. The effect of necrotic size and rotational degree on the stress reduction in transtrochanteric rotational osteotomy for femoral head osteonecrosis—a three-dimensional finite-element simulation. *Clin Biomech*. 2006;21:969–76.
- Camp JF, Colwell CW. Core decompression of the femoral head for osteonecrosis. *J Bone Jt Surg Am*. 1986;68:1313–9.
- Koo KH, Kim R, Ko GH, et al. Preventing collapse in early osteonecrosis of the femoral head, a randomized clinical trial of core decompression. *J Bone Jt Surg Br*. 1995;77:870–974.
- Urbaniak JR, Congan PG, Gunneson EB, et al. Treatment of osteonecrosis of the femoral head with free vascularized fibular grafting: a long term follow-up study of one hundred and three hips. *J Bone Jt Surg Am*. 1995;77:6881–94.
- Wang BL, Sun W, Shi ZC, et al. Treatment of nontraumatic osteonecrosis of the femoral head using bone impaction grafting through a femoral neck window. *Int Orthop*. 2010;34:635–9.
- Mel SL, Tai CL, Senan V, et al. The effect of necrotic lesion and rotational degree on the stress reduction in transtrochanteric rotational osteotomy for femoral head osteonecrosis: a three dimensional finite element simulation. *Clin Biomech*. 2006;21:969–76.
- Cui YF, Yuan H. A clinical study of Yuan's plastic therapy for treatment of osteonecrosis of femoral head. *Chin Gen Pract*. 2005;8(21):1760–1.
- Sun KM, Wang HM, Chen RQ. Three dimensional finite element analysis on the treatment of avascular necrosis of the femoral head with Anti-Chaplin gait. *Chin J Tradit Med Traumatol Orthop*. 2006;14(2):15–7.
- Sugano N, Atsumi T, Ohzono K, et al. The 2001 revised criteria for diagnosis, classification, and staging of idiopathic osteonecrosis of the femoral head. *J Orthop Sci*. 2002;7:601–5.
- Sugano N, Kubo T, Takaoka K, et al. Diagnostic criteria for non-traumatic osteonecrosis of the femoral head: a multicenter study. *J Bone Jt Surg*. 1999;81-B:590–5.
- Sverdllova NS, Witzel U. Principles of determination and verification of muscle forces in the human musculoskeletal system: muscle force to minimize bending stress. *J Biomech*. 2010;43:387–96.
- Gwun JI, Yong KI. Computational study of Wolff's law with trabecular architecture in the human proximal femur using topology optimization. *J Biomech*. 2008;41:2353–61.

19. Gwun J, Youg KI. Computation simulation of trabecular adaptation progression in human proximal femur during growth. *J Biomech.* 2009;42:573–80.
20. Boyle C, Yong KI. Three-dimensional micro-level computational study of Wolff's law via trabecular bone remodeling in the human proximal femur using design space topology optimization. *J Biomech.* 2011;44:935–42.
21. Penrod JD, Boockvar KS, Litke A, et al. Physical therapy and mobility 2 and 6 months after hip fracture. *J Am Geriatr Soc.* 2004;52:1114–20.
22. Roddy E, Zhang W, Doherty M, et al. Evidence-based recommendations for the role exercise in the management of osteoarthritis of the hip or knee—the Move consensus. *Rheumatology.* 2005;44:67–73.
23. Smidt N, de Vet HC, Bouter LM, et al. Exercise therapy group. Effectiveness of exercise therapy: a best-evidence summary of systematic reviews. *Austral J Physiother.* 2005;51:71–85.
24. Hoeksma HL, Dekker J, Ronday HK, et al. Comparison of manual therapy and exercise therapy in osteoarthritis of the hip: a randomized clinical trial. *Arthritis Rheum.* 2004;51:722–9.
25. Jan MH, Hung JY, Lin JC, et al. Effects of a home program on strength, walking speed, and function after total hip replacement. *Arch Phys Med Rehabil.* 2004;85:1943–51.
26. Richard JVA, Luca M, Andrew TMP, et al. Hip Abduction can prevent posterior edge loading of hip replacements. *J Orthop Res.* 2013;31(8):1172–9.
27. Chang A, Hayes K, Dunlop D, et al. Hip Abduction moment and protection against medial tibiofemoral osteoarthritis progression. *Arthritis Rheum.* 2005;52(11):3515–9.
28. Lorna K. Case study: physical therapy management of hip osteoarthritis prior to total hip arthroplasty. *JOSPT.* 1997;26(1):35–8.
29. Cameron WM, Julie MW, Joshua AC, et al. Clinical outcomes following manual physical therapy and exercise for hip osteoarthritis: a case series. *JOSPT.* 2006;36(8):588–99.
30. Cynn HS, Oh JS, Kwon OY, et al. Effects of lumbar stabilization using a pressure biofeedback unit on muscle activity and lateral pelvic tilt during hip abduction in sidelying. *Arch Phys Med Rehabil.* 2006;87:1454–7.
31. Lynne DN, Christine A, Ann NE, et al. Physical therapy alone compared with core decompression and physical therapy for femoral head osteonecrosis in sickle cell disease. *J Bone Jt Surg.* 2006;88-A:2573–82.
32. McComack T, Karaikovic E, Robert WG. The load sharing classification of spine fracture. *Spine.* 1994;19:1741–4.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.