



香港城市大學  
City University of Hong Kong

專業 創新 胸懷全球  
Professional · Creative  
For The World

## CityU Scholars

### Resilience of human gut microbial communities for the long stay with multiple dietary shifts

Liu, Hong; Han, Maozhen; Li, Shuai Cheng; Tan, Guangming; Sun, Shiwei; Hu, Zhiqiang; Yang, Pengshuo; Wang, Rui; Liu, Yawen; Chen, Feng; Peng, Jianjun; Peng, Hong; Song, Hongxing; Xia, Yang; Chu, Liqun; Zhou, Quan; Guan, Feng; Wu, Jing; Bu, Dongbo; Ning, Kang

**Published in:**  
Gut

**Published:** 01/12/2019

**Document Version:**

Final Published version, also known as Publisher's PDF, Publisher's Final version or Version of Record

**License:**  
CC BY-NC

**Publication record in CityU Scholars:**  
[Go to record](#)

**Published version (DOI):**  
[10.1136/gutjnl-2018-317298](https://doi.org/10.1136/gutjnl-2018-317298)

**Publication details:**

Liu, H., Han, M., Li, S. C., Tan, G., Sun, S., Hu, Z., Yang, P., Wang, R., Liu, Y., Chen, F., Peng, J., Peng, H., Song, H., Xia, Y., Chu, L., Zhou, Q., Guan, F., Wu, J., Bu, D., & Ning, K. (2019). Resilience of human gut microbial communities for the long stay with multiple dietary shifts. *Gut*, 68(12), 2254-2255.  
<https://doi.org/10.1136/gutjnl-2018-317298>

**Citing this paper**

Please note that where the full-text provided on CityU Scholars is the Post-print version (also known as Accepted Author Manuscript, Peer-reviewed or Author Final version), it may differ from the Final Published version. When citing, ensure that you check and use the publisher's definitive version for pagination and other details.

**General rights**

Copyright for the publications made accessible via the CityU Scholars portal is retained by the author(s) and/or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights. Users may not further distribute the material or use it for any profit-making activity or commercial gain.

**Publisher permission**

Permission for previously published items are in accordance with publisher's copyright policies sourced from the SHERPA RoMEO database. Links to full text versions (either Published or Post-print) are only available if corresponding publishers allow open access.

**Take down policy**


Contact [lbscholars@cityu.edu.hk](mailto:lbscholars@cityu.edu.hk) if you believe that this document breaches copyright and provide us with details. We will remove access to the work immediately and investigate your claim.

## Resilience of human gut microbial communities for the long stay with multiple dietary shifts

We read with great interest the article by Bennet *et al*<sup>1</sup> that investigated the impact of dietary shifts on gut microbiota. Together with several recent studies,<sup>2–5</sup> they have shown that short-term dietary shifts can alter the composition of gut microbiota for both patients and healthy humans. However, these works drawn the conclusion with only a few time points and the subjects under investigation had undergone just one dietary shift. The dynamic patterns of microbial communities across longer time scales with multiple dietary shifts remain unclear.

In this study, we recruited a Chinese volunteer team (VT) composing of 10 people who departed from Beijing, conducted a long stay of 6 months in Trinidad and Tobago (TAT) and returned to Beijing. Their faecal samples (188 samples), together with detailed dietary information, were collected using a high-density longitudinal sampling strategy (19 time points on average for VT members). We partitioned the whole longitudinal study into six phases: T1 represents pre-travel phase (20 samples), T2 (28 samples), T3 (60 samples) and T4 (21 samples) represent three time-slots when VT stayed in TAT, and T5 (35 samples) and T6 (20 samples) represent two time-slots after VT returned to Beijing. Meanwhile, we also collected samples from healthy natives of Beijing (BJN, 57 samples), TAT healthy natives (TTN, 28 samples), TAT patients (TTP, six samples) and TAT Chinese (TTC, eight samples) as control datasets. Finally, we sequenced the V4 hypervariable region of microbial 16S rRNA genes for 287 faecal samples from 41 individuals and analysed the high-quality reads with QIIME.<sup>6</sup>

Interestingly, we found that gut microbial communities are bidirectionally plastic and resilient across the long stay with multiple dietary shifts. First of all, BJN and TTN exhibited different patterns of microbial communities (figure 1A). The microbial communities of VT members, however, shifted from those similar to BJN towards TTN pattern accompanying their stays in TAT,



**Figure 1** Pattern of human gut microbial community during the long stay with multiple dietary shifts. The dynamics of human gut microbial communities are shown through comparing different groups/phases of samples, including (A) BJN and TTN groups, (B) BJN, TTN and T1 groups, (C) BJN, TTN, T1 and T2-T4 groups, (D) BJN, TTN, T1, T2-T4 and T5-T6 groups, (E) BJN, TTN, T1, T2-T4, T5-T6 and TTC groups, (F) BJN, TTN, T1, T2-T4, T5-T6, TTC and TTP groups. The large arrows indicate the progressive add-in of groups for comparison. The profile of all faecal samples labelled by (G) collected locations, and (H) populations (Chinese and TAT). In (H), the label 'Chinese' represents all samples from BJN, VT and TTC, while the label 'TAT' represents all samples from TTN and TTP. The dynamic changes of gut microbiota for two members of the VT, including (I) VT5, and (J) VT6. The curves with arrows in (I) and (J) represent the directions of changes of microbial communities. Panels (K), (L), (M) and (N) show the dynamics of the average relative abundance of Firmicutes, Bacteroidetes, Proteobacteria and Actinobacteria along the temporal axis. Results were visualised using PCoA of weighted UniFrac distances of 16S rRNA amplicon profiles. OTUs shared by at least 10% VT samples during same phase (green bars) are tracked using Sankey plots for (O) Firmicutes, (P) Bacteroidetes, (Q) Proteobacteria and (R) Actinobacteria. The heights of bars are proportional to the number of OTUs, and phases are arranged in chronological order. The lines represent the transfer of OTUs between phases and are coloured according to the first phase of their appearance. The abbreviations used in this figure are listed as follows: BJN, Beijing natives; OTUs, operational taxonomic units; PCoA, principal coordinates analysis; TTN, Trinidad and Tobago natives; T1-T6, six phases; TTC, Trinidad and Tobago Chinese; TTP, Trinidad and Tobago patient; VT, volunteer team.

and reverted to their original patterns within 1 month after VT returned to Beijing (figure 1B-F). Such plastic pattern is also verified by DeBlur<sup>7</sup> analysis. Besides, though we have found that locations and populations have strong effects on differentiation of samples (figure 1G,H), the dynamic changes for each member of VT showed specific trends (figure 1I,J), indicating the

possible enterotypes-dependent plastic patterns among VT members. Additionally, the relative abundances of phylum Firmicutes and Bacteroidetes have shown a strong resilience along the timeline, and they are negatively correlated along the timeline (figure 1K,L). Similarly, the relative abundance of Proteobacteria and Actinobacteria also exhibited plastic patterns (figure 1M,N).

By tracing and comparing the common operational taxonomic units (OTUs) shared by at least 10% of VT members, we found that individual OTUs within Firmicutes, Bacteroidetes, Proteobacteria and Actinobacteria have distinctive temporal dynamics during VT's long stay (figure 1O–R).

The bi-directional plasticity of human microbial community was largely mediated by dietary shifts. During their stay in TAT, VT members consumed local foods (online supplementary table 1) mainly including fish and seafood, dairy products, and refined grains, which were considerably different from those in Beijing (online supplementary table 2). Significant correlation was observed between dietary shifts and the dynamics of gut microbial community structures ( $p < 0.05$ , Mantel's test, online supplementary table 3 and 4).

The bi-directional plasticity of gut microbiome shown in this study could elicit effects on clinical practice for gut microbiome-related diseases in at least two ways: First, it indicated that diagnostics should incorporate dietary habits and travel records. Second, it suggested that due to the strong resilience of gut microbial communities, clinical practices, such as faecal microbiota transplantation and antibiotic treatment, should monitor therapeutic effects across a longer time scale.<sup>8</sup>

Hong Liu,<sup>1,2</sup> Maozhen Han <sup>1</sup>, Shuai Cheng Li,<sup>3</sup> Guangming Tan,<sup>4,5</sup> Shiwei Sun,<sup>4,5</sup> Zhiqiang Hu,<sup>2</sup> Pengshuo Yang,<sup>1</sup> Rui Wang,<sup>2</sup> Yawen Liu,<sup>2</sup> Feng Chen,<sup>2</sup> Jianjun Peng,<sup>2</sup> Hong Peng,<sup>2</sup> Hongxing Song,<sup>2</sup> Yang Xia,<sup>2</sup> Liqun Chu,<sup>2</sup> Quan Zhou,<sup>2</sup> Feng Guan,<sup>2</sup> Jing Wu,<sup>2</sup> Dongbo Bu,<sup>4,5</sup> Kang Ning<sup>1</sup>

<sup>1</sup>Key Laboratory of Molecular Biophysics of the Ministry of Education, Hubei Key Laboratory of Bioinformatics and Molecular-imaging, Department of Bioinformatics and Systems Biology, College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan, China

<sup>2</sup>Beijing Shijitan Hospital, Capital Medical University, Beijing, China

<sup>3</sup>Department of Computer Science, City University of Hong Kong, Hong Kong, China

<sup>4</sup>Key Lab of Intelligent Information Processing, State Lab of Computer Architecture, Institute of Computing Technology, Chinese Academy of Sciences, Beijing, China

<sup>5</sup>School of Computer and Control Engineering, University of Chinese Academy of Sciences, Beijing, China

**Correspondence to** Dr Jing Wu, Beijing Shijitan Hospital, Capital Medical University, Beijing, 100038, China; wujingsjt@gmail.com, Professor Dongbo Bu, Key Lab of Intelligent Information Processing, Institute of Computing Technology, Chinese Academy of Sciences, Beijing, 100190, China; dbu@ict.ac.cn and Professor Kang Ning, Department of Bioinformatics and Systems Biology, College of Life Science and Technology, Huazhong University of Science and

Technology, Wuhan 430074, China; ningkang@hust.edu.cn

**Acknowledgements** We thank Yumin Song and all other members of Embassy of the People's Republic of China in the Republic of Trinidad and Tobago, for their support in sample collection for this work. We thank Le Cao and Chaofang Zhong, from College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan, China, for their help in preparing the figures.

**Contributors** This study was designed by HL, JW, DBB and KN. MZH, HL, RW, YWL, ZQH, FC, JJP, HP, QZ, LQC, HXS, FG and YX collected samples. MZH, HL, RW and PSY analyzed the data. MZH, HL, SWS, SCL, GMT, DBB and KN wrote the initial draft of the manuscript. All authors revised the manuscript.

**Funding** This work was partially supported by National Key Research and Development Program of China (2018YFC0910502 and 2018YFC0910405), National Natural Science Foundation of China (31671374, 31871334, 31327901, 31671369, 31770775, 61521092, 91430218, 61472395, and 61432018), Ministry of Science and Technology's high-tech (863) grants (2014AA021502, 2016YFB0201305, 2016YFB0200803 and 2016YFB0200300), and the GRF Research Project 9042348(CityU11257316).

**Competing interests** None declared.

**Patient consent** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.



## OPEN ACCESS

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/gutjnl-2018-317298>).

HL, MH and SCL contributed equally.



**To cite** Liu H, Han M, Li SC, *et al.* *Gut* 2019;**68**:2254–2255.

Received 31 July 2018

Revised 18 September 2018

Accepted 11 October 2018

Published Online First 12 November 2018

*Gut* 2019;**68**:2254–2255. doi:10.1136/gutjnl-2018-317298

### ORCID iD

Maozhen Han <http://orcid.org/0000-0002-5958-1941>

### REFERENCES

- 1 Bennet SMP, Böhn L, Störsrud S, *et al.* Multivariate modelling of faecal bacterial profiles of patients with

IBS predicts responsiveness to a diet low in FODMAPs. *Gut* 2018;**67**:872–81.

2 Claesson MJ, Jeffery IB, Conde S, *et al.* Gut microbiota composition correlates with diet and health in the elderly. *Nature* 2012;**488**:178–184.

3 David LA, Maurice CF, Carmody RN, *et al.* Diet rapidly and reproducibly alters the human gut microbiome. *Nature* 2014;**505**:559–563.

4 Chassaing B, Van de Wiele T, De Bodt J, *et al.* Dietary emulsifiers directly alter human microbiota composition and gene expression ex vivo potentiating intestinal inflammation. *Gut* 2017;**66**:1414–27.

5 Smits SA, Leach J, Sonnenburg ED, *et al.* Seasonal cycling in the gut microbiome of the Hadza hunter-gatherers of Tanzania. *Science* 2017;**357**:802–6.

6 Caporaso JG, Kuczynski J, Stombaugh J, *et al.* QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* 2010;**7**:335–6.

7 Amir A, McDonald D, Navas-Molina JA, *et al.* Deblur rapidly resolves single-nucleotide community sequence patterns. *mSystems* 2017;**2**:e00191–16.

8 Sommer F, Anderson JM, Bharti R, *et al.* The resilience of the intestinal microbiota influences health and disease. *Nat Rev Microbiol* 2017;**15**:630–8.