



香港城市大學
City University of Hong Kong

專業 創新 胸懷全球
Professional · Creative
For The World

CityU Scholars

WELL-POSEDNESS FOR MOVING INTERFACES WITH SURFACE TENSION IN IDEAL COMPRESSIBLE MHD

TRAKHININ, Yuri; WANG, Tao

Published in:
SIAM Journal on Mathematical Analysis

Published: 01/12/2022

Document Version:
Final Published version, also known as Publisher's PDF, Publisher's Final version or Version of Record

Publication record in CityU Scholars:
[Go to record](#)

Published version (DOI):
[10.1137/22M1488429](https://doi.org/10.1137/22M1488429)

Publication details:
TRAKHININ, Y., & WANG, T. (2022). WELL-POSEDNESS FOR MOVING INTERFACES WITH SURFACE TENSION IN IDEAL COMPRESSIBLE MHD. *SIAM Journal on Mathematical Analysis*, 54(6), 5888-5921. Advance online publication. <https://doi.org/10.1137/22M1488429>

Citing this paper

Please note that where the full-text provided on CityU Scholars is the Post-print version (also known as Accepted Author Manuscript, Peer-reviewed or Author Final version), it may differ from the Final Published version. When citing, ensure that you check and use the publisher's definitive version for pagination and other details.

General rights

Copyright for the publications made accessible via the CityU Scholars portal is retained by the author(s) and/or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights. Users may not further distribute the material or use it for any profit-making activity or commercial gain.

Publisher permission

Permission for previously published items are in accordance with publisher's copyright policies sourced from the SHERPA RoMEO database. Links to full text versions (either Published or Post-print) are only available if corresponding publishers allow open access.

Take down policy

Contact lbscholars@cityu.edu.hk if you believe that this document breaches copyright and provide us with details. We will remove access to the work immediately and investigate your claim.

WELL-POSEDNESS FOR MOVING INTERFACES WITH SURFACE TENSION IN IDEAL COMPRESSIBLE MHD*

YURI TRAKHININ[†] AND TAO WANG[‡]

Abstract. We study the local well-posedness for an interface with surface tension that separates a perfectly conducting inviscid fluid from a vacuum. The fluid flow is governed by the equations of three-dimensional ideal compressible magnetohydrodynamics (MHD), while the vacuum magnetic and electric fields are supposed to satisfy the pre-Maxwell equations. The fluid and vacuum magnetic fields are tangential to the interface. This renders a nonlinear hyperbolic-elliptic coupled problem with a characteristic free boundary. We introduce some suitable regularization to establish the solvability and tame estimates for the linearized problem. Combining the linear well-posedness result with a modified Nash–Moser iteration scheme, we prove the local existence and uniqueness of solutions of the nonlinear problem. The non-collinearity condition required by Secchi and Trakhinin [*Nonlinearity*, 27 (2014), pp. 105–169] for the case of zero surface tension becomes unnecessary in our result, which verifies the stabilizing effect of surface tension on the evolution of moving vacuum interfaces in ideal compressible MHD.

Key words. ideal compressible MHD, pre-Maxwell equations, surface tension, moving interface, well-posedness

MSC codes. 76W05, 35L65, 35R35

DOI. 10.1137/22M1488429

1. Introduction. We study the local well-posedness for an interface $\Sigma(t)$ with surface tension that separates a perfectly conducting inviscid fluid (e.g., plasma or liquid metal) from a vacuum. In the moving domain $\Omega^+(t) \subset \mathbb{R}^3$, we consider the following equations of ideal compressible magnetohydrodynamics (MHD) describing the motion of an inviscid perfectly conducting fluid interacting with a magnetic field (see Landau and Lifshitz [14, section 65]):

$$(1.1) \quad \begin{cases} \partial_t \rho + \nabla \cdot (\rho v) = 0, \\ \partial_t (\rho v) + \nabla \cdot (\rho v \otimes v - H \otimes H) + \nabla q = 0, \\ \partial_t H - \nabla \times (v \times H) = 0, \\ \partial_t (\rho \mathbf{e} + \frac{1}{2} \rho |v|^2 + \frac{1}{2} |H|^2) + \nabla \cdot (v(\rho \mathbf{e} + \frac{1}{2} \rho |v|^2 + p) + H \times (v \times H)) = 0, \end{cases}$$

together with the divergence constraint

$$(1.2) \quad \nabla \cdot H = 0.$$

*Received by the editors April 4, 2022; accepted for publication (in revised form) August 5, 2022; published electronically November 9, 2022.

<https://doi.org/10.1137/22M1488429>

Funding: The work of the first author was supported by the Mathematical Center in Akademgorodok under grant 075-15-2022-282 from the Ministry of Science and Higher Education of the Russian Federation. The work of the second author was supported by the Fundamental Research Funds for the Central Universities grant 2042022kf1183, the National Natural Science Foundation of China grants 11731008 and 11971359, and the Hong Kong Institute for Advanced Study grant 9360157.

[†]Sobolev Institute of Mathematics, Koptyug av. 4, 630090 Novosibirsk, Russia; Novosibirsk State University, Pirogova str. 1, 630090 Novosibirsk, Russia (trakhin@math.nsc.ru).

[‡]Corresponding author. School of Mathematics and Statistics, Wuhan University, Wuhan 430072, China; Department of Mathematics, City University of Hong Kong, Hong Kong, China (tao.wang@whu.edu.cn).

Here density ρ , fluid velocity $v = (v_1, v_2, v_3)^\top$, magnetic field $H = (H_1, H_2, H_3)^\top$, and pressure p are unknown functions of time t and space variable $x = (x_1, x_2, x_3)$. We denote by $q = p + \frac{1}{2}|H|^2$ the total pressure. According to thermodynamics, the internal energy ϵ and the density ρ are given functions of the pressure p and the entropy S , which renders the system of equations (1.1) closed for the unknowns $U := (q, v, H, S)^\top \in \mathbb{R}^8$.

Since the displacement current is neglected in the MHD equations (1.1)–(1.2), we assume that in the vacuum domain $\Omega^-(t) \subset \mathbb{R}^3$, the magnetic field $h = (h_1, h_2, h_3)^\top$ and the electric field $e = (e_1, e_2, e_3)^\top$ satisfy the so-called pre-Maxwell equations, that is, Maxwell's equations in vacuum without the displacement current [3]:

$$(1.3) \quad \nabla \times h = 0, \quad \nabla \cdot h = 0,$$

$$(1.4) \quad \nabla \times e = -\partial_t h, \quad \nabla \cdot e = 0.$$

The vacuum electric field e in (1.3)–(1.4) is a secondary variable, so that only one basic variable is needed, viz., h , satisfying the elliptic system (1.3).

Let $\Omega := (-1, 1) \times \mathbb{T}^2$ be the reference domain occupied by the conducting fluid and the vacuum, where \mathbb{T}^2 denotes the 2-torus and will be thought of as the unit square with periodic boundary conditions. For simplicity, we assume that the moving interface $\Sigma(t)$ has the form of a graph $\Sigma(t) := \{x_1 = \varphi(t, x')\} \cap \Omega$ with $x' = (x_2, x_3)$, where the interface function $\varphi : \mathbb{R}_+ \times \mathbb{T}^2 \rightarrow (-1, 1)$ is to be determined. The general case of the moving interface can be treated by the standard but more technically involved arguments as in [16, section 2] and [28, Remark 2.3]. Let us denote by $\Sigma^\pm := \{\pm 1\} \times \mathbb{T}^2$ the fixed boundaries of the domains $\Omega^\pm(t) := \{x_1 \gtrless \varphi(t, x')\} \cap \Omega$, respectively. Then the boundary conditions read as

$$(1.5a) \quad \partial_t \varphi = v \cdot N \quad \text{on } \Sigma(t),$$

$$(1.5b) \quad q - \frac{1}{2}|h|^2 = \mathfrak{s}\mathcal{H}(\varphi) \quad \text{on } \Sigma(t),$$

$$(1.5c) \quad H \cdot N = 0, \quad h \cdot N = 0 \quad \text{on } \Sigma(t),$$

$$(1.5d) \quad H_1 = 0, \quad v_1 = 0 \quad \text{on } \Sigma^+,$$

$$(1.5e) \quad h \times \mathbf{e}_1 = \mathbf{j}_c \quad \text{on } \Sigma^-,$$

with $N := (1, -\partial_2 \varphi, -\partial_3 \varphi)^\top$ and $\mathbf{e}_1 := (1, 0, 0)^\top$. Here \mathbf{j}_c is a given smooth vector function, $\mathfrak{s} > 0$ is the constant coefficient of surface tension, and $\mathcal{H}(\varphi)$ is twice the mean curvature of $\Sigma(t)$ defined by

$$(1.6) \quad \mathcal{H}(\varphi) := D_{x'} \cdot \left(\frac{D_{x'} \varphi}{\sqrt{1 + |D_{x'} \varphi|^2}} \right) \quad \text{with } D_{x'} := \begin{pmatrix} \partial_2 \\ \partial_3 \end{pmatrix}.$$

Condition (1.5a) states that the interface moves with the motion of the conducting fluid, which renders the free interface $\Sigma(t)$ *characteristic*. Condition (1.5b) results from the presence of surface tension and the balance of the normal stresses at the interface [10]. Note that the effect of surface tension becomes especially important in modeling the flows of liquid metals [19]. Condition (1.5c) means that the fluid and vacuum magnetic fields are tangential to the interface. Conditions (1.5d) are the perfectly conducting wall and impermeable conditions. The function \mathbf{j}_c in (1.5e) represents a given surface current that forces oscillations onto the plasma-vacuum system. For laboratory plasmas this external excitation may be caused by a system

of coils. We refer to [12, section 4.6] for a thorough discussion of the condition (1.5e).

We supplement (1.1)–(1.3) and (1.5) with the initial data

$$(1.7) \quad \varphi|_{t=0} = \varphi_0 \quad \text{on } \mathbb{T}^2, \quad U|_{t=0} = U_0 \quad \text{in } \Omega^+(0),$$

where $\|\varphi_0\|_{L^\infty(\mathbb{T}^2)} < 1$. It is important to point out that if the interface function φ is given, then the vacuum magnetic field h can be uniquely determined from the elliptic problem (1.3), (1.5c), and (1.5e) [3].

In the absence of surface tension, Secchi and Trakhinin [22, 23] obtained the linear and nonlinear well-posedness for the free-boundary problem (1.1)–(1.7) with $\mathfrak{s} = 0$, provided the fluid and vacuum magnetic fields are not collinear at each point of the interface. The non-collinearity condition stems from the stability analysis of compressible current-vortex sheets (cf. [5, 26, 27]). On the one hand, this condition enhances the regularity of the moving interface $\Sigma(t)$, since it allows one to express $\partial_t \varphi$ and $\nabla \varphi$ as functions of the traces of the velocity and magnetic fields. But on the other hand, it excludes some important cases such as the case of the vanishing vacuum magnetic field. Thus in [30] we considered the free-boundary problem in ideal compressible MHD and proved the first local existence result for the case of vanishing vacuum magnetic field under the generalized Taylor sign condition. We mention that the local existence is still unsolved for the nontrivial vacuum magnetic field without the non-collinearity condition (see [29] for further discussions).

It is known that surface tension provides a stabilizing effect on the motion of free vacuum interfaces; see, for instance, Coutand and Shkoller [8] and Shatah and Zeng [24, 25] for the incompressible Euler equations with surface tension and Coutand, Hole, and Shkoller [9] for the compressible isentropic case. Motivated by these works, the authors [31] recently investigated the free-boundary ideal compressible MHD equations with surface tension and constructed the unique solution without assuming any Taylor-type sign condition. The result obtained in [31] corresponds to the special case of a vanishing vacuum magnetic field. Therefore, it is natural to examine the stabilizing effect of surface tension on the evolution of moving interfaces in ideal compressible MHD for a general vacuum magnetic field, that is, to study the local well-posedness for problem (1.1)–(1.7) with $\mathfrak{s} > 0$.

Different from the nonlinear hyperbolic problem studied in [31], the problem (1.1)–(1.7) under consideration is a nonlinear hyperbolic-elliptic coupled problem with a characteristic free boundary. To handle this, we consider very general constitutive relations satisfying the physical assumption that the sound speed is positive. The first step in our analysis is to reformulate the free-boundary problem (1.1)–(1.7) into an equivalent fixed-boundary problem by use of a simple lift of the graph.

We establish the solvability and high-order energy estimates for the linearized problem around a certain basic state by passing to the limit from a well-chosen regularization. For this purpose, we rewrite the linearized vacuum equations into a div-curl system, so that we can introduce some suitable decompositions and scalar potential ξ to obtain the reduced problem (3.32) with homogeneous boundary conditions and homogeneous vacuum equations as in [29]. Similar to our previous work [31], the L^2 estimate for the problem (3.32) is not closed. To deal with this situation, we design an elaborate ε -regularization that has a unique solution satisfying uniform-in- ε energy estimates in certain Sobolev spaces of sufficiently large regularity.

More precisely, we add some terms in the fluid equations and the boundary conditions to close the L^2 estimates for both the regularization (3.34) and its dual problem (3.50). This enables us to derive the existence of L^2 weak solutions to the

ε -regularization for any small fixed parameter $\varepsilon > 0$ by applying the duality argument. Then we build for the regularized problem (3.34) high-order energy estimates uniformly in ε . Noting that the fluid variables W satisfy a symmetric hyperbolic problem with characteristic boundary, we choose to work in the anisotropic Sobolev spaces H_*^m first introduced by Chen [6] with different regularity in the normal and tangential directions (see Secchi [20] for a general theory). In the estimate of tangential derivatives (cf. section 3.5.4), we have to deal with the troublesome boundary term, i.e., the integral of Q_{2c} (cf. (3.77)). To overcome this difficulty, we add another regularized term $-\varepsilon\Delta_{x'}^2\xi$ in the boundary condition (3.34e) so that the good term \mathcal{J}_2 (cf. (3.78)–(3.79)) can be used to estimate the integral of Q_{2c} as in (3.87), where $\Delta_{x'}^2$ denotes the biharmonic operator in the tangential space coordinates. Moreover, since the $L^2(\Sigma_t)$ -norm of $\partial_t^m\psi$ rather than its instant $L^2(\Sigma)$ -norm is controllable because of the boundary condition (3.34c), we consider the estimate for Q_4 in the cases with $\alpha_0 < m$ and $\alpha = m$ separately (see section 3.5.4 for the details). The elliptic equation for the gradient of the potential ξ helps to gain the normal derivatives of $\nabla\xi$, which along with the spatial boundary regularity enhanced from surface tension allows us to obtain the high-order energy estimates for the ε -regularization (3.34) uniformly in ε . Then we achieve the resolution and high-order energy estimate for the linearized problem (3.14) by passing to the limit $\varepsilon \rightarrow 0$.

Our energy estimate (3.20) for the linearized problem exhibits a *fixed* loss of regularity from the basic state and source terms to the solution and hence is a so-called tame estimate. Based on the linear well-posedness result, we can construct local solutions for the nonlinear problem by an appropriate modification of the Nash–Moser iteration scheme developed by Hörmander [13] and Coulombel and Secchi [7]. In particular, a smooth intermediate state should be introduced and estimated, so that the state around which we linearize at each iteration step can satisfy certain constraints for the linear solvability.

The manuscript is organized as follows. In section 2, we reformulate the nonlinear free-boundary problem and present the main result of this paper (i.e., Theorem 2.3). Section 3 is devoted to proving the energy estimates and unique solvability for the linearized problem around a suitable basic state (cf. Theorem 3.1). In section 4, we show the local existence of solutions for the nonlinear problem.

2. Main result. In this section we reformulate the nonlinear free-boundary problem (1.1)–(1.7) into an equivalent fixed-boundary problem, state the main result of this paper, and present the notation for later use.

2.1. Nonlinear problem. We consider very general, smooth constitutive relations $\rho = \rho(p, S)$ and $\mathfrak{e} = \mathfrak{e}(p, S)$ for the perfectly conducting fluid. All we need is that the sound speed $a = a(\rho, S)$ satisfies

$$(2.1) \quad a(\rho, S) := \sqrt{p_\rho(\rho, S)} > 0 \quad \text{for all } \rho \in (\rho_*, \rho^*),$$

where ρ_* and ρ^* are some nonnegative constants. Using the constraint (1.2) and the Gibbs relation $\vartheta dS = d\mathfrak{e} + p d(1/\rho)$ with $\vartheta > 0$ being the temperature, we can reduce the fluid equations (1.1) to the symmetric hyperbolic system

$$(2.2) \quad A_0^+(U)\partial_t U + \sum_{i=1}^3 A_i^+(U)\partial_i U = 0 \quad \text{in } \Omega^+(t)$$

for smooth solutions $U = (q, v, H, S)^\top$, where

$$A_0^+(U) := \begin{pmatrix} \frac{1}{\rho a^2} & 0 & -\frac{1}{\rho a^2} H^\top & 0 \\ 0 & \rho I_3 & O_3 & 0 \\ -\frac{1}{\rho a^2} H & O_3 & I_3 + \frac{1}{\rho a^2} H \otimes H & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

and

$$A_i^+(U) := \begin{pmatrix} \frac{v_i}{\rho a^2} & \mathbf{e}_i^\top & -\frac{v_i}{\rho a^2} H^\top & 0 \\ \mathbf{e}_i & \rho v_i I_3 & -H_i I_3 & 0 \\ -\frac{v_i}{\rho a^2} H & -H_i I_3 & v_i I_3 + \frac{v_i}{\rho a^2} H \otimes H & 0 \\ 0 & 0 & 0 & v_i \end{pmatrix} \quad \text{for } i = 1, 2, 3.$$

Here and below, O_m and I_m are the zero and identity matrices of order m , respectively, $\mathbf{e}_1 := (1, 0, 0)^\top$, $\mathbf{e}_2 := (0, 1, 0)^\top$, and $\mathbf{e}_3 := (0, 0, 1)^\top$. Moreover, it is convenient to rewrite the vacuum equations (1.3) as

$$(2.3) \quad \sum_{i=1}^3 A_i^- \partial_i h = 0 \quad \text{in } \Omega^-(t),$$

where A_1^- , A_2^- , and A_3^- are the constant matrices given by

$$A_1^- := \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad A_2^- := \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad A_3^- := \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Let us reformulate the free-boundary problem (2.2)–(2.3) and (1.5)–(1.7) into an equivalent fixed-boundary problem. To this end, we take the lifting function Φ as

$$\Phi(t, x) := x_1 + \chi(x_1) \varphi(t, x'),$$

where $\chi \in C_0^\infty(-1, 1)$ is a cut-off function satisfying

$$(2.4) \quad \|\chi'\|_{L^\infty(\mathbb{R})} < \frac{4}{\|\varphi_0\|_{L^\infty(\mathbb{T}^2)} + 3}, \quad \chi \equiv 1 \quad \text{on } [-\delta_0, \delta_0],$$

for some small constant $\delta_0 > 0$. Then the change of variables $(t, x) \mapsto (t, \tilde{x})$ with $x = (\Phi(t, \tilde{x}), \tilde{x}_2, \tilde{x}_3)$ maps Σ^\pm and $\Sigma(t)$ to Σ^\pm and $\Sigma := \{0\} \times \mathbb{T}^2$, respectively. The domains $\Omega^+(t)$ and $\Omega^-(t)$ are mapped to $\Omega^+ := (0, 1) \times \mathbb{T}^2$ and $\Omega^- := (-1, 0) \times \mathbb{T}^2$, respectively. Since $\|\varphi_0\|_{L^\infty(\mathbb{T}^2)} < 1$, we know that $\|\varphi\|_{L^\infty([0, T] \times \mathbb{T}^2)} \leq \frac{1}{2}(\|\varphi_0\|_{L^\infty(\mathbb{T}^2)} + 1) < 1$ for some small $T > 0$. As a result, the change of variables is admissible on the time interval $[0, T]$.

Let us introduce $\tilde{U}(t, \tilde{x}) = U(t, x)$ and $\tilde{h}(t, \tilde{x}) = h(t, x)$. Then the free-boundary problem (2.2)–(2.3), (1.5)–(1.7) can be reduced to the following nonlinear fixed-boundary problem:

$$\begin{aligned}
 (2.5a) \quad & \mathbb{L}_+(U, \Phi) := L_+(U, \Phi)U = 0 && \text{in } [0, T] \times \Omega^+, \\
 (2.5b) \quad & \mathbb{L}_-(h, \Phi) := L_-(\Phi)h = 0 && \text{in } [0, T] \times \Omega^-, \\
 (2.5c) \quad & \partial_t \varphi = v \cdot N, \quad q - \frac{1}{2}|h|^2 = \mathfrak{s}\mathcal{H}(\varphi), \quad h \cdot N = 0 && \text{on } [0, T] \times \Sigma, \\
 (2.5d) \quad & v_1 = 0 \quad \text{on } [0, T] \times \Sigma^+, \quad h \times \mathbf{e}_1 = \mathbf{j}_c && \text{on } [0, T] \times \Sigma^-, \\
 (2.5e) \quad & (U, \varphi)|_{t=0} = (U_0, \varphi_0),
 \end{aligned}$$

where we drop the tildes for notational simplicity. The operators L_{\pm} are defined as

$$(2.6) \quad L_+(U, \Phi) := A_0^+(U)\partial_t + \tilde{A}_1^+(U, \Phi)\partial_1 + A_2^+(U)\partial_2 + A_3^+(U)\partial_3,$$

$$(2.7) \quad L_-(\Phi) := \tilde{A}_1^-(\Phi)\partial_1 + A_2^-\partial_2 + A_3^-\partial_3,$$

with $\tilde{A}_1^-(\Phi) := (A_1^- - \partial_2\Phi A_2^- - \partial_3\Phi A_3^-)/\partial_1\Phi$ and

$$\tilde{A}_1^+(U, \Phi) := \frac{1}{\partial_1\Phi} (A_1^+(U) - \partial_t\Phi A_0^+(U) - \partial_2\Phi A_2^+(U) - \partial_3\Phi A_3^+(U)).$$

For later use, we introduce the following boundary operators (cf. (2.5c)–(2.5d)):

$$(2.8) \quad \mathbb{B}_+(U, h, \varphi) := \begin{pmatrix} \partial_t\varphi - v \cdot N \\ q - \frac{1}{2}|h|^2 - \mathfrak{s}\mathcal{H}(\varphi) \\ v_1 \end{pmatrix}, \quad \mathbb{B}_-(h, \varphi) := \begin{pmatrix} h \cdot N \\ h \times \mathbf{e}_1 - \mathbf{j}_c \end{pmatrix}.$$

In the new coordinates, (1.2) and the first conditions in (1.5c)–(1.5d) become

$$(2.9) \quad \frac{\partial_1 H_1}{\partial_1 \Phi} + \sum_{i=2}^3 \left(\partial_i - \frac{\partial_i \Phi}{\partial_1 \Phi} \partial_1 \right) H_i = 0 \quad \text{in } \Omega^+,$$

$$(2.10) \quad H \cdot N = 0 \quad \text{on } \Sigma, \quad H_1 = 0 \quad \text{on } \Sigma^+.$$

The identities (2.9)–(2.10) can be regarded as initial constraints, meaning that they are satisfied for $t > 0$ as long as they hold at the initial time, for which the proof can be found in [27, Appendix A].

2.2. Main result. Before stating the main theorem, we introduce the compatibility conditions on the initial data. To this end, we suppose that the initial data $U_0 \in H^{m+3/2}(\Omega^+)$ and $\varphi_0 \in H^{m+2}(\mathbb{T}^2)$ satisfy $\|\varphi_0\|_{L^\infty(\mathbb{T}^2)} < 1$ and the hyperbolicity condition

$$(2.11) \quad \rho_* < \inf_{\Omega^+} \rho(U_0) \leq \sup_{\Omega^+} \rho(U_0) < \rho^*,$$

where the nonnegative constants ρ_* and ρ^* are specified in (2.1) and $m \geq 3$ is an integer. It follows from (2.4) that

$$\partial_1 \Phi_0 \geq \frac{3 - 3\|\varphi_0\|_{L^\infty(\mathbb{T}^2)}}{3 + \|\varphi_0\|_{L^\infty(\mathbb{T}^2)}} > 0 \quad \text{for } \Phi_0(x) := x_1 + \chi(x_1)\varphi_0(x').$$

Then the initial vacuum magnetic field $h_0 \in \mathbb{R}^3$ is uniquely determined by the following div-curl system (cf. (2.5b)–(2.5d) and Lemma 3.2):

$$L_-(\Phi_0)h_0 = 0 \quad \text{in } \Omega^-, \quad h_0 \cdot N_0 = 0 \quad \text{on } \Sigma, \quad h_0 \times \mathbf{e}_1 = \mathbf{j}_c(0) \quad \text{on } \Sigma^-,$$

where the operator L_- is defined by (2.7) and $N_0 := (1, -\partial_2\varphi_0, -\partial_3\varphi_0)^\top$. Let us denote $U_{(\ell)} := \partial_t^\ell U|_{t=0}$ and $\varphi_{(\ell)} := \partial_t^\ell \varphi|_{t=0}$ for any $\ell \in \mathbb{N}$. Taking ℓ time derivatives of the interior equations (2.5a) and the first condition in (2.5c), we evaluate the resulting identities at the initial time to determine $U_{(\ell)}$ and $\varphi_{(\ell)}$ inductively. Then we set $h_{(\ell)} := \partial_t^\ell h|_{t=0}$ as the unique solution of the elliptic problem that results from taking ℓ time derivatives of the equations (2.5b), the third condition in (2.5c), and the second condition in (2.5d). More precisely, we have the following result (see [23, Lemma 19] for the detailed proof).

LEMMA 2.1. *Suppose that $m \geq 3$ is an integer, the surface current \mathbf{j}_c belongs to $H^{m+3/2}([0, T_0] \times \Sigma^-)$ for some $T_0 > 0$, and the initial data $(U_0, \varphi_0) \in H^{m+3/2}(\Omega^+) \times H^{m+2}(\mathbb{T}^2)$ satisfy $\|\varphi_0\|_{L^\infty(\mathbb{T}^2)} < 1$ and (2.11). Then the procedure described above determines $U_{(\ell)} \in H^{m+3/2-\ell}(\Omega^+)$, $\varphi_{(\ell)} \in H^{m+2-\ell}(\mathbb{T}^2)$, and $h_{(\ell)} \in H^{m+3/2-\ell}(\Omega^-)$, for $\ell = 0, 1, \dots, m$, which satisfy*

$$\sum_{\ell=0}^m (\|U_{(\ell)}\|_{H^{m+3/2-\ell}(\Omega^+)} + \|\varphi_{(\ell)}\|_{H^{m+2-\ell}(\mathbb{T}^2)} + \|h_{(\ell)}\|_{H^{m+3/2-\ell}(\Omega^-)}) \leq C(M_0),$$

for some positive constant $C(M_0)$ depending on

$$(2.12) \quad M_0 := \|U_0\|_{H^{m+3/2}(\Omega^+)} + \|\varphi_0\|_{H^{m+2}(\mathbb{T}^2)} + \|\mathbf{j}_c\|_{H^{m+3/2}([0, T_0] \times \Sigma^-)}.$$

Taking ℓ time derivatives of the second condition in (2.5c) as in [31, (3.4)] leads to the following terminology for the compatibility conditions on the initial data.

DEFINITION 2.2. *Suppose that all the conditions of Lemma 2.1 are satisfied. The initial data (U_0, φ_0) are said to satisfy the compatibility conditions up to order m if $U_{(\ell)}$, $\varphi_{(\ell)}$, and $h_{(\ell)}$ satisfy the boundary conditions $v_{1(\ell)}|_{\Sigma^+} = 0$ and*

$$(2.13) \quad q_{(\ell)} = \sum_{i=0}^{\ell-1} \binom{\ell-1}{i} h_{(i)} \cdot h_{(\ell-i)} + \mathfrak{s} \sum_{\substack{\alpha_i \in \mathbb{N}^2 \\ |\alpha_1| + \dots + |\alpha_\ell| = \ell}} D_{x'} \cdot \left(D_\zeta^{\alpha_1 + \dots + \alpha_\ell} \mathfrak{f}(\zeta_{(0)}) \ell! \prod_{i=1}^{\ell} \frac{1}{\alpha_i!} \left(\frac{\zeta_{(i)}}{i!} \right)^{\alpha_i} \right) \quad \text{on } \Sigma,$$

for $\ell = 0, \dots, m$, where $\zeta_{(i)} := D_{x'} \varphi_{(i)} \in \mathbb{R}^2$ and $\mathfrak{f}(\zeta) := \zeta / \sqrt{1 + |\zeta|^2}$.

Denote by $\lfloor s \rfloor$ the floor function of $s \in \mathbb{R}$ that maps s to the greatest integer less than or equal to s . We now state the main result of this paper, that is, the local existence theorem for the nonlinear problem (2.5).

THEOREM 2.3. *Let $\mathbf{j}_c \in H^{m+3/2}([0, T_0] \times \Sigma^-)$ for some $T_0 > 0$ and integer $m \geq 20$. Suppose that the initial data $(U_0, \varphi_0) \in H^{m+3/2}(\Omega^+) \times H^{m+2}(\mathbb{T}^2)$ satisfy $\|\varphi_0\|_{L^\infty(\mathbb{T}^2)} < 1$, the constraints (2.9)–(2.10), the hyperbolicity condition (2.11), and the compatibility conditions up to order m . Then there exists a small time $T > 0$ such that the problem (2.5) has a unique solution (U, h, φ) in $H^{\lfloor (m-9)/2 \rfloor}([0, T] \times \Omega^+) \times H^{m-9}([0, T] \times \Omega^-) \times H^{m-9}([0, T] \times \mathbb{T}^2)$ satisfying $D_{x'} \varphi \in H^{m-9}([0, T] \times \mathbb{T}^2)$.*

The solution U constructed in this paper belongs to the anisotropic Sobolev space $H_*^{m-9}([0, T] \times \Omega^+)$, and hence $U \in H^{\lfloor (m-9)/2 \rfloor}([0, T] \times \Omega^+)$ due to the embedding $H_*^m \hookrightarrow H^{\lfloor m/2 \rfloor}$. For the case of zero vacuum magnetic field and surface tension, we [30] showed the first local existence result with a regularity loss from the initial data,

while Lindblad and Zhang [15] recently obtained the a priori estimates without loss of regularity in the anisotropic Sobolev space H_*^8 . It is an interesting open problem to establish the a priori estimates and existence of solutions without loss of (anisotropic) regularity for the nonlinear free-boundary problem (1.1)–(1.7).

2.3. Notation. We adopt the following notation throughout the paper:

- (a) We write C for some universal positive constant and $C(\cdot)$ for some positive constant depending on the quantities listed in the parentheses. Symbol $A \lesssim B$ denotes $A \leq CB$, while $A \lesssim_{a_1, \dots, a_m} B$ means that $A \leq C(a_1, \dots, a_m)B$ for given parameters a_1, \dots, a_m .
- (b) Recall that $\Sigma^\pm := \{\pm 1\} \times \mathbb{T}^2$ are the boundaries of the reference domain $\Omega := (-1, 1) \times \mathbb{T}^2$ and $\Sigma := \{0\} \times \mathbb{T}^2$ is the common boundary of $\Omega^\pm := \{0 < \pm x_1 < 1\} \cap \Omega$. For $T > 0$, we set $\Omega_T := (-\infty, T) \times \Omega$, $\Sigma_T := (-\infty, T) \times \Sigma$, and

$$\Omega_T^\pm := (-\infty, T) \times \Omega^\pm, \quad \Sigma_T^\pm := (-\infty, T) \times \Sigma^\pm.$$

- (c) We denote by ∂_t (or ∂_0) the time derivative $\frac{\partial}{\partial t}$. Set $\nabla := (\partial_1, \partial_2, \partial_3)^\top$ and $D := (\partial_0, \partial_1, \partial_2, \partial_3)^\top$, where $\partial_i := \frac{\partial}{\partial x_i}$ for $i = 1, 2, 3$. For any $\beta = (\beta_1, \dots, \beta_n) \in \mathbb{N}^n$ and $z = (z_1, \dots, z_n) \in \mathbb{R}^n$, we define

$$\beta! := \beta_1! \cdots \beta_n!, \quad |\beta| := \beta_1 + \cdots + \beta_n, \quad z^\beta := z_1^{\beta_1} \cdots z_n^{\beta_n},$$

$$D_z := \left(\frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_n} \right)^\top, \quad D_z^\beta := \left(\frac{\partial}{\partial z_1} \right)^{\beta_1} \cdots \left(\frac{\partial}{\partial z_n} \right)^{\beta_n}.$$

- (d) We abbreviate $z' := (z_2, z_3)^\top$ for $z := (z_1, z_2, z_3)^\top$, so that $x' := (x_2, x_3)$. We use $D_{x'} := (\partial_2, \partial_3)^\top$, $\Delta_{x'} := D_{x'} \cdot D_{x'}$, and $\Delta_{x'}^2 := \Delta_{x'} \Delta_{x'}$ to denote the gradient, Laplacian, and biharmonic operators in the tangential space coordinates x' , respectively. For any integer $m \geq 2$, we use $D_{x'}^m := (\partial_2^m, \partial_2^{m-1} \partial_3, \dots, \partial_2 \partial_3^{m-1}, \partial_3^m)^\top$ to represent the vector of all partial derivatives in x' of order m .
- (e) We denote $D_*^\alpha := \partial_t^{\alpha_0} (\sigma \partial_1)^{\alpha_1} \partial_2^{\alpha_2} \partial_3^{\alpha_3} \partial_1^{\alpha_4}$ for $\alpha := (\alpha_0, \dots, \alpha_4) \in \mathbb{N}^5$ and $\sigma := x_1(1 - x_1)$. For $m \in \mathbb{N}$ and $I \subset \mathbb{R}$, the anisotropic Sobolev space $H_*^m(I \times \Omega^+)$ is defined as

$$H_*^m(I \times \Omega^+) := \{u \in L^2(I \times \Omega^+) : D_*^\alpha u \in L^2(I \times \Omega^+) \text{ for } \langle \alpha \rangle \leq m\},$$

and equipped with the norm $\|\cdot\|_{H_*^m(I \times \Omega^+)}$, where

$$\|u\|_{H_*^m(I \times \Omega^+)}^2 := \sum_{\langle \alpha \rangle \leq m} \|D_*^\alpha u\|_{L^2(I \times \Omega^+)}^2, \quad \langle \alpha \rangle := \sum_{i=0}^3 \alpha_i + 2\alpha_4.$$

By definition, $H^m(I \times \Omega^+) \hookrightarrow H_*^m(I \times \Omega^+) \hookrightarrow H^{\lfloor m/2 \rfloor}(I \times \Omega^+)$ for all $m \in \mathbb{N}$ and $I \subset \mathbb{R}$.

- (e) For any $m \in \mathbb{N}$, we denote by \mathring{c}_m a generic and smooth matrix-valued function of $\{(D^\alpha \mathring{U}, D^\alpha \mathring{h}, D^\alpha \mathring{\varphi}) : |\alpha| \leq m\}$, where $D^\alpha := \partial_t^{\alpha_0} \partial_1^{\alpha_1} \partial_2^{\alpha_2} \partial_3^{\alpha_3}$ for $\alpha := (\alpha_0, \dots, \alpha_3) \in \mathbb{N}^4$. The exact form of \mathring{c}_m may vary at different places.

3. Linear well-posedness. This section is devoted to showing the high-order energy estimates and unique solvability for the linearization of the problem (2.5) around a suitable basic state $(\mathring{U}, \mathring{h}, \mathring{\varphi})$.

3.1. Main theorem for the linearized problem. We assume that the basic state $(\mathring{U}(t, x), \mathring{h}(t, x), \mathring{\varphi}(t, x'))$ with $\mathring{U} = (\mathring{q}, \mathring{v}, \mathring{H}, \mathring{S})^\top$ and $\mathring{h} = (\mathring{h}_1, \mathring{h}_2, \mathring{h}_3)^\top$ is sufficiently smooth and satisfies

$$(3.1) \quad \|\mathring{\varphi}\|_{L^\infty(\Sigma_T)} \leq \frac{1}{2}(\|\varphi_0\|_{L^\infty(\mathbb{T}^2)} + 1) < 1,$$

$$(3.2) \quad \rho_* < \rho(\mathring{U}) < \rho^* \quad \text{on } \overline{\Omega_T^+},$$

$$(3.3) \quad \|\mathring{U}\|_{W^{3,\infty}(\Omega_T^+)} + \|\mathring{h}\|_{W^{3,\infty}(\Omega_T^-)} + \|\mathring{\varphi}\|_{W^{4,\infty}(\Sigma_T)} \leq K$$

for some constant $K > 0$. We also assume that

$$(3.4) \quad \partial_t \mathring{\varphi} = \mathring{v} \cdot \mathring{N}, \quad \mathring{h} \cdot \mathring{N} = 0, \quad \mathring{H} \cdot \mathring{N} = 0 \quad \text{on } \Sigma_T,$$

$$(3.5) \quad \partial_1 \mathring{h} \cdot \mathring{N} + \partial_2 \mathring{h}_2 + \partial_3 \mathring{h}_3 = 0 \quad \text{on } \Sigma_T,$$

$$(3.6) \quad \mathring{H}_1 = 0, \quad \mathring{v}_1 = 0 \quad \text{on } \Sigma_T^+, \quad \mathring{h} \times \mathbf{e}_1 = \mathbf{j}_c \quad \text{on } \Sigma_T^-,$$

where $\mathring{N} := (1, -\partial_2 \mathring{\varphi}, -\partial_3 \mathring{\varphi})^\top$. Constraint (3.5) comes from restricting the last equation in (2.5b) on boundary Σ_T for the basic state. Moreover, $\partial_1 \mathring{\Phi} > 0$ on $\overline{\Omega_T}$ for

$$\mathring{\Phi}(t, x) := x_1 + \mathring{\Psi}(t, x), \quad \mathring{\Psi}(t, x) := \chi(x_1)\mathring{\varphi}(t, x').$$

Let us introduce the good unknowns of Alinhac [1]:

$$(3.7) \quad \mathring{V} := V - \frac{\Psi}{\partial_1 \mathring{\Phi}} \partial_1 \mathring{U}, \quad \mathring{h} := h - \frac{\Psi}{\partial_1 \mathring{\Phi}} \partial_1 \mathring{h}$$

for $V = (q, v, H, S)^\top$ and $\Psi(t, x) := \chi(x_1)\psi(t, x')$. Then the linearized operators for the interior equations (2.5a)–(2.5b) around the basic state $(\mathring{U}, \mathring{h}, \mathring{\varphi})$ are defined and simplified as

$$(3.8) \quad \mathbb{L}'_+(\mathring{U}, \mathring{\Phi})(V, \Psi) := \frac{d}{d\theta} \mathbb{L}_+(\mathring{U} + \theta V, \mathring{\Phi} + \theta \Psi) \Big|_{\theta=0} \\ = L_+(\mathring{U}, \mathring{\Phi})\mathring{V} + \mathcal{C}_+(\mathring{U}, \mathring{\Phi})\mathring{V} + \frac{\Psi}{\partial_1 \mathring{\Phi}} \partial_1 \mathbb{L}_+(\mathring{U}, \mathring{\Phi}),$$

$$(3.9) \quad \mathbb{L}'_-(\mathring{h}, \mathring{\Phi})(h, \Psi) := \frac{d}{d\theta} \mathbb{L}_-(\mathring{h} + \theta h, \mathring{\Phi} + \theta \Psi) \Big|_{\theta=0} = L_-(\mathring{\Phi})\mathring{h} + \frac{\Psi}{\partial_1 \mathring{\Phi}} \partial_1 \mathbb{L}_-(\mathring{h}, \mathring{\Phi}),$$

where

$$(3.10) \quad \mathcal{C}_+(\mathring{U}, \mathring{\Phi})V := \sum_{k=1}^8 V_k \left(\frac{\partial \tilde{A}_1^+}{\partial U_k}(\mathring{U}, \mathring{\Phi}) \partial_1 U + \sum_{i=0,2,3} \frac{\partial A_i^+}{\partial U_k}(\mathring{U}) \partial_i U \right).$$

To linearize the boundary conditions (2.5c), we recall (1.6) and calculate

$$\frac{d}{d\theta} \mathcal{H}(\mathring{\varphi} + \theta \psi) \Big|_{\theta=0} = D_{x'} \cdot \frac{d}{d\theta} \left(\frac{D_{x'}(\mathring{\varphi} + \theta \psi)}{\sqrt{1 + |D_{x'}(\mathring{\varphi} + \theta \psi)|^2}} \right) \Big|_{\theta=0} = D_{x'} \cdot (\mathring{B} D_{x'} \psi),$$

where \mathring{B} is the positive definite matrix defined by

$$(3.11) \quad \mathring{B} := \frac{I_2}{|\mathring{N}|} - \frac{D_{x'} \mathring{\varphi} \otimes D_{x'} \mathring{\varphi}}{|\mathring{N}|^3}.$$

Consequently, for the boundary operators \mathbb{B}_\pm defined by (2.8), we have

$$(3.12) \quad \mathbb{B}'_+(\dot{U}, \dot{h}, \dot{\varphi})(V, h, \psi) := \frac{d}{d\theta} \mathbb{B}_+(\dot{U} + \theta V, \dot{h} + \theta h, \dot{\varphi} + \theta \psi) \Big|_{\theta=0} = \begin{pmatrix} (\partial_t + \dot{v}' \cdot D_{x'})\psi - v \cdot \dot{N} \\ q - \dot{h} \cdot h - \mathfrak{s} D_{x'} \cdot (\dot{B} D_{x'} \psi) \\ v_1 \end{pmatrix},$$

$$(3.13) \quad \mathbb{B}'_-(\dot{h}, \dot{\varphi})(h, \psi) := \frac{d}{d\theta} \mathbb{B}_-(\dot{h} + \theta h, \dot{\varphi} + \theta \psi) \Big|_{\theta=0} = \begin{pmatrix} h \cdot \dot{N} - \dot{h}' \cdot D_{x'} \psi \\ h \times \mathbf{e}_1 \end{pmatrix},$$

where we denote $z' := (z_2, z_3)^\top$ for any vector $z := (z_1, z_2, z_3)^\top$.

Apply the good unknowns (3.7) and neglect the last terms in (3.8)–(3.9) to obtain the following effective linear problem:

$$(3.14a) \quad \mathbb{L}'_{e+}(\dot{U}, \dot{\Phi})\dot{V} := L_+(\dot{U}, \dot{\Phi})\dot{V} + \mathcal{C}_+(\dot{U}, \dot{\Phi})\dot{V} = f^+ \quad \text{in } \Omega_T^+,$$

$$(3.14b) \quad L_-(\dot{\Phi})\dot{h} = f^- \quad \text{in } \Omega_T^-,$$

$$(3.14c) \quad \mathbb{B}'_{e+}(\dot{U}, \dot{h}, \dot{\varphi})(\dot{V}, \dot{h}, \psi) = g^+ \quad \text{on } \Sigma_T^2 \times \Sigma_T^+,$$

$$(3.14d) \quad \mathbb{B}'_{e-}(\dot{h}, \dot{\varphi})(\dot{h}, \psi) = g^- \quad \text{on } \Sigma_T \times \Sigma_T^-,$$

$$(3.14e) \quad (\dot{V}, \psi) = 0, \quad \dot{h} = 0 \quad \text{if } t < 0,$$

where operators L_\pm and \mathcal{C}_+ are defined by (2.6)–(2.7) and (3.10), respectively. According to the identities

$$\mathbb{B}'_{e+}(\dot{U}, \dot{h}, \dot{\varphi})(\dot{V}, \dot{h}, \psi) = \mathbb{B}'_+(\dot{U}, \dot{h}, \dot{\varphi})(V, h, \psi) \quad \text{and} \quad \mathbb{B}'_{e-}(\dot{h}, \dot{\varphi})(\dot{h}, \psi) = \mathbb{B}'_-(\dot{h}, \dot{\varphi})(h, \psi),$$

we get from definition (3.7) and constraint (3.5) that

$$(3.15) \quad \mathbb{B}'_{e+}(\dot{U}, \dot{h}, \dot{\varphi})(\dot{V}, \dot{h}, \psi) := \begin{pmatrix} (\partial_t + \dot{v}' \cdot D_{x'} + \dot{b}_1)\psi - \dot{v} \cdot \dot{N} \\ \dot{q} - \dot{h} \cdot \dot{h} - \mathfrak{s} D_{x'} \cdot (\dot{B} D_{x'} \psi) + \dot{b}_2 \psi \\ \dot{v}_1 \end{pmatrix},$$

$$(3.16) \quad \mathbb{B}'_{e-}(\dot{h}, \dot{\varphi})(\dot{h}, \psi) := \begin{pmatrix} \dot{h} \cdot \dot{N} - D_{x'} \cdot (\dot{h}' \psi) \\ \dot{h} \times \mathbf{e}_1 \end{pmatrix}$$

for $\dot{b}_1 := -\partial_1 \dot{v} \cdot \dot{N}$ and $\dot{b}_2 := \partial_1 \dot{q} - \dot{h} \cdot \partial_1 \dot{h}$. In (3.14c) we employ the notation $\Sigma_T^2 \times \Sigma_T^+$ to denote that the first two components of this vector equation are taken on Σ_T and the third component on Σ_T^+ . Similar notation applies also for (3.14d). The source terms f^\pm and g^\pm are supposed to vanish in the past, so that the second equation in (3.14e) follows from (3.14b), (3.14d), and the first equation in (3.14e).

The well-posedness result for the effective linear problem (3.14) is presented in the following theorem. Hereafter, we use the shorthand notation

$$(3.17) \quad \|(U, h, \varphi)\|_m := \|U\|_{H_*^m(\Omega_T^+)} + \|h\|_{H^m(\Omega_T^-)} + \|\varphi\|_{H^m(\Sigma_T)},$$

$$(3.18) \quad \|(g^+, g^-)\|_{H^m \times H^{m+1}} := \|(g_1^+, g_2^+)\|_{H^m(\Sigma_T)} + \|g_3^+\|_{H^m(\Sigma_T^+)} + \|g_1^-\|_{H^{m+1}(\Sigma_T)} + \|g_2^-\|_{H^{m+1}(\Sigma_T^-)}.$$

THEOREM 3.1. *Let $K_0 > 0$ and $m \in \mathbb{N}$ with $m \geq 6$. Then there exist constants $T_0 > 0$ and $C(K_0) > 0$ such that if $(\dot{U}, \dot{h}, \dot{\varphi}) \in H_*^{m+4}(\Omega_T^+) \times H^{m+4}(\Omega_T^-) \times H^{m+4}(\Sigma_T)$ satisfies (3.1)–(3.6) and*

$$(3.19) \quad \|\dot{U}\|_{H_*^{10}(\Omega_T^+)} + \|\dot{h}\|_{H^{10}(\Omega_T^-)} + \|\dot{\varphi}\|_{H^{10}(\Sigma_T)} \leq K_0,$$

and source terms $(f^+, f^-) \in H_*^m(\Omega_T^+) \times H^{m+1}(\Omega_T^-)$, $g^+ \in H^{m+1}(\Sigma_T)^2 \times H^{m+1}(\Sigma_T^+)$, and $g^- \in H^{m+2}(\Sigma_T) \times H^{m+2}(\Sigma_T^-)$ vanish in the past and satisfy the compatibility conditions (3.25) for some $0 < T \leq T_0$, then problem (3.14) admits a unique solution $(\dot{V}, \dot{h}, \dot{\psi}) \in H_*^m(\Omega_T^+) \times H^m(\Omega_T^-) \times H^m(\Sigma_T)$ satisfying

$$(3.20) \quad \begin{aligned} & \|\dot{V}\|_{H_*^m(\Omega_T^+)} + \|\dot{h}\|_{H^m(\Omega_T^-)} + \|\Psi\|_{H^m(\Omega_T)} + \|(\psi, D_{x'}\psi)\|_{H^m(\Sigma_T)} \\ & \lesssim_{K_0} \|(\dot{U}, \dot{h}, \dot{\varphi})\|_{m+4} \left(\|f^+\|_{H_*^6(\Omega_T^+)} + \|f^-\|_{H^7(\Omega_T^-)} + \|(g^+, g^-)\|_{H^7 \times H^8} \right) \\ & + \|f^+\|_{H_*^m(\Omega_T^+)} + \|f^-\|_{H^{m+1}(\Omega_T^-)} + \|(g^+, g^-)\|_{H^{m+1} \times H^{m+2}}. \end{aligned}$$

The rest of this section is devoted to proving the above theorem.

3.2. Reformulation. To reduce the problem (3.14), we compute that the vacuum equations (3.14b) are equivalent to

$$(3.21) \quad \begin{pmatrix} \nabla \times (\partial_1 \dot{\Phi} \dot{\eta}^{-T} \dot{h}) \\ \nabla \cdot (\dot{\eta} \dot{h}) \end{pmatrix} = \begin{pmatrix} \dot{\eta} & 0 \\ 0 & \partial_1 \dot{\Phi} \end{pmatrix} f^- =: \tilde{f}^- \quad \text{in } \Omega_T^-,$$

where $\dot{\eta}$ is the invertible matrix defined by

$$(3.22) \quad \dot{\eta} := \begin{pmatrix} 1 & -\partial_2 \dot{\Phi} & -\partial_3 \dot{\Phi} \\ 0 & \partial_1 \dot{\Phi} & 0 \\ 0 & 0 & \partial_1 \dot{\Phi} \end{pmatrix}.$$

Then we decompose \dot{h} as $\dot{h} = h_b + h_{\natural}$, where h_{\natural} is required to solve the following div-curl boundary value problem (cf. (3.21), (3.14d), and (3.16)):

$$(3.23) \quad \begin{cases} \begin{pmatrix} \nabla \times (\partial_1 \dot{\Phi} \dot{\eta}^{-T} h_{\natural}) \\ \nabla \cdot (\dot{\eta} h_{\natural}) \end{pmatrix} = \tilde{f}^- & \text{in } \Omega_T^-, \\ h_{\natural} \cdot \dot{N} = g_1^- & \text{on } \Sigma_T, \quad h_{\natural} \times \mathbf{e}_1 = g_2^- & \text{on } \Sigma_T^-, \\ (x_2, x_3) \rightarrow h_{\natural}(t, x_1, x_2, x_3) & \text{is 1-periodic.} \end{cases}$$

The resolution and H^2 -estimate for (3.23) have been proved in Secchi and Trakhinin [23, section 6]. To get the H^m -estimate for $m \geq 6$, as in the proof of [23, Theorem 13], we decompose $\partial_1 \dot{\Phi} \dot{\eta}^{-T} h_{\natural} = \zeta^{\natural} + \nabla \xi_{\natural}$ for the vector ζ^{\natural} and the scalar ξ_{\natural} satisfying

$$\begin{cases} \nabla \times \zeta^{\natural} = (\tilde{f}_1^-, \tilde{f}_2^-, \tilde{f}_3^-)^T, \quad \nabla \cdot \zeta^{\natural} = 0 & \text{in } \Omega_T^-, \\ \zeta_1^{\natural} = 0 & \text{on } \Sigma_T, \quad \zeta^{\natural} \times \mathbf{e}_1 = g_2^- & \text{on } \Sigma_T^-, \\ (x_2, x_3) \rightarrow \zeta^{\natural}(t, x_1, x_2, x_3) & \text{is 1-periodic,} \\ \nabla \cdot (\dot{A} \nabla \xi_{\natural}) = \tilde{f}_4^- - \nabla \cdot (\dot{A} \zeta^{\natural}) & \text{in } \Omega_T^-, \\ (\dot{A} \nabla \xi_{\natural})_1 = g_1^- - (\dot{A} \zeta^{\natural})_1 & \text{on } \Sigma_T, \quad \xi_{\natural} = 0 & \text{on } \Sigma_T^-, \\ (x_2, x_3) \rightarrow \xi_{\natural}(t, x_1, x_2, x_3) & \text{is 1-periodic,} \end{cases}$$

where $(\mathring{A}\nabla\xi_{\mathfrak{h}})_1$ denotes the first component of vector $\mathring{A}\nabla\xi_{\mathfrak{h}}$ and \mathring{A} is the positive definite matrix defined by

$$(3.24) \quad \mathring{A} := \frac{1}{\partial_1 \mathring{\Phi}} \mathring{\eta} \mathring{\eta}^\top.$$

Then we can obtain the next lemma by using the elliptic regularization and the Moser-type calculus inequalities.

LEMMA 3.2. *Let $(\tilde{f}^-, g_1^-, g_2^-)$ belong to $H^{m-1}(\Omega_T^-) \times H^{m-1/2}(\Sigma_T) \times H^{m-1/2}(\Sigma_T^-)$ for some integer $m \geq 6$. Assume that the compatibility conditions*

$$(3.25) \quad g_2^- \cdot \mathbf{e}_1|_{\Sigma^-} = 0, \quad \int_{\Sigma^-} u \cdot g_2^- = \int_{\Omega^-} u \cdot (\tilde{f}_1^-, \tilde{f}_2^-, \tilde{f}_3^-)^\top$$

hold for all vectors $u \in H^1(\Omega^-)$ satisfying $u_2 = u_3 = 0$ on Σ and $\nabla \times u = 0$ in Ω^- . Then the problem (3.23) has a unique solution $h_{\mathfrak{h}}$ in $H^m(\Omega_T^-)$ and

$$(3.26) \quad \begin{aligned} \|h_{\mathfrak{h}}\|_{H^m(\Omega_T^-)} \lesssim & K (1 + \|\mathring{\varphi}\|_{H^{m+1}(\Sigma_T)}) \|(\tilde{f}^-, g_1^-, g_2^-)\|_{H^5(\Omega_T^-) \times H^5(\Sigma_T) \times H^5(\Sigma_T^-)} \\ & + \|\tilde{f}^-\|_{H^{m-1}(\Omega_T^-)} + \|g_1^-\|_{H^{m-1/2}(\Sigma_T)} + \|g_2^-\|_{H^{m-1/2}(\Sigma_T^-)}. \end{aligned}$$

To transform (3.14) into a problem with homogeneous boundary conditions, we introduce the decomposition $\mathring{V} = V_b + V_{\mathfrak{h}}$ with $V_{\mathfrak{h}} := (q_{\mathfrak{h}}, v_{\mathfrak{h}}^{\mathfrak{h}}, 0)^\top \in \mathbb{R}^8$ satisfying

$$(3.27) \quad q_{\mathfrak{h}} := \mathfrak{R}_T(g_2^+ + \mathring{h} \cdot h_{\mathfrak{h}}), \quad v_{\mathfrak{h}}^{\mathfrak{h}} := \chi(x_1)\mathfrak{R}_T(-g_1^+) + \chi(1-x_1)\tilde{\mathfrak{R}}_T g_3^+,$$

where $\mathfrak{R}_T : H^m(\Sigma_T) \rightarrow H_*^{m+1}(\Omega_T^+)$ and $\tilde{\mathfrak{R}}_T : H^m(\Sigma_T^+) \rightarrow H_*^{m+1}(\Omega_T^+)$ are the continuous extension operators (see [18] for more details). It follows from (3.14), (3.23), and (3.27) that vectors V_b and h_b solve the problem

$$\begin{aligned} (3.28a) \quad & \mathbb{L}'_{e+}(\mathring{U}, \mathring{\Phi})V = \tilde{f}^+ := f^+ - \mathbb{L}'_{e+}(\mathring{U}, \mathring{\Phi})V_{\mathfrak{h}} && \text{in } \Omega_T^+, \\ (3.28b) \quad & \nabla \times (\partial_1 \mathring{\Phi} \mathring{\eta}^{-\top} h) = 0, \quad \nabla \cdot (\mathring{\eta} h) = 0 && \text{in } \Omega_T^-, \\ (3.28c) \quad & \mathbb{B}'_{e+}(\mathring{U}, \mathring{h}, \mathring{\varphi})(V, h, \psi) = 0 && \text{on } \Sigma_T^2 \times \Sigma_T^+, \\ (3.28d) \quad & \mathbb{B}'_{e-}(\mathring{h}, \mathring{\varphi})(h, \psi) = 0 && \text{on } \Sigma_T \times \Sigma_T^-, \\ (3.28e) \quad & (V, h, \psi) = 0 && \text{if } t < 0, \end{aligned}$$

where we drop the subscript “b” for simplicity of notation and operators $\mathbb{B}'_{e\pm}$ are given in (3.15)–(3.16).

Let us derive a suitable reformulation for problem (3.28). In view of (3.28b), we can introduce the scalar potential ξ by

$$(3.29) \quad \nabla \xi = \partial_1 \mathring{\Phi} \mathring{\eta}^{-\top} h \quad \text{in } \Omega_T^-.$$

Then $\nabla \cdot (\mathring{A}\nabla\xi) = \nabla \cdot (\mathring{\eta}h) = 0$ in Ω_T^- , where \mathring{A} is the positive definite matrix defined by (3.24). To rewrite (3.28c)–(3.28d) in terms of $\nabla\xi$, we get from (3.29), (3.22), and the second condition in (3.4) that

$$(3.30) \quad \mathring{h} \cdot h = \mathring{\eta} \mathring{h} \cdot \nabla \xi = \mathring{h}' \cdot D_{x'} \xi, \quad h \cdot \mathring{N} = (\mathring{A}\nabla\xi)_1 \quad \text{on } \Sigma_T.$$

Moreover, as in [31, 30], we set

$$\begin{aligned}
 (3.31) \quad W &:= (q, v_1 - \partial_2 \dot{\Phi} v_2 - \partial_3 \dot{\Phi} v_3, v_2, v_3, H, S)^\top \\
 &= J(\dot{\Phi})^{-1} V \quad \text{with} \quad J(\dot{\Phi}) := \begin{pmatrix} 1 & 0 & 0 & 0 & & \\ 0 & 1 & \partial_2 \dot{\Phi} & \partial_3 \dot{\Phi} & & \\ 0 & 0 & 1 & 0 & & \\ 0 & 0 & 0 & 1 & & \\ & & & & O_4 & \\ & & & & & I_4 \end{pmatrix}.
 \end{aligned}$$

In view of (3.28)–(3.31), it suffices to study the reduced problem

$$\begin{aligned}
 (3.32a) \quad \mathbf{L}W &:= \sum_{i=0}^3 \mathbf{A}_i \partial_i W + \mathbf{A}_4 W = \mathbf{f} := J(\dot{\Phi})^\top \tilde{f}^+ && \text{in } \Omega_T^+, \\
 (3.32b) \quad \nabla \cdot (\dot{A} \nabla \xi) &= 0 && \text{in } \Omega_T^-, \\
 (3.32c) \quad W_2 &= (\partial_t + \dot{v}' \cdot D_{x'} + \dot{b}_1) \psi && \text{on } \Sigma_T, \\
 (3.32d) \quad W_1 &= \dot{h}' \cdot D_{x'} \xi + \mathfrak{s} D_{x'} \cdot (\dot{B} D_{x'} \psi) - \dot{b}_2 \psi && \text{on } \Sigma_T, \\
 (3.32e) \quad (\dot{A} \nabla \xi)_1 &= D_{x'} \cdot (\dot{h}' \psi) && \text{on } \Sigma_T, \\
 (3.32f) \quad W_2 &= 0 \quad \text{on } \Sigma_T^+, \quad \xi = 0 \quad \text{on } \Sigma_T^-, \quad (W, \xi, \psi)|_{t < 0} = 0,
 \end{aligned}$$

where $\mathbf{A}_i := J(\dot{\Phi})^\top \mathbf{A}_i^+(\dot{U}) J(\dot{\Phi})$ for $i = 0, 2, 3$, $\mathbf{A}_1 := J(\dot{\Phi})^\top \tilde{\mathbf{A}}_1^+(\dot{U}, \dot{\Phi}) J(\dot{\Phi})$, and $\mathbf{A}_4 := J(\dot{\Phi})^\top \mathbb{L}'_{e^+}(\dot{U}, \dot{\Phi}) J(\dot{\Phi})$. We deduce from (3.4) and (3.6) that

$$(3.33) \quad \mathbf{A}_1|_\Sigma = \mathbf{A}_1|_{\Sigma^+} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & O_6 \end{pmatrix} =: \mathbf{A}_1^{(1)}.$$

Define $\mathbf{A}_1^{(0)} := \mathbf{A} - \mathbf{A}_1^{(1)}$ so that $\mathbf{A}_1^{(0)}|_\Sigma = \mathbf{A}_1^{(0)}|_{\Sigma^+} = 0$.

3.3. L^2 estimate for the regularization. To solve problem (3.32), we introduce the following ε -regularization:

$$\begin{aligned}
 (3.34a) \quad \mathbf{L}_\varepsilon W &:= \sum_{i=0}^3 \mathbf{A}_i \partial_i W - \varepsilon \mathbf{J} \partial_1 W + \mathbf{A}_4 W = \mathbf{f} && \text{in } \Omega_T^+, \\
 (3.34b) \quad \nabla \cdot (\dot{A} \nabla \xi) &= 0 && \text{in } \Omega_T^-, \\
 (3.34c) \quad W_2 &= (\partial_t + \dot{v}' \cdot D_{x'} + \dot{b}_1) \psi + \varepsilon \Delta_{x'}^2 \psi && \text{on } \Sigma_T, \\
 (3.34d) \quad W_1 &= \dot{h}' \cdot D_{x'} \xi + \mathfrak{s} D_{x'} \cdot (\dot{B} D_{x'} \psi) - \dot{b}_2 \psi && \text{on } \Sigma_T, \\
 (3.34e) \quad (\dot{A} \nabla \xi)_1 &= D_{x'} \cdot (\dot{h}' \psi) + \varepsilon \Delta_{x'} \xi - \varepsilon \Delta_{x'}^2 \xi && \text{on } \Sigma_T, \\
 (3.34f) \quad W_2 &= 0 \quad \text{on } \Sigma_T^+, \quad \xi = 0 \quad \text{on } \Sigma_T^-, \quad (W, \xi, \psi)|_{t < 0} = 0,
 \end{aligned}$$

where $\mathbf{J} := \text{diag}(0, 1, 0, 0, 0, 0, 0)$, the matrices \dot{A} and \dot{B} are defined in (3.24) and (3.11), respectively, $\Delta_{x'} := D_{x'} \cdot D_{x'}$, and $\Delta_{x'}^2 := \Delta_{x'} \Delta_{x'}$. As in [31, section 2.3] for the problem with vanishing vacuum magnetic field, we add the term $-\varepsilon \mathbf{J} \partial_1 W$ in (3.34a) to derive the L^2 estimate for regularization (3.34). The terms $\varepsilon \Delta_{x'}^2 \psi$ and $\varepsilon \Delta_{x'} \xi$ contained respectively in (3.34c) and (3.34e) allow us to obtain the L^2 estimate for the dual problem of (3.34). Moreover, the term $-\varepsilon \Delta_{x'}^2 \xi$ will become especially useful in establishing the uniform-in- ε energy estimates for the problem (3.34).

Let us show the L^2 energy estimate for regularization (3.34). Taking the scalar product of (3.34a) with W , we utilize (3.33) and (3.34f) to obtain

$$(3.35) \quad \int_{\Omega^+} \mathbf{A}_0 W \cdot W + \varepsilon \|W_2\|_{L^2(\Sigma_t)}^2 - 2 \int_{\Sigma_t} W_1 W_2 \lesssim_K \|(\mathbf{f}, W)\|_{L^2(\Omega_t^+)}^2.$$

It follows from the boundary conditions (3.34c)–(3.34d) that

$$(3.36) \quad \begin{aligned} 2W_1 W_2 = & \underbrace{2\mathring{h}' \cdot D_{x'} \xi \partial_t \psi}_{\mathcal{T}_1} + \underbrace{2\mathring{h}' \cdot D_{x'} \xi (\mathring{v}' \cdot D_{x'} + \mathring{b}_1) \psi}_{\mathcal{T}_2} + \underbrace{2\varepsilon W_1 \Delta_{x'}^2 \psi}_{\mathcal{T}_3} \\ & + 2 \underbrace{\{\mathring{s} D_{x'} \cdot (\mathring{B} D_{x'} \psi) - \mathring{b}_2 \psi\}}_{\mathcal{T}_4} (\partial_t + \mathring{v}' \cdot D_{x'} + \mathring{b}_1) \psi \quad \text{on } \Sigma_t. \end{aligned}$$

Using $\mathcal{T}_1 = 2D_{x'} \cdot (\xi \partial_t(\mathring{h}' \psi)) - 2\xi D_{x'} \cdot \partial_t(\mathring{h}' \psi) - 2D_{x'} \xi \cdot \partial_t \mathring{h}' \psi$ and (3.34e) yields

$$(3.37) \quad - \int_{\Sigma_t} \mathcal{T}_1 = 2 \int_{\Sigma_t} \xi \partial_t (\mathring{A} \nabla \xi)_1 + \varepsilon \int_{\Sigma_t} (|D_{x'} \xi|^2 + |\Delta_{x'} \xi|^2) + 2 \int_{\Sigma_t} \partial_t \mathring{h}' \psi \cdot D_{x'} \xi.$$

Passing in the first term on the right-hand side of (3.37) to the volume integral, we utilize (3.34f) and (3.34b) to infer

$$(3.38) \quad \begin{aligned} 2 \int_{\Sigma_t} \xi \partial_t (\mathring{A} \nabla \xi)_1 &= 2 \int_{\Omega_t^-} \partial_t (\xi \partial_t (\mathring{A} \nabla \xi)_1) = 2 \int_{\Omega_t^-} \nabla \cdot (\xi \partial_t (\mathring{A} \nabla \xi)) \\ &= 2 \int_{\Omega_t^-} \nabla \xi \cdot \partial_t (\mathring{A} \nabla \xi) = \int_{\Omega_t^-} \mathring{A} \nabla \xi \cdot \nabla \xi + \int_{\Omega_t^-} \partial_t \mathring{A} \nabla \xi \cdot \nabla \xi, \end{aligned}$$

where the last identity results from the fact that the matrix \mathring{A} is symmetric (cf. (3.24)). The second term on the right-hand side of (3.37) is a good term, while the last term will be estimated below.

Regarding the term \mathcal{T}_2 defined in (3.36), we compute

$$(3.39) \quad \begin{aligned} \mathring{h}' \cdot D_{x'} \xi (\mathring{v}' \cdot D_{x'}) \psi &= \mathring{v}' \cdot D_{x'} \xi (\mathring{h}' \cdot D_{x'}) \psi - (\mathring{h}_2 \mathring{v}_3 - \mathring{h}_3 \mathring{v}_2) (\partial_2 (\psi \partial_3 \xi) - \partial_3 (\psi \partial_2 \xi)) \\ &= \mathring{v}' \cdot D_{x'} \xi (\mathring{h}' \cdot D_{x'}) \psi + D_{x'} \cdot (\mathring{c}_0 \psi D_{x'} \xi) + \mathring{c}_1 \psi D_{x'} \xi, \end{aligned}$$

where for any $m \in \mathbb{N}$ we denote by \mathring{c}_m a generic and smooth matrix-valued function of $\{(D^\alpha \mathring{U}, D^\alpha \mathring{h}, D^\alpha \mathring{\Psi}) : |\alpha| \leq m\}$. Using (3.39) and (3.34e) leads to

$$(3.40) \quad \begin{aligned} \int_{\Sigma_t} \mathcal{T}_2 &= 2 \int_{\Sigma_t} \left\{ \mathring{v}' \cdot D_{x'} \xi (D_{x'} \cdot (\mathring{h}' \psi) - D_{x'} \cdot \mathring{h}' \psi) + \mathring{c}_1 \psi D_{x'} \xi \right\} \\ &= 2 \int_{\Sigma_t} \left\{ \mathring{v}' \cdot D_{x'} \xi ((\mathring{A} \nabla \xi)_1 - \varepsilon \Delta_{x'} \xi + \varepsilon \Delta_{x'}^2 \xi) + \mathring{c}_1 \psi D_{x'} \xi \right\}. \end{aligned}$$

Let us make the estimate for each term in (3.40). Similar to (3.38), we discover

$$(3.41) \quad 2 \int_{\Sigma_t} \mathring{v}' \cdot D_{x'} \xi (\mathring{A} \nabla \xi)_1 = \sum_{i=2,3} \int_{\Omega_t^-} \left(2 \nabla \mathring{v}_i \partial_i \xi \cdot (\mathring{A} \nabla \xi) - \partial_i (\mathring{v}_i \mathring{A}) \nabla \xi \cdot \nabla \xi \right).$$

Since

$$2 \int_{\Sigma_t} \mathring{v}' \cdot D_{x'} \xi \Delta_{x'}^2 \xi = \int_{\Sigma_t} (2[\Delta_{x'}, \mathring{v}' \cdot D_{x'}] \xi \Delta_{x'} \xi - D_{x'} \cdot \mathring{v}' |\Delta_{x'} \xi|^2),$$

we have

$$(3.42) \quad 2\varepsilon \int_{\Sigma_t} \check{\nu}' \cdot D_{x'} \xi (\Delta_{x'}^2 \xi - \Delta_{x'} \xi) \lesssim_K \varepsilon \| (D_{x'} \xi, \Delta_{x'} \xi) \|_{L^2(\Sigma_t)}^2,$$

$$(3.43) \quad \int_{\Sigma_t} \check{c}_1 \psi D_{x'} \xi = \int_{\Sigma_t} (\check{c}_2 \psi + \check{c}_1 D_{x'} \psi) \xi \lesssim_K \| (\psi, D_{x'} \psi) \|_{L^2(\Sigma_t)}^2 + \| \xi \|_{L^2(\Sigma_t)}^2.$$

For the last term in (3.43), we employ integration by parts and Poincaré’s inequality (see, e.g., Evans [11, section 5.8.1]) to get

$$(3.44) \quad \| \xi \|_{L^2(\Sigma)}^2 \lesssim \| (\xi, \partial_1 \xi) \|_{L^2(\Omega^-)}^2 \lesssim \| \nabla \xi \|_{L^2(\Omega^-)}^2, \quad \| \xi \|_{L^2(\Sigma_t)}^2 \lesssim \| \nabla \xi \|_{L^2(\Omega_t^-)}^2.$$

Utilizing the boundary condition (3.34d) yields

$$\begin{aligned} - \int_{\Sigma_t} \mathcal{T}_3 &= 2\varepsilon \mathfrak{s} \int_{\Sigma_t} \Delta_{x'} (\mathring{B} D_{x'} \psi) \cdot \Delta_{x'} D_{x'} \psi \\ &\quad + 2\varepsilon \int_{\Sigma_t} \Delta_{x'} D_{x'} \psi \cdot D_{x'} (\mathring{h}' \cdot D_{x'} \xi) + 2\varepsilon \int_{\Sigma_t} \Delta_{x'} \psi \Delta_{x'} (\mathring{b}_2 \psi). \end{aligned}$$

Recalling definition (3.11) for matrix \mathring{B} , we obtain

$$(3.45) \quad \int_{\Sigma_t} \mathcal{T}_3 \leq -\varepsilon \mathfrak{s} \int_{\Sigma_t} \frac{|\Delta_{x'} D_{x'} \psi|^2}{|\mathring{N}|^3} + \varepsilon C(K) \sum_{|\alpha| \leq 2} \| (D_{x'}^\alpha \psi, D_{x'} \xi, D_{x'}^2 \xi) \|_{L^2(\Sigma_t)}^2,$$

where $D_{x'}^m := (\partial_2^m, \partial_2^{m-1} \partial_3, \dots, \partial_2 \partial_3^{m-1}, \partial_3^m)^\top$ denotes the vector of all partial derivatives in x' of order $m \geq 2$.

A lengthy calculation implies (cf. [31, (2.20)])

$$(3.46) \quad \int_{\Sigma_t} \mathcal{T}_4 \leq -\mathfrak{s} \int_{\Sigma} \frac{|D_{x'} \psi|^2}{|\mathring{N}|^3} - \int_{\Sigma} \mathring{b}_2 \psi^2 + C(K) \| (\psi, D_{x'} \psi) \|_{L^2(\Sigma_t)}^2.$$

Plugging (3.37)–(3.38) and (3.40)–(3.46) into (3.35)–(3.36) and using the identity $\| D_{x'}^2 \xi \|_{L^2(\Sigma)} \leq \| \Delta_{x'} \xi \|_{L^2(\Sigma)}$ imply

$$\begin{aligned} &\| W(t) \|_{L^2(\Omega^+)}^2 + \| D_{x'} \psi(t) \|_{L^2(\Sigma)}^2 + \| (\xi, \nabla \xi)(t) \|_{L^2(\Omega^-)}^2 \\ &\quad + \varepsilon \| (W_2, D_{x'}^3 \psi) \|_{L^2(\Sigma_t)}^2 + \varepsilon \| (D_{x'} \xi, D_{x'}^2 \xi)(t) \|_{L^2(\Sigma)}^2 \\ &\lesssim_K \| (\mathbf{f}, W) \|_{L^2(\Omega_t^+)}^2 + \| (\psi, D_{x'} \psi) \|_{L^2(\Sigma_t)}^2 + \| \nabla \xi \|_{L^2(\Omega_t^-)}^2 \\ (3.47) \quad &\quad + \varepsilon \| (D_{x'}^2 \psi, D_{x'} \xi, D_{x'}^2 \xi) \|_{L^2(\Sigma_t)}^2 + \| \psi(t) \|_{L^2(\Sigma)}^2. \end{aligned}$$

We emphasize that the L^2 estimate (3.47) is valid also for the case $\varepsilon = 0$, that is, for the linear problem (3.32).

To control the last term in (3.47), we multiply (3.34c) with ψ to infer

$$(3.48) \quad \| \psi(t) \|_{L^2(\Sigma)}^2 + 2\varepsilon \| \Delta_{x'} \psi \|_{L^2(\Sigma_t)}^2 \leq \epsilon \varepsilon \| W_2 \|_{L^2(\Sigma_t)}^2 + C(K, \epsilon \varepsilon) \| \psi \|_{L^2(\Sigma_t)}^2$$

for all $\epsilon > 0$. Combining (3.47) with (3.48), taking $\epsilon > 0$ small enough, and employing Grönwall’s inequality, we have

$$\begin{aligned} (3.49) \quad &\| W(t) \|_{L^2(\Omega^+)}^2 + \| (\xi, \nabla \xi)(t) \|_{L^2(\Omega^-)}^2 + \| (\psi, D_{x'} \psi, D_{x'} \xi, D_{x'}^2 \xi)(t) \|_{L^2(\Sigma)}^2 \\ &\quad + \| (W_2, W_1, D_{x'}^2 \psi, D_{x'}^3 \psi) \|_{L^2(\Sigma_t)}^2 \lesssim_{K, \varepsilon} \| \mathbf{f} \|_{L^2(\Omega_t^+)}^2. \end{aligned}$$

This is the desired ε -dependent L^2 energy estimate for regularization (3.34).

3.4. Existence for the regularization. We prove the existence of solutions to the regularization (3.34) by applying the duality argument. For this purpose, we introduce the dual problem of (3.34), which reads as

$$(3.50a) \quad \mathbf{L}_\varepsilon^* W^* := \left(- \sum_{i=0}^3 \mathbf{A}_i \partial_i + \varepsilon \mathbf{J} \partial_1 + \mathbf{A}_4^\top - \sum_{i=0}^3 \partial_i \mathbf{A}_i \right) W^* = \mathbf{f}^* \quad \text{in } \Omega^+,$$

$$(3.50b) \quad \nabla \cdot (\mathring{A} \nabla \xi^*) = 0 \quad \text{in } \Omega^-,$$

$$(3.50c) \quad \begin{aligned} & \partial_t w^* + \mathbf{D}_{x'} \cdot (\mathring{v}' w^*) - \varepsilon \Delta_{x'}^2 w^* - \mathring{b}_1 w^* \\ & - \mathring{h}' \cdot \mathbf{D}_{x'} \xi^* + \mathring{b}_2 W_2^* - \mathfrak{s} \mathbf{D}_{x'} \cdot (\mathring{B} \mathbf{D}_{x'} W_2^*) = 0 \end{aligned} \quad \text{on } \Sigma,$$

$$(3.50d) \quad (\mathring{A} \nabla \xi^*)_1 = \mathbf{D}_{x'} \cdot (\mathring{h}' W_2^*) + \varepsilon \Delta_{x'} \xi^* - \varepsilon \Delta_{x'}^2 \xi^* \quad \text{on } \Sigma,$$

$$(3.50e) \quad W_2^* = 0 \quad \text{on } \Sigma^+, \quad \xi^* = 0 \quad \text{on } \Sigma^-, \quad (W^*, \xi^*)|_{t>T} = 0,$$

with $w^* := W_1^* - \varepsilon W_2^*$. The conditions (3.50c)–(3.50e) are imposed to ensure that

$$\begin{aligned} & \int_{\Omega_T^+} (\mathbf{L}_\varepsilon W \cdot W^* - W \cdot \mathbf{L}_\varepsilon^* W^*) + \int_{\Omega_T^-} (\xi^* \nabla \cdot (\mathring{A} \nabla \xi) - \xi \nabla \cdot (\mathring{A} \nabla \xi^*)) \\ & = \int_{\Sigma_T^+} W_1 W_2^* - \int_{\Sigma_T} (W_2 w^* + W_1 W_2^* - \xi^* (\mathring{A} \nabla \xi)_1 + \xi (\mathring{A} \nabla \xi^*)_1) - \int_{\Sigma_T^-} \xi^* \partial_1 \xi = 0, \end{aligned}$$

where we have used (3.34c)–(3.34f). Passing then to the back time $\tilde{t} := T - t$, we find that $\tilde{W}^*(\tilde{t}, x) := W^*(t, x)$ and $\tilde{\xi}^*(\tilde{t}, x) := \xi^*(t, x)$ satisfy

$$(3.51a) \quad \left(\mathbf{A}_0 \partial_t - \sum_{i=1}^3 \mathbf{A}_i \partial_i + \varepsilon \mathbf{J} \partial_1 + \mathbf{A}_4^\top - \sum_{i=0}^3 \partial_i \mathbf{A}_i \right) W^* = \mathbf{f}^* \quad \text{in } \Omega^+,$$

$$(3.51b) \quad \nabla \cdot (\mathring{A} \nabla \xi^*) = 0 \quad \text{in } \Omega^-,$$

$$(3.51c) \quad \begin{aligned} & \partial_t w^* - \mathbf{D}_{x'} \cdot (\mathring{v}' w^*) + \varepsilon \Delta_{x'}^2 w^* + \mathring{b}_1 w^* \\ & + \mathring{h}' \cdot \mathbf{D}_{x'} \xi^* - \mathring{b}_2 W_2^* + \mathfrak{s} \mathbf{D}_{x'} \cdot (\mathring{B} \mathbf{D}_{x'} W_2^*) = 0 \end{aligned} \quad \text{on } \Sigma,$$

$$(3.51d) \quad (\mathring{A} \nabla \xi^*)_1 = \mathbf{D}_{x'} \cdot (\mathring{h}' W_2^*) + \varepsilon \Delta_{x'} \xi^* - \varepsilon \Delta_{x'}^2 \xi^* \quad \text{on } \Sigma,$$

$$(3.51e) \quad W_2^* = 0 \quad \text{on } \Sigma^+, \quad \xi^* = 0 \quad \text{on } \Sigma^-, \quad (W^*, \xi^*)|_{t<0} = 0,$$

where for convenience we have dropped the tildes. Taking the scalar product of (3.51a) with W^* and recalling $w^* := W_1^* - \varepsilon W_2^*$, we use (3.33) and (3.51e) to get

$$(3.52) \quad \int_{\Omega^+} \mathbf{A}_0 W^* \cdot W^* + \int_{\Sigma_t} (\varepsilon |W_2^*|^2 + 2w^* W_2^*) \lesssim_K \|(\mathbf{f}^*, W^*)\|_{L^2(\Omega_t^+)}^2.$$

It follows from (3.51b) and (3.51d)–(3.51e) that

$$\begin{aligned} \int_{\Omega_t^-} \mathring{A} \nabla \xi^* \cdot \nabla \xi^* &= \int_{\Omega_t^-} \nabla \cdot (\xi^* \mathring{A} \nabla \xi^*) = \int_{\Sigma_t} \xi^* (\mathring{A} \nabla \xi^*)_1 \\ &= - \int_{\Sigma_t} W_2^* \mathring{h}' \cdot \mathbf{D}_{x'} \xi^* - \varepsilon \int_{\Sigma_t} (|\mathbf{D}_{x'} \xi^*|^2 + |\Delta_{x'} \xi^*|^2), \end{aligned}$$

from which we have

$$(3.53) \quad \|\nabla \xi^*\|_{L^2(\Omega_t^-)}^2 + \|(\mathbf{D}_{x'} \xi^*, \mathbf{D}_{x'}^2 \xi^*)\|_{L^2(\Sigma_t)}^2 \lesssim_{K, \varepsilon} \|W_2^*\|_{L^2(\Sigma_t)}^2.$$

Multiplying the boundary condition (3.51c) by w^* leads to

$$\begin{aligned} & \|w^*(t)\|_{L^2(\Sigma)}^2 + 2\varepsilon \|\Delta_{x'} w^*\|_{L^2(\Sigma_t)}^2 \\ & \leq \varepsilon \varepsilon \|(D_{x'} w^*, D_{x'}^2 w^*)\|_{L^2(\Sigma_t)}^2 + C(K, \varepsilon \varepsilon) \|(w^*, W_2^*, D_{x'} \xi^*)\|_{L^2(\Sigma_t)}^2 \end{aligned}$$

for all $\varepsilon > 0$. Substitute $\|(D_{x'} w^*, D_{x'}^2 w^*)\|_{L^2(\Sigma_t)} \lesssim \|(w^*, \Delta_{x'} w^*)\|_{L^2(\Sigma_t)}$ into the last estimate and take $\varepsilon > 0$ suitably small to infer

$$(3.54) \quad \|w^*(t)\|_{L^2(\Sigma)}^2 + \|(D_{x'} w^*, D_{x'}^2 w^*)\|_{L^2(\Sigma_t)}^2 \lesssim_{K,\varepsilon} \|(w^*, W_2^*, D_{x'} \xi^*)\|_{L^2(\Sigma_t)}^2.$$

Then we combine (3.52)–(3.54), utilize (3.44) with ξ replaced by ξ^* , and apply Grönwall’s inequality to obtain

$$(3.55) \quad \begin{aligned} & \|W^*(t)\|_{L^2(\Omega^+)}^2 + \|w^*(t)\|_{L^2(\Sigma)}^2 + \|(\xi^*, \nabla \xi^*)\|_{L^2(\Omega_t^-)}^2 \\ & + \|(W_2^*, D_{x'} w^*, D_{x'}^2 w^*, D_{x'} \xi^*, D_{x'}^2 \xi^*)\|_{L^2(\Sigma_t)}^2 \lesssim_{K,\varepsilon} \|\mathbf{f}^*\|_{L^2(\Omega_t^+)}^2. \end{aligned}$$

With the ε -dependent L^2 estimates (3.49) and (3.55), we can deduce the existence of weak solutions $(W, \xi) \in L^2(\Omega_T^+) \times L^2(\Omega_T^-)$ to regularization (3.34) for any small but fixed parameter $\varepsilon \in (0, 1)$ by the standard duality argument in [4]. Regarding (3.34c) as a fourth-order parabolic equation for ψ with given source term $W_2|_{x_1=0} \in L^2(\Sigma_T)$ and zero initial data $\psi|_{t=0} = 0$, as in [4, Theorem 5.2], we can obtain that the Cauchy problem for this parabolic equation has a unique solution $\psi \in C([0, T], H^4(\mathbb{T}^2)) \cap C^1([0, T], L^2(\mathbb{T}^2))$. Therefore, for any small and fixed parameter $\varepsilon > 0$, we obtain the existence of solutions $(W, \xi, \psi) \in L^2(\Omega_T^+) \times L^2(\Omega_T^-) \times L^2((-\infty, T]; H^4(\mathbb{T}^2))$ to the regularized problem (3.34).

3.5. Uniform energy estimates. We now show the uniform-in- ε high-order energy estimates for solutions to the regularization (3.34). Let $m \geq 1$ be an integer and $\alpha = (\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4) \in \mathbb{N}^5$ satisfy $\langle \alpha \rangle := \sum_{i=0}^3 \alpha_i + 2\alpha_4 \leq m$. For clear presentation, we divide this section into five parts.

3.5.1. Prelude. Applying $D_*^\alpha := \partial_t^{\alpha_0} (\sigma \partial_1)^{\alpha_1} \partial_2^{\alpha_2} \partial_3^{\alpha_3} \partial_1^{\alpha_4}$ to (3.34a) with $\sigma := x_1(1 - x_1)$ and taking the scalar product of the resulting equations with $D_*^\alpha W$, we utilize (3.33) and (3.34f) to deduce

$$(3.56) \quad \int_{\Omega^+} \mathbf{A}_0 D_*^\alpha W \cdot D_*^\alpha W + \varepsilon \|D_*^\alpha W_2\|_{L^2(\Sigma_t)}^2 = \mathcal{Q}_\alpha(t) + \mathcal{R}_\alpha(t)$$

for

$$(3.57) \quad \begin{aligned} \mathcal{Q}_\alpha(t) & := 2 \int_{\Sigma_t} D_*^\alpha W_1 D_*^\alpha W_2 + \int_{\Sigma_t^+} (\varepsilon |D_*^\alpha W_2|^2 - 2D_*^\alpha W_1 D_*^\alpha W_2), \\ \mathcal{R}_\alpha(t) & := \int_{\Omega_t^+} D_*^\alpha W \cdot \left(2D_*^\alpha (\mathbf{f} - \mathbf{A}_4 W) - \sum_{i=0}^3 (2[D_*^\alpha, \mathbf{A}_i \partial_i] W - \partial_i \mathbf{A}_i D_*^\alpha W) \right). \end{aligned}$$

To estimate the integral \mathcal{R}_α , we obtain from (3.34a) and (3.33) that

$$(3.58) \quad \begin{pmatrix} \partial_1 W_2 \\ \partial_1 W_1 - \varepsilon \partial_1 W_2 \\ 0 \end{pmatrix} = \mathbf{f} - \mathbf{A}_4 W - \sum_{i=0,2,3} \mathbf{A}_i \partial_i W - \mathbf{A}_1^{(0)} \partial_1 W,$$

where matrix $\mathbf{A}_1^{(0)}$ vanishes on the boundaries Σ_T and Σ_T^+ . Then we can follow the proof of [30, Lemma 3.5] and use decomposition (3.33) to infer

$$(3.59) \quad \mathcal{R}_\alpha(t) \lesssim_K \mathcal{M}_1(t) := \|(\mathbf{f}, W)\|_{H_*^m(\Omega_t^+)}^2 + \mathring{C}_{m+4} \|(\mathbf{f}, W)\|_{W_*^{2,\infty}(\Omega_t^+)}^2,$$

for all $\langle \alpha \rangle \leq m$, where $\mathring{C}_m := 1 + \|(\mathring{U}, \mathring{h}, \mathring{\varphi})\|_m^2$ (cf. (3.17)) and

$$\|u\|_{W_*^{2,\infty}(\Omega_t^+)} := \sum_{\langle \alpha \rangle \leq 1} \|D_*^\alpha u\|_{W^{1,\infty}(\Omega_t^+)}.$$

3.5.2. Case $\alpha_1 > 0$. Since $\mathcal{Q}_\alpha(t) = 0$ for $\alpha_1 > 0$, we plug (3.59) into (3.56) to get

$$(3.60) \quad \sum_{\langle \alpha \rangle \leq m, \alpha_1 > 0} \|D_*^\alpha W(t)\|_{L^2(\Omega^+)}^2 \lesssim_K \mathcal{M}_1(t),$$

where $\mathcal{M}_1(t)$ is given in (3.59).

3.5.3. Case $\alpha_1 = 0$ and $\alpha_4 > 0$. It follows from the identity (3.58) that

$$(3.61) \quad \mathcal{Q}_\alpha(t) \lesssim \sum_{i=0,2,3} \|D_*^{\alpha-\mathbf{e}}(\mathbf{f}, \mathbf{A}_4 W, \mathbf{A}_i \partial_i W, \mathbf{A}_1^{(0)} \partial_1 W)\|_{L^2(\Sigma_t \cup \Sigma_t^+)}^2$$

for $\mathbf{e} := (0, 0, 0, 0, 1)$. Use the trace theorem (cf. [18]) and the Moser-type calculus inequalities (cf. [17, Theorem B.3]) for anisotropic Sobolev spaces to obtain

$$(3.62) \quad \|D_*^{\alpha-\mathbf{e}}(\mathbf{f}, \mathbf{A}_4 W)\|_{L^2(\Sigma_t \cup \Sigma_t^+)}^2 \lesssim_K \mathcal{M}_1(t),$$

and

$$(3.63) \quad \begin{aligned} \|D_*^{\alpha-\mathbf{e}}(\mathbf{A}_i \partial_i W)\|_{L^2(\Sigma_t \cup \Sigma_t^+)}^2 &\lesssim_K \sum_{0 < \beta \leq \alpha - \mathbf{e}} \|(\partial_i D_*^{\alpha-\mathbf{e}} W, D_*^\beta \mathbf{A}_i D_*^{\alpha-\mathbf{e}-\beta} \partial_i W)\|_{L^2(\Sigma_t \cup \Sigma_t^+)}^2 \\ &\lesssim_K \|D_*^{\alpha-\mathbf{e}} W\|_{H^1(\Sigma_t \cup \Sigma_t^+)}^2 + \sum_{0 < \beta \leq \alpha - \mathbf{e}} \|D_*^\beta \mathbf{A}_i D_*^{\alpha-\mathbf{e}-\beta} \partial_i W\|_{H_*^2(\Omega_t^+)}^2 \\ &\lesssim_K \|W\|_{H_*^m(\Omega_t^+)}^2 + \mathring{C}_{m+4} \|W\|_{W_*^{2,\infty}(\Omega_t^+)}^2 \lesssim_K \mathcal{M}_1(t) \quad \text{for } i = 0, 2, 3. \end{aligned}$$

Since $(D_*^\beta \mathbf{A}_1^{(0)})|_{\Sigma_T \cup \Sigma_T^+} = 0$ for $\beta = (\beta_0, \beta_1, \beta_2, \beta_3, \beta_4) \in \mathbb{N}^5$ with $\beta_4 = 0$, we derive

$$(3.64) \quad \begin{aligned} \|D_*^{\alpha-\mathbf{e}}(\mathbf{A}_1^{(0)} \partial_1 W)\|_{L^2(\Sigma_t \cup \Sigma_t^+)}^2 &\lesssim \sum_{\mathbf{e} \leq \beta \leq \alpha - \mathbf{e}} \|D_*^\beta \mathbf{A}_1^{(0)} D_*^{\alpha-\mathbf{e}-\beta} \partial_1 W\|_{L^2(\Sigma_t \cup \Sigma_t^+)}^2 \\ &\lesssim \sum_{\mathbf{e} \leq \beta \leq \alpha - \mathbf{e}} \|D_*^{\beta-\mathbf{e}}(\partial_1 \mathbf{A}_1^{(0)}) D_*^{\alpha-\beta} W\|_{H_*^2(\Omega_t^+)}^2 \lesssim_K \mathcal{M}_1(t). \end{aligned}$$

Substituting (3.59) and (3.61)–(3.64) into (3.56) implies

$$(3.65) \quad \sum_{\langle \alpha \rangle \leq m, \alpha_1 = 0, \alpha_4 > 0} \left(\|D_*^\alpha W(t)\|_{L^2(\Omega^+)}^2 + \varepsilon \|D_*^\alpha W_2\|_{L^2(\Sigma_t)}^2 \right) \lesssim_K \mathcal{M}_1(t).$$

3.5.4. Case $\alpha_1 = \alpha_4 = 0$. We have $D_*^\alpha = \partial_t^{\alpha_0} \partial_2^{\alpha_2} \partial_3^{\alpha_3}$ and $\alpha_0 + \alpha_2 + \alpha_3 \leq m$. It follows from the boundary conditions (3.34f) and (3.34d) that

$$(3.66) \quad \mathcal{Q}_\alpha(t) = 2 \int_{\Sigma_t} D_*^\alpha W_1 D_*^\alpha W_2 = \sum_{i=1}^4 \int_{\Sigma_t} Q_i,$$

where

$$(3.67) \quad Q_1 := 2[D_*^\alpha, \dot{h}' \cdot D_{x'}] \xi D_*^\alpha W_2, \quad Q_2 := 2\dot{h}' \cdot D_{x'} D_*^\alpha \xi D_*^\alpha W_2,$$

$$(3.68) \quad Q_3 := 2\mathfrak{s} D_*^\alpha D_{x'} \cdot (\dot{B} D_{x'} \psi) D_*^\alpha W_2, \quad Q_4 := -2D_*^\alpha (\dot{b}_2 \psi) D_*^\alpha W_2.$$

Let us present the estimates for Q_i in the following four steps.

Step 1: Estimate for Q_1 . Passing to the volume integral and using (3.34f) yield

$$(3.69) \quad \begin{aligned} \int_{\Sigma_t} Q_1 &= -2 \int_{\Omega_t^+} \partial_1 \left([D_*^\alpha, \dot{h}'_\# \cdot D_{x'}] \xi_\# D_*^\alpha W_2 \right) \\ &= \underbrace{-2 \int_{\Omega_t^+} \partial_1 [D_*^\alpha, \dot{h}'_\# \cdot D_{x'}] \xi_\# D_*^\alpha W_2}_{\mathcal{Q}_{1a}} - \underbrace{2 \int_{\Omega_t^+} [D_*^\alpha, \dot{h}'_\# \cdot D_{x'}] \xi_\# D_*^\alpha \partial_1 W_2}_{\mathcal{Q}_{1b}}, \end{aligned}$$

where we denote $\dot{h}'_\#(t, x_1, x') := \dot{h}'(t, -x_1, x')$ and $\xi_\#(t, x_1, x') := \xi(t, -x_1, x')$. It follows directly from Cauchy's inequality that

$$(3.70) \quad \mathcal{Q}_{1a} \lesssim \|\partial_1 [D_*^\alpha, \dot{c}_0] D_{x'} \xi\|_{L^2(\Omega_t^-)}^2 + \|W_2\|_{H_*^m(\Omega_t^+)}^2.$$

If $\langle \alpha \rangle \leq m - 1$, then the integral \mathcal{Q}_{1b} can be estimated as

$$(3.71) \quad \mathcal{Q}_{1b} \lesssim \|[D_*^\alpha, \dot{c}_0] D_{x'} \xi\|_{L^2(\Omega_t^-)}^2 + \|D_*^\alpha \partial_1 W_2\|_{L^2(\Omega_t^+)}^2.$$

If $\langle \alpha \rangle = m$, then we choose $\beta < \alpha$ with $\langle \beta \rangle = m - 1$ and employ integration by parts to derive

$$(3.72) \quad \begin{aligned} \mathcal{Q}_{1b} &\lesssim \int_{\Omega^+} \left| [D_*^\alpha, \dot{h}'_\# \cdot D_{x'}] \xi_\# D_*^\beta \partial_1 W_2 \right| + \|[D_*^\alpha, \dot{h}' \cdot D_{x'}] \xi\|_{H^1(\Omega_t^-)}^2 + \|D_*^\beta \partial_1 W_2\|_{L^2(\Omega_t^+)}^2 \\ &\lesssim \epsilon \|D_*^\beta \partial_1 W_2\|_{L^2(\Omega^+)}^2 + C(\epsilon) \|[D_*^\alpha, \dot{c}_0] D_{x'} \xi\|_{H^1(\Omega_t^-)}^2 + \|D_*^\beta \partial_1 W_2\|_{L^2(\Omega_t^+)}^2 \end{aligned}$$

for all $\epsilon > 0$. To estimate the last terms in (3.71)–(3.72), we compute from identities (3.58) and (3.33) that for all $\gamma = (\gamma_0, 0, \gamma_2, \gamma_3, 0)$ with $\langle \gamma \rangle \leq m - 1$,

$$(3.73) \quad \begin{aligned} \|D_*^\gamma \partial_1 W_2(t)\|_{L^2(\Omega^+)}^2 &\lesssim_K \sum_{\langle \beta \rangle \leq m} \|D_*^\beta W(t)\|_{L^2(\Omega^+)}^2 + \|(\mathbf{f}, \mathbf{A}_4 W)\|_{H_*^m(\Omega_t^+)}^2 \\ &\quad + \sum_{i=0,2,3} \|[D_*^\gamma, \mathbf{A}_i \partial_i] W, [D_*^\gamma, \mathbf{A}_1^{(0)} \partial_1] W(t)\|_{L^2(\Omega^+)}^2 \\ &\lesssim_K \sum_{\langle \beta \rangle \leq m} \|D_*^\beta W(t)\|_{L^2(\Omega^+)}^2 + \mathcal{M}_1(t), \end{aligned}$$

$$(3.74) \quad \|D_*^\gamma \partial_1 W_2\|_{L^2(\Omega_t^+)}^2 \lesssim_K \mathcal{M}_1(t),$$

where $\mathcal{M}_1(t)$ is defined in (3.59). Plugging (3.70)–(3.72) into (3.69), we use (3.73)–(3.74) and the Moser-type calculus inequalities to discover

$$(3.75) \quad \int_{\Sigma_t} Q_1 \lesssim_K \epsilon \sum_{\langle \beta \rangle \leq m} \|D_*^\beta W(t)\|_{L^2(\Omega^+)}^2 + C(\epsilon)\mathcal{M}_2(t) + C(\epsilon)\mathcal{M}_1(t)$$

for all $\epsilon > 0$, where

$$(3.76) \quad \mathcal{M}_2(t) := \|\nabla \xi\|_{H^m(\Omega_t^-)}^2 + \dot{C}_{m+4} \|\nabla \xi\|_{L^\infty(\Omega_t^-)}^2.$$

Step 2: Estimate for Q_2 . For Q_2 defined in (3.67), from (3.34c), we have

$$(3.77) \quad \begin{aligned} Q_2 = & \underbrace{2D_{x'}D_*^\alpha \xi \cdot D_*^\alpha \partial_t(\dot{h}'\psi)}_{Q_{2a}} + \underbrace{2\dot{h}' \cdot D_{x'}D_*^\alpha \xi (\dot{v}' \cdot D_{x'})D_*^\alpha \psi}_{Q_{2b}} + \underbrace{2\epsilon \dot{h}' \cdot D_{x'}D_*^\alpha \xi D_*^\alpha \Delta_{x'}^2 \psi}_{Q_{2c}} \\ & \underbrace{-2D_{x'}D_*^\alpha \xi \cdot [D_*^\alpha \partial_t, \dot{h}']\psi + 2\dot{h}' \cdot D_{x'}D_*^\alpha \xi \{ [D_*^\alpha, \dot{v}' \cdot D_{x'}]\psi + D_*^\alpha(\dot{b}_1\psi) \}}_{Q_{2d}}. \end{aligned}$$

In view of the boundary condition (3.34e), we find

$$(3.78) \quad \int_{\Sigma_t} Q_{2a} = -2 \underbrace{\int_{\Sigma_t} D_*^\alpha \xi D_*^\alpha \partial_t(\dot{A}\nabla \xi)_1}_{\mathcal{J}_1} + 2\epsilon \underbrace{\int_{\Sigma_t} D_*^\alpha \xi D_*^\alpha \partial_t(\Delta_{x'} \xi - \Delta_{x'}^2 \xi)}_{\mathcal{J}_2}.$$

Pass \mathcal{J}_1 to the volume integral and use the elliptic equation (3.34b) to derive

$$\begin{aligned} \mathcal{J}_1 &= -2 \int_{\Omega_t^-} \nabla \cdot (D_*^\alpha \partial_t(\dot{A}\nabla \xi) D_*^\alpha \xi) = -2 \int_{\Omega_t^-} \partial_t D_*^\alpha(\dot{A}\nabla \xi) \cdot D_*^\alpha \nabla \xi \\ &= - \int_{\Omega^-} \dot{A} D_*^\alpha \nabla \xi \cdot D_*^\alpha \nabla \xi + \int_{\Omega_t^-} (\partial_t \dot{A} D_*^\alpha \nabla \xi - 2[\partial_t D_*^\alpha, \dot{A}]\nabla \xi) \cdot D_*^\alpha \nabla \xi. \end{aligned}$$

Regarding term \mathcal{J}_2 , we see that it is a good term, since

$$(3.79) \quad \mathcal{J}_2 = -\epsilon \int_{\Sigma} (|D_*^\alpha D_{x'} \xi|^2 + |D_*^\alpha \Delta_{x'} \xi|^2) \leq -\epsilon \|(D_*^\alpha D_{x'} \xi, D_*^\alpha D_{x'}^2 \xi)(t)\|_{L^2(\Sigma)}^2.$$

Substituting the above estimates for \mathcal{J}_1 and \mathcal{J}_2 into (3.78) yields

$$(3.80) \quad \int_{\Sigma_t} Q_{2a} + \int_{\Omega^-} \dot{A} D_*^\alpha \nabla \xi \cdot D_*^\alpha \nabla \xi + \epsilon \|(D_*^\alpha D_{x'} \xi, D_*^\alpha D_{x'}^2 \xi)(t)\|_{L^2(\Sigma)}^2 \lesssim_K \mathcal{M}_2(t),$$

where $\mathcal{M}_2(t)$ is defined by (3.76). For term Q_{2b} given in (3.77), we use identity (3.39) with ξ and ψ replaced respectively by $D_*^\alpha \xi$ and $D_*^\alpha \psi$ to get

$$(3.81) \quad \int_{\Sigma_t} Q_{2b} = 2 \underbrace{\int_{\Sigma_t} \dot{v}' \cdot D_{x'} D_*^\alpha \xi D_*^\alpha D_{x'} \cdot (\dot{h}'\psi)}_{\mathcal{J}_3} + \mathcal{J}_4 + \mathcal{J}_5,$$

where

$$\mathcal{J}_4 := -2 \sum_{i=2,3} \int_{\Sigma_t} \dot{v}' \cdot D_{x'} D_*^\alpha \xi [D_*^\alpha \partial_i, \dot{h}_i]\psi, \quad \mathcal{J}_5 := \int_{\Sigma_t} \dot{c}_1 D_*^\alpha \psi D_*^\alpha D_{x'} \xi.$$

In light of (3.34e), we obtain

$$(3.82) \quad \mathcal{J}_3 = 2 \underbrace{\int_{\Sigma_t} \dot{v}' \cdot D_{x'} D_*^\alpha \xi D_*^\alpha (\mathring{A} \nabla \xi)}_{\mathcal{J}_{3a}} - 2\varepsilon \underbrace{\int_{\Sigma_t} \dot{v}' \cdot D_{x'} D_*^\alpha \xi D_*^\alpha (\Delta_{x'} - \Delta_{x'}^2) \xi}_{\mathcal{J}_{3b}}.$$

It follows from (3.34b) and integration by parts that

$$(3.83) \quad \begin{aligned} \mathcal{J}_{3a} &= 2 \int_{\Omega_t^-} \nabla \cdot (\dot{v}'_\# \cdot D_{x'} D_*^\alpha \xi D_*^\alpha (\mathring{A} \nabla \xi)) = 2 \int_{\Omega_t^-} \nabla (\dot{v}'_\# \cdot D_{x'} D_*^\alpha \xi) \cdot D_*^\alpha (\mathring{A} \nabla \xi) \\ &= \int_{\Omega_t^-} \dot{c}_2 \nabla D_*^\alpha \xi \cdot \{D_*^\alpha (\dot{c}_1 \nabla \xi) + [D_{x'} D_*^\alpha, \dot{c}_1] \nabla \xi\} \lesssim_K \mathcal{M}_2(t), \end{aligned}$$

where we denote $\dot{v}'_\#(t, x_1, x') := \dot{v}'(t, -x_1, x')$ and $\mathcal{M}_2(t)$ is defined by (3.76). And

$$(3.84) \quad \begin{aligned} \mathcal{J}_{3b} &= 2\varepsilon \int_{\Sigma_t} \{D_{x'} (\dot{v}' \cdot D_{x'}) D_*^\alpha \xi \cdot D_{x'} D_*^\alpha \xi + \Delta_{x'} (\dot{v}' \cdot D_{x'}) D_*^\alpha \xi \Delta_{x'} D_*^\alpha \xi\} \\ &\lesssim_K \varepsilon \| (D_{x'} \xi, D_{x'}^2 \xi) \|_{H^m(\Sigma_t)}^2. \end{aligned}$$

For the terms $\mathcal{J}_4, \mathcal{J}_5$ and the integral of Q_{2d} defined in (3.77), we use the Moser-type calculus inequalities and (3.44) with ξ replaced by $D_*^\alpha \xi$ to infer

$$(3.85) \quad \begin{aligned} \mathcal{J}_4 + \mathcal{J}_5 + \int_{\Sigma_t} Q_{2d} &= \sum_{i=0,2,3} \int_{\Sigma_t} \{\dot{c}_1 [D_*^\alpha \partial_i, \dot{c}_0] \psi + \dot{c}_1 D_*^\alpha (\dot{c}_1 \psi)\} D_{x'} D_*^\alpha \xi \\ &= \sum_{i=0,2,3} \int_{\Sigma_t} D_{x'} \cdot \{\dot{c}_1 [D_*^\alpha \partial_i, \dot{c}_0] \psi + \dot{c}_1 D_*^\alpha (\dot{c}_1 \psi)\} D_*^\alpha \xi \\ &\lesssim_K \mathcal{M}_3(t) + \|D_*^\alpha \xi\|_{L^2(\Sigma_t)}^2 \lesssim_K \mathcal{M}_3(t) + \mathcal{M}_2(t), \end{aligned}$$

where

$$(3.86) \quad \mathcal{M}_3(t) := \|(\psi, D_{x'} \psi)\|_{H^m(\Sigma_t)}^2 + \dot{C}_{m+4} \|(\psi, D_{x'} \psi)\|_{L^\infty(\Sigma_t)}^2.$$

The integral of Q_{2c} (cf. (3.77)) can be estimated as

$$(3.87) \quad \begin{aligned} \int_{\Sigma_t} Q_{2c} &= -2\varepsilon \int_{\Sigma_t} D_{x'} (\dot{h}' \cdot D_{x'}) D_*^\alpha \xi \cdot D_*^\alpha \Delta_{x'} D_{x'} \psi \\ &\leq \varepsilon \varepsilon \|D_{x'}^3 \psi\|_{H^m(\Sigma_t)}^2 + C(K, \varepsilon) \varepsilon \| (D_{x'} \xi, D_{x'}^2 \xi) \|_{H^m(\Sigma_t)}^2 \quad \text{for all } \varepsilon > 0. \end{aligned}$$

In view of decomposition (3.77), we combine (3.80)–(3.85) with (3.87) to get

$$(3.88) \quad \begin{aligned} \int_{\Sigma_t} Q_2 + \int_{\Omega^-} \mathring{A} D_*^\alpha \nabla \xi \cdot D_*^\alpha \nabla \xi + \varepsilon \| (D_*^\alpha D_{x'} \xi, D_*^\alpha D_{x'}^2 \xi) (t) \|_{L^2(\Sigma)}^2 \\ \lesssim_K \mathcal{M}_2(t) + \mathcal{M}_3(t) + \varepsilon \varepsilon \|D_{x'}^3 \psi\|_{H^m(\Sigma_t)}^2 + C(K, \varepsilon) \varepsilon \| (D_{x'} \xi, D_{x'}^2 \xi) \|_{H^m(\Sigma_t)}^2 \end{aligned}$$

for all $\varepsilon > 0$, where $\mathcal{M}_2(t)$ and $\mathcal{M}_3(t)$ are defined in (3.76) and (3.86), respectively.

Step 3: Estimate for Q_3 . Next we consider the integral of Q_3 defined in (3.68). Thanks to the boundary condition (3.34c), we infer

$$(3.89) \quad \int_{\Sigma_t} Q_3 = -2\varepsilon \underbrace{\int_{\Sigma_t} D_*^\alpha (\mathring{B} D_{x'} \psi) \cdot (\partial_t + \dot{v}' \cdot D_{x'}) D_*^\alpha D_{x'} \psi}_{\mathcal{Q}_{3a}} + \mathcal{Q}_{3b} + \mathcal{Q}_{3c},$$

where

$$\begin{aligned} \mathcal{Q}_{3b} &:= -2\mathfrak{s} \int_{\Sigma_t} D_*^\alpha (\mathring{B} D_{x'} \psi) \cdot \left\{ [D_*^\alpha D_{x'}, \mathring{v}' \cdot D_{x'}] \psi + D_*^\alpha D_{x'} (\mathring{b}_1 \psi) \right\}, \\ \mathcal{Q}_{3c} &:= -2\mathfrak{s} \varepsilon \int_{\Sigma_t} \Delta_{x'} D_*^\alpha (\mathring{B} D_{x'} \psi) \cdot \Delta_{x'} D_*^\alpha D_{x'} \psi. \end{aligned}$$

We have derived in [31, section 2.4] the estimate for \mathcal{Q}_{3a} and \mathcal{Q}_{3b} (denoted respectively as $\mathcal{Q}_\alpha^{(2)}(t)$ and $\mathcal{Q}_\alpha^{(4)}(t)$ therein), which reads as

$$(3.90) \quad \mathcal{Q}_{3a} + \mathcal{Q}_{3b} \leq -\frac{\mathfrak{s}}{2} \int_\Sigma \frac{|D_*^\alpha D_{x'} \psi|^2}{|\mathring{N}|^3} + C(K) \mathcal{M}_3(t).$$

Noting that \mathring{B} defined in (3.11) is positive definite, we apply the Cauchy and Moser-type calculus inequalities to have

$$\begin{aligned} \mathcal{Q}_{3c} &\leq -\mathfrak{s} \varepsilon \int_{\Sigma_t} \mathring{B} D_*^\alpha \Delta_{x'} D_{x'} \psi \cdot D_*^\alpha \Delta_{x'} D_{x'} \psi + \varepsilon C(K) \|[D_*^\alpha \Delta_{x'}, \mathring{B}] D_{x'} \psi\|_{L^2(\Sigma_t)}^2 \\ (3.91) \quad &\leq -\mathfrak{s} \varepsilon \int_{\Sigma_t} \mathring{B} D_*^\alpha \Delta_{x'} D_{x'} \psi \cdot D_*^\alpha \Delta_{x'} D_{x'} \psi + \varepsilon C(K) \mathcal{M}_4(t), \end{aligned}$$

where

$$\mathcal{M}_4(t) := \|(D_{x'} \psi, D_{x'}^2 \psi)\|_{H^m(\Sigma_t)}^2 + \mathring{C}_{m+4} \|(D_{x'} \psi, D_{x'}^2 \psi)\|_{L^\infty(\Sigma_t)}^2.$$

Utilizing (3.89)–(3.91) and $\|D_{x'}^2 \psi\|_{H^m(\Sigma_t)}^2 \lesssim \|D_{x'} \psi\|_{H^m(\Sigma_t)} \|D_{x'}^3 \psi\|_{H^m(\Sigma_t)}$ leads to

$$\begin{aligned} (3.92) \quad \int_{\Sigma_t} \mathcal{Q}_3 + \frac{\mathfrak{s}}{2} \int_\Sigma \frac{|D_*^\alpha D_{x'} \psi|^2}{|\mathring{N}|^3} + \mathfrak{s} \varepsilon \int_{\Sigma_t} \mathring{B} D_*^\alpha \Delta_{x'} D_{x'} \psi \cdot D_*^\alpha \Delta_{x'} D_{x'} \psi \\ \lesssim_K \varepsilon \|D_{x'}^3 \psi\|_{H^m(\Sigma_t)}^2 + C(\varepsilon) \mathcal{M}_3(t) + \varepsilon \mathring{C}_{m+4} \|D_{x'}^2 \psi\|_{L^\infty(\Sigma_t)}^2 \end{aligned}$$

for all $\varepsilon > 0$, where $\mathcal{M}_3(t)$ is defined in (3.86).

Plugging (3.75), (3.88), and (3.92) into (3.66), we use (3.56) and (3.59) to get

$$\begin{aligned} &\|D_*^\alpha W(t)\|_{L^2(\Omega^+)}^2 + \|D_*^\alpha \nabla \xi(t)\|_{L^2(\Omega^-)}^2 + \|D_*^\alpha D_{x'} \psi(t)\|_{L^2(\Sigma)}^2 \\ &\quad + \varepsilon \|(D_*^\alpha W_2, D_*^\alpha D_{x'}^3 \psi)\|_{L^2(\Sigma_t)}^2 + \varepsilon \|(D_*^\alpha D_{x'} \xi, D_*^\alpha D_{x'}^2 \xi)(t)\|_{L^2(\Sigma)}^2 \\ &\lesssim_K C(\varepsilon) \mathcal{M}(t) + \varepsilon \mathring{C}_{m+4} \|D_{x'}^2 \psi\|_{L^\infty(\Sigma_t)}^2 + C(K, \varepsilon) \varepsilon \|(D_{x'} \xi, D_{x'}^2 \xi)\|_{H^m(\Sigma_t)}^2 \\ (3.93) \quad &\quad + \varepsilon \varepsilon \|D_{x'}^3 \psi\|_{H^m(\Sigma_t)}^2 + \varepsilon \sum_{\langle \beta \rangle \leq m} \|D_*^\beta W(t)\|_{L^2(\Omega^+)}^2 + \left| \int_{\Sigma_t} \mathcal{Q}_4 \right| \end{aligned}$$

for $\alpha_1 = \alpha_4 = 0$, where

$$(3.94) \quad \mathcal{M}(t) := \mathcal{M}_1(t) + \mathcal{M}_2(t) + \mathcal{M}_3(t).$$

Step 4: Estimate for \mathcal{Q}_4 . Let us now consider the final term \mathcal{Q}_4 given in (3.68). Utilize (3.34c) to get

$$(3.95) \quad \int_{\Sigma_t} \mathcal{Q}_4 = -2 \underbrace{\int_{\Sigma_t} D_*^\alpha (\mathring{b}_2 \psi) \cdot (\partial_t + \mathring{v}' \cdot D_{x'}) D_*^\alpha \psi}_{\mathcal{Q}_{4a}} + \mathcal{Q}_{4b} + \mathcal{Q}_{4c},$$

where

$$\begin{aligned} \mathcal{Q}_{4b} &:= -2 \int_{\Sigma_t} D_*^\alpha(\mathring{b}_2\psi) \cdot \left\{ [D_*^\alpha, \mathring{v}' \cdot D_{x'}]\psi + D_*^\alpha(\mathring{b}_1\psi) \right\}, \\ \mathcal{Q}_{4c} &:= 2\varepsilon \int_{\Sigma_t} D_{x'} D_*^\alpha(\mathring{b}_2\psi) \cdot D_*^\alpha D_{x'} \Delta_{x'} \psi. \end{aligned}$$

The estimate for \mathcal{Q}_{4a} and \mathcal{Q}_{4b} can be obtained similarly to that for the integrals $\mathcal{Q}_\alpha^{(1)}(t)$ and $\mathcal{Q}_\alpha^{(3)}(t)$ in [31, section 2.4]. Precisely, we can have

$$(3.96) \quad |\mathcal{Q}_{4a} + \mathcal{Q}_{4b}| \lesssim \|D_*^\alpha \psi(t)\|_{L^2(\Sigma)}^2 + \mathcal{M}_3(t).$$

If $\alpha_0 < m$, then we infer

$$(3.97) \quad \|D_*^\alpha \psi(t)\|_{L^2(\Sigma)}^2 \lesssim \int_{\Sigma_t} |D_*^\alpha \psi| |\partial_t D_*^\alpha \psi| \lesssim \|(\psi, D_{x'} \psi)\|_{H^m(\Sigma_t)}^2.$$

Term \mathcal{Q}_{4c} can be estimated by use of the Moser-type calculus inequalities as

$$(3.98) \quad |\mathcal{Q}_{4c}| \lesssim_K \varepsilon \varepsilon \|D_{x'}^3 \psi\|_{H^m(\Sigma_t)}^2 + C(\varepsilon) \varepsilon \mathcal{M}_3(t).$$

Plugging (3.96)–(3.98) into (3.95) implies

$$(3.99) \quad \left| \int_{\Sigma_t} Q_4 \right| \lesssim_K \varepsilon \varepsilon \|D_{x'}^3 \psi\|_{H^m(\Sigma_t)}^2 + C(\varepsilon) \mathcal{M}_3(t) \quad \text{if } \alpha_0 < m.$$

If $\alpha_0 = m$, then we get from (3.34c) that

$$Q_4 = -2\partial_t^m W_2 \left\{ [\partial_t^m, \mathring{b}_2]\psi + \mathring{b}_2 \partial_t^{m-1} \left(W_2 - (\mathring{v}' \cdot D_{x'} + \mathring{b}_1)\psi - \varepsilon \Delta_{x'}^2 \psi \right) \right\},$$

which leads to

$$(3.100) \quad \int_{\Sigma_t} Q_4 = \tilde{\mathcal{Q}}_{4a} + \tilde{\mathcal{Q}}_{4b} + 2\varepsilon \int_{\Sigma_t} \mathring{b}_2 \partial_t^m W_2 \partial_t^{m-1} \Delta_{x'}^2 \psi,$$

where

$$\begin{aligned} \tilde{\mathcal{Q}}_{4a} &:= - \int_{\Sigma} \partial_t^{m-1} W_2 \left\{ \mathring{b}_2 \partial_t^{m-1} W_2 - 2\mathring{b}_2 \partial_t^{m-1} (\mathring{v}' \cdot D_{x'} + \mathring{b}_1)\psi + 2[\partial_t^m, \mathring{b}_2]\psi \right\}, \\ \tilde{\mathcal{Q}}_{4b} &:= \int_{\Sigma_t} \partial_t^{m-1} W_2 \left\{ \partial_t \mathring{b}_2 \partial_t^{m-1} W_2 - 2\partial_t \left(\mathring{b}_2 \partial_t^{m-1} (\mathring{v}' \cdot D_{x'} + \mathring{b}_1)\psi - [\partial_t^m, \mathring{b}_2]\psi \right) \right\}. \end{aligned}$$

Applying integration by parts and using Cauchy’s inequality yield

$$\begin{aligned} \left| \tilde{\mathcal{Q}}_{4a} + \tilde{\mathcal{Q}}_{4b} \right| &\lesssim_K \|\partial_t^{m-1} W_2(t)\|_{L^2(\Sigma)}^2 + \|\partial_t^{m-1} W_2\|_{L^2(\Sigma_t)}^2 \\ &\quad + \|\mathring{b}_2 \partial_t^{m-1} (\mathring{v}' \cdot D_{x'} + \mathring{b}_1)\psi - [\partial_t^m, \mathring{b}_2]\psi\|_{H^1(\Sigma_t)}^2. \end{aligned}$$

Noting from (3.73) that

$$(3.101) \quad \begin{aligned} \|\partial_t^{m-1} W_2(t)\|_{L^2(\Sigma)}^2 &\lesssim_K \varepsilon \|\partial_t^{m-1} \partial_1 W_2(t)\|_{L^2(\Omega^+)}^2 + C(\varepsilon) \|W_2\|_{H_*^m(\Omega_t^+)}^2 \\ &\lesssim_K \varepsilon \sum_{\langle \beta \rangle \leq m} \|D_*^\beta W(t)\|_{L^2(\Omega^+)}^2 + C(\varepsilon) \mathcal{M}_1(t), \end{aligned}$$

we employ the Moser-type calculus inequalities to get

$$(3.102) \quad \left| \tilde{Q}_{4a} + \tilde{Q}_{4b} \right| \lesssim_K \epsilon \sum_{\langle \beta \rangle \leq m} \|D_*^\beta W(t)\|_{L^2(\Omega^+)}^2 + C(\epsilon)\mathcal{M}(t)$$

for all $\epsilon > 0$, where $\mathcal{M}(t)$ is defined by (3.94). Plugging (3.102) into (3.100), we find

$$(3.103) \quad \left| \int_{\Sigma_t} Q_4 \right| \lesssim_K \epsilon \varepsilon \|\partial_t^m W_2\|_{L^2(\Sigma_t)}^2 + C(\epsilon, K)\varepsilon \|\partial_t^{m-1} D_{x'}^4 \psi\|_{L^2(\Sigma_t)}^2 \\ + \epsilon \sum_{\langle \beta \rangle \leq m} \|D_*^\beta W(t)\|_{L^2(\Omega^+)}^2 + C(\epsilon)\mathcal{M}(t) \quad \text{if } \alpha_0 = m.$$

The first and second terms on the right-hand side of (3.103) can be absorbed by the left-hand side of (3.93) with $D_*^\alpha = \partial_t^m$ and $D_*^\alpha = \partial_t^{m-1} D_{x'}$, respectively. Therefore, we plug (3.99) and (3.103) into (3.93), let $\epsilon > 0$ be sufficiently small, and take a suitable combination of the resulting identities to discover

$$(3.104) \quad \|D_*^\alpha W(t)\|_{L^2(\Omega^+)}^2 + \|D_*^\alpha \nabla \xi(t)\|_{L^2(\Omega^-)}^2 + \|D_*^\alpha D_{x'} \psi(t)\|_{L^2(\Sigma)}^2 \\ + \varepsilon \| (D_*^\alpha W_2, D_*^\alpha D_{x'}^3 \psi) \|_{L^2(\Sigma_t)}^2 + \varepsilon \| (D_*^\alpha D_{x'} \xi, D_*^\alpha D_{x'}^2 \xi)(t) \|_{L^2(\Sigma)}^2 \\ \lesssim_K C(\epsilon)\mathcal{M}(t) + \varepsilon \dot{C}_{m+4} \|D_{x'}^2 \psi\|_{L^\infty(\Sigma_t)}^2 + C(K, \epsilon)\varepsilon \| (D_{x'} \xi, D_{x'}^2 \xi) \|_{H^m(\Sigma_t)}^2 \\ + \epsilon \varepsilon \|D_{x'}^3 \psi\|_{H^m(\Sigma_t)}^2 + \epsilon \sum_{\langle \beta \rangle \leq m} \|D_*^\beta W(t)\|_{L^2(\Omega^+)}^2 \quad \text{for } \alpha_1 = \alpha_4 = 0.$$

3.5.5. Conclusion. Combining (3.104) with (3.60) and (3.65), we take $\epsilon > 0$ small enough and use Grönwall’s inequality to derive

$$(3.105) \quad \sum_{\langle \alpha \rangle \leq m} \|D_*^\alpha W(t)\|_{L^2(\Omega^+)}^2 + \sum_{\langle \alpha \rangle \leq m, \alpha_1 = \alpha_4 = 0} \left(\|D_*^\alpha \nabla \xi(t)\|_{L^2(\Omega^-)}^2 + \|D_*^\alpha D_{x'} \psi(t)\|_{L^2(\Sigma)}^2 \right) \\ + \varepsilon \| (D_{x'}^3 \psi, D_{x'} \xi, D_{x'}^2 \xi) \|_{H^m(\Sigma_t)}^2 \lesssim_K \mathcal{M}(t) + \varepsilon \dot{C}_{m+4} \|D_{x'}^2 \psi\|_{L^\infty(\Sigma_t)}^2,$$

where $\mathcal{M}(t)$ is defined by (3.94) (cf. (3.59), (3.76), and (3.86)). To close the above estimate (3.105), we first obtain from (3.34c) that

$$\|\partial_t^m \psi\|_{L^2(\Sigma_t)}^2 \lesssim \|\partial_t^{m-1} W_2 - \varepsilon \partial_t^{m-1} \Delta_{x'}^2 \psi\|_{L^2(\Sigma_t)}^2 + \|(\hat{v}' \cdot D_{x'} + \hat{b}_1)\psi\|_{H^{m-1}(\Sigma_t)}^2,$$

and hence

$$(3.106) \quad \mathcal{M}_3(t) \lesssim_K \mathcal{M}_1(t) + \varepsilon^2 \|D_{x'}^3 \psi\|_{H^m(\Sigma_t)}^2 + \sum_{\alpha_0 < m, \alpha_1 = \alpha_4 = 0} \|D_*^\alpha \psi\|_{L^2(\Sigma_t)}^2 \\ + \|D_{x'} \psi\|_{H^m(\Sigma_t)}^2 + \dot{C}_{m+4} \|(\psi, D_{x'} \psi)\|_{L^\infty(\Sigma_t)}^2.$$

It follows from (3.24) and (3.34b) that

$$\partial_1 \partial_1 \xi = \hat{c}_2 \nabla \xi + \hat{c}_1 D_{x'} \nabla \xi \quad \text{in } \Omega_T^-.$$

Using the last identity and the Moser-type calculus inequalities, by induction in $k = 0, 1, \dots, m - 1$, we can deduce that

$$(3.107) \quad \|D_*^\gamma \partial_1^{k+1} \partial_1 \xi(t)\|_{L^2(\Omega^-)}^2 \lesssim_K \mathcal{M}_2(t) + \sum_{\langle \alpha \rangle \leq m, \alpha_1 = \alpha_4 = 0} \|D_*^\alpha \nabla \xi(t)\|_{L^2(\Omega^-)}^2$$

for all $\langle \gamma \rangle \leq m - k - 1$. In view of (3.97) and (3.105)–(3.107), we define the energy functional

$$\begin{aligned} \mathcal{I}(t) := & \sum_{\langle \alpha \rangle \leq m} \|D_*^\alpha W(t)\|_{L^2(\Omega^+)}^2 + \sum_{|\beta| \leq m} \|D^\beta \nabla \xi(t)\|_{L^2(\Omega^-)}^2 \\ & + \sum_{\langle \alpha \rangle \leq m, \alpha_1 = \alpha_4 = 0} \|D_*^\alpha D_{x'} \psi(t)\|_{L^2(\Sigma)}^2 + \sum_{\alpha_0 < m, \alpha_1 = \alpha_4 = 0} \|D_*^\alpha \psi(t)\|_{L^2(\Sigma)}^2 \end{aligned}$$

and find

$$(3.108) \quad \mathcal{I}(t) + \varepsilon \|(D_{x'}^3 \psi, D_{x'} \xi, D_{x'}^2 \xi)\|_{H^m(\Sigma_t)}^2 \lesssim_K \int_0^t \mathcal{I}(\tau) d\tau + \mathcal{N}(t)$$

for $\varepsilon > 0$ small enough, where

$$\mathcal{N}(t) := \|\mathbf{f}\|_{H_*^m(\Omega_t^+)}^2 + \mathring{C}_{m+4} \left(\|(\mathbf{f}, W)\|_{W_*^{2,\infty}(\Omega_t^+)}^2 + \|\nabla \xi\|_{L^\infty(\Omega_t^-)}^2 + \|\psi\|_{W^{2,\infty}(\Sigma_t)}^2 \right).$$

Apply Grönwall’s inequality to (3.108) and use the embedding inequalities to infer

$$(3.109) \quad \begin{aligned} & \mathcal{I}(t) + \varepsilon \|(D_{x'}^3 \psi, D_{x'} \xi, D_{x'}^2 \xi)\|_{H^m(\Sigma_t)}^2 \lesssim_K \mathcal{N}(t) \\ & \lesssim_K \|\mathbf{f}\|_{H_*^m(\Omega_t^+)}^2 + \mathring{C}_{m+4} \left(\|(\mathbf{f}, W)\|_{H_*^6(\Omega_t^+)}^2 + \|\nabla \xi\|_{H^6(\Omega_t^-)}^2 + \|\psi\|_{H^5(\Sigma_t)}^2 \right) \end{aligned}$$

for all $0 \leq t \leq T$ provided $T, \varepsilon > 0$ are suitably small. Integrating (3.109) over $[0, T]$, we can find $T_0 > 0$ depending on K_0 (cf. (3.19)), such that

$$(3.110) \quad \begin{aligned} & \|W\|_{H_*^m(\Omega_T^+)}^2 + \|\nabla \xi\|_{H^m(\Omega_T^-)}^2 + \|D_{x'} \psi\|_{H^m(\Sigma_T)}^2 + \|\psi\|_{H^{m-1}(\Sigma_T)}^2 \\ & \lesssim_{K_0} \|\mathbf{f}\|_{H_*^m(\Omega_T^+)}^2 + \|\mathbf{f}\|_{H_*^6(\Omega_T^+)}^2 \|(\mathring{U}, \mathring{h}, \mathring{\varphi})\|_{m+4}^2 \quad \text{for } 0 \leq T \leq T_0, \quad m \geq 6. \end{aligned}$$

Combine (3.106), (3.108), and (3.110) to get

$$(3.111) \quad \begin{aligned} & \|W\|_{H_*^m(\Omega_T^+)}^2 + \|(\xi, \nabla \xi)\|_{H^m(\Omega_T^-)}^2 + \|(\psi, D_{x'} \psi)\|_{H^m(\Sigma_T)}^2 + \varepsilon \|(D_{x'}^3 \psi, D_{x'} \xi, D_{x'}^2 \xi)\|_{H^m(\Sigma_T)}^2 \\ & \lesssim_{K_0} \|\mathbf{f}\|_{H_*^m(\Omega_T^+)}^2 + \|\mathbf{f}\|_{H_*^6(\Omega_T^+)}^2 \|(\mathring{U}, \mathring{h}, \mathring{\varphi})\|_{m+4}^2 \quad \text{for } 0 \leq T \leq T_0, \quad m \geq 6, \end{aligned}$$

provided $\varepsilon > 0$ is sufficiently small. Estimate (3.111) provides the desired uniform-in- ε estimate for solutions to regularization (3.34).

3.6. Proof of Theorem 3.1. The uniform-in- ε high-order estimate (3.111) enables us to establish the solvability of problem (3.32) by passing to the limit $\varepsilon \rightarrow 0$. Indeed, according to (3.111), we can extract a subsequence weakly convergent to $(W, \xi, \psi) \in H_*^m(\Omega_T^+) \times H^m(\Omega_T^-) \times H^m(\Sigma_T)$ satisfying estimate (3.111) with $\varepsilon = 0$. Noting that $\partial_1 W_2$ and $\sqrt{\varepsilon}(\Delta_{x'}^2 \psi, \Delta_{x'} \xi, \Delta_{x'}^2 \xi)$ are uniformly bounded in $H_*^{m-2}(\Omega_T^+)$ and $H^{m-2}(\Sigma_T)$, respectively, the passage to the limit $\varepsilon \rightarrow 0$ in (3.34) verifies that (W, ξ, ψ) solves the reduced problem (3.32). Furthermore, the uniqueness of solutions follows from estimate (3.111) with $\varepsilon = 0$.

Recall from (3.31), (3.29), (3.32a), and (3.28a) that

$$\dot{V} = V_{\mathring{\eta}} + J(\mathring{\Phi})W, \quad \dot{h} = h_{\mathring{\eta}} + \partial_1 \mathring{\Phi} \mathring{\eta}^\top \nabla \xi, \quad \mathbf{f} = J(\mathring{\Phi})^\top (f^+ - \mathbb{L}'_{e^+}(\mathring{U}, \mathring{\Phi})V_{\mathring{\eta}}).$$

Then using (3.26)–(3.27) and (3.111) with $\varepsilon = 0$, we can apply the embedding and Moser-type calculus inequalities to obtain the tame energy estimate (3.20) for the effective linear problem (3.14). This completes the proof of Theorem 3.1.

4. Nonlinear analysis. In this section, we employ a suitable Nash–Moser iteration scheme to prove Theorem 2.3, that is, the solvability of the nonlinear problem (2.5). See, for instance, [2] or [21] for a more general presentation of this method.

4.1. Approximate solutions. To apply Theorem 3.1, which is valid for functions vanishing in the past, we reduce the nonlinear problem (2.5) to that with zero initial data via the approximate solution. The compatibility conditions on the initial data introduced in section 2.2 are necessary for constructing the approximate solution in the following lemma; see [23, Lemma 21] and [32, Lemma 5.2] for the proof.

LEMMA 4.1. *Suppose that all the conditions of Lemma 2.1 are satisfied. Suppose further that the initial data (U_0, φ_0) are compatible up to order m and satisfy the constraints (2.9)–(2.10). Then there exist positive constants $C(M_0)$ and $T_1(M_0)$ depending on M_0 (cf. (2.12)), such that if $0 < T \leq T_1(M_0)$, then we can find (U^a, h^a, φ^a) that belongs to $H^{m+1}(\Omega_T^+) \times H^{m+1}(\Omega_T^-) \times H^{m+5/2}(\Sigma_T)$ and satisfies*

$$\begin{aligned} (4.1) \quad & \partial_t^\ell \mathbb{L}_+(U^a, \Phi^a)|_{t=0} = 0 && \text{in } \Omega^+ \quad \text{for } \ell = 0, \dots, m-1, \\ (4.2) \quad & \mathbb{L}_-(h^a, \Phi^a) = 0 && \text{in } \Omega_T^-, \\ (4.3) \quad & \mathbb{B}_+(U^a, h^a, \varphi^a) = 0 && \text{on } \Sigma_T^2 \times \Sigma_T^+, \\ (4.4) \quad & \mathbb{B}_-(h^a, \varphi^a) = 0 && \text{on } \Sigma_T \times \Sigma_T^-, \\ (4.5) \quad & (U^a, h^a, \varphi^a) = (U_0, h_0, \varphi_0) && \text{if } t < 0, \end{aligned}$$

where operators \mathbb{L}_\pm and \mathbb{B}_\pm are defined in (2.5a)–(2.5b) and (2.8), respectively, and $\Phi^a(t, x) := x_1 + \chi(x_1)\varphi^a(t, x')$. Moreover,

$$\begin{aligned} (4.6) \quad & \|U^a\|_{H^{m+1}(\Omega_T^+)} + \|h^a\|_{H^{m+1}(\Omega_T^-)} + \|\varphi^a\|_{H^{m+5/2}(\Sigma_T)} \leq C(M_0), \\ (4.7) \quad & \rho_* < \inf_{\Omega_T^+} \rho(U^a) \leq \sup_{\Omega_T^+} \rho(U^a) < \rho^*, \quad \|\varphi^a\|_{L^\infty(\Sigma_T)} \leq \frac{3\|\varphi_0\|_{L^\infty(\mathbb{T}^2)} + 1}{4}, \\ & H_1^a - H_2^a \partial_2 \varphi^a - H_3^a \partial_3 \varphi^a = 0 \quad \text{on } \Sigma_T, \quad H_1^a = 0 \quad \text{on } \Sigma_T^+. \end{aligned}$$

The vector function (U^a, h^a, φ^a) constructed above is called the *approximate solution* to the nonlinear problem (2.5). For

$$f^a := \begin{cases} -\mathbb{L}_+(U^a, \Phi^a) & \text{if } t > 0, \\ 0 & \text{if } t < 0, \end{cases}$$

we have from (4.1) and (4.6) that

$$(4.8) \quad f^a \in H^m(\Omega_T^+), \quad \|f^a\|_{H^m(\Omega_T^+)} \leq \delta_0(T),$$

where $\delta_0(T) \rightarrow 0$ as $T \rightarrow 0$. Using (4.1)–(4.5) implies that (U, \hat{h}, φ) solves the nonlinear problem (2.5) on the time interval $[0, T]$, if $(V, h, \psi) := (U, \hat{h}, \varphi) - (U^a, h^a, \varphi^a)$ satisfies

$$(4.9) \quad \begin{cases} \mathcal{L}_+(V, \Psi) := \mathbb{L}_+(U^a + V, \Phi^a + \Psi) - \mathbb{L}_+(U^a, \Phi^a) = f^a & \text{in } \Omega_T^+, \\ \mathcal{L}_-(h, \Psi) := \mathbb{L}_-(h^a + h, \Phi^a + \Psi) = 0 & \text{in } \Omega_T^-, \\ \mathcal{B}_+(V, h, \psi) := \mathbb{B}_+(U^a + V, h^a + h, \varphi^a + \psi) = 0 & \text{on } \Sigma_T^2 \times \Sigma_T^+, \\ \mathcal{B}_-(h, \psi) := \mathbb{B}_-(h^a + h, \varphi^a + \psi) = 0 & \text{on } \Sigma_T \times \Sigma_T^-, \\ (V, h, \psi) = 0 & \text{if } t < 0, \end{cases}$$

for $\Psi(t, x) := \chi(x_1)\psi(t, x')$. Note here that as in (3.14c) the notation $\Sigma_T^2 \times \Sigma_T^+$ means that the first two components of the corresponding vector equation are taken on Σ_T and the third one on Σ_T^+ .

4.2. Nash–Moser iteration. We first quote the properties on the smoothing operators from [1, 7, 27]. Denote by $\mathcal{F}_*^s(\Omega_T^+)$ (resp., $\mathcal{F}^s(\Omega_T^-)$) the class of $H_*^s(\Omega_T^+)$ (resp., $H^s(\Omega_T^-)$) vanishing in the past.

PROPOSITION 4.2. *Let $T > 0$ and $m \in \mathbb{N}$ with $m \geq 3$. Then there is a family of smoothing operators $\{\mathcal{S}_\theta\}_{\theta \geq 1} : \mathcal{F}_*^3(\Omega_T^+) \rightarrow \bigcap_{s \geq 3} \mathcal{F}_*^s(\Omega_T^+)$, such that*

$$(4.10a) \quad \|\mathcal{S}_\theta u\|_{H_*^k(\Omega_T^+)} \lesssim_m \theta^{(k-j)_+} \|u\|_{H_*^j(\Omega_T^+)} \quad \text{for } k, j = 1, \dots, m,$$

$$(4.10b) \quad \|\mathcal{S}_\theta u - u\|_{H_*^k(\Omega_T^+)} \lesssim_m \theta^{k-j} \|u\|_{H_*^j(\Omega_T^+)} \quad \text{for } 1 \leq k \leq j \leq m,$$

$$(4.10c) \quad \left\| \frac{d}{d\theta} \mathcal{S}_\theta u \right\|_{H_*^k(\Omega_T^+)} \lesssim_m \theta^{k-j-1} \|u\|_{H_*^j(\Omega_T^+)} \quad \text{for } k, j = 1, \dots, m,$$

where $k, j \in \mathbb{N}$ and $(k - j)_+ := \max\{0, k - j\}$. Moreover, there exist two families of smoothing operators (still denoted by \mathcal{S}_θ) acting respectively on $\mathcal{F}^3(\Omega_T^-)$ and functions defined on Σ_T , and satisfying the properties in (4.10) with norms $\|\cdot\|_{H^j(\Omega_T^-)}$ and $\|\cdot\|_{H^j(\Sigma_T)}$, respectively.

We follow [7, 27, 31, 30, 23] to describe the iteration scheme for problem (4.9).

Assumption (A-1). *Set $(V_0, h_0, \psi_0) = 0$. Let (V_k, h_k, ψ_k) be already given, vanish in the past, and satisfy $V_{k,2}|_{\Sigma_T^+} = 0$ and $h_k \times \mathbf{e}_1|_{\Sigma_T^-} = 0$ for $k = 0, 1, \dots, n$. Define $\Psi_k := \chi(x_1)\psi_k$.*

The differences $\delta V_n := V_{n+1} - V_n$, $\delta h_n := h_{n+1} - h_n$, and $\delta \psi_n := \psi_{n+1} - \psi_n$ will be determined through the effective linear problem

$$(4.11) \quad \begin{cases} \mathbb{L}'_{e^+}(U^a + V_{n+1/2}, \Phi^a + \Psi_{n+1/2})\delta \dot{V}_n = f_n^+ & \text{in } \Omega_T^+, \\ L_-(\Phi^a + \Psi_{n+1/2})\delta \dot{h}_n = f_n^- & \text{in } \Omega_T^-, \\ \mathbb{B}'_{n+1/2}(\delta \dot{V}_n, \delta \dot{h}_n, \delta \psi_n) = g_n^+ & \text{on } \Sigma_T^2 \times \Sigma_T^+, \\ \mathbb{B}'_{e^-}(h^a + h_{n+1/2}, \varphi^a + \psi_{n+1/2})(\delta \dot{h}_n, \delta \psi_n) = g_n^- & \text{on } \Sigma_T \times \Sigma_T^-, \\ (\delta \dot{V}_n, \delta \dot{h}_n, \delta \psi_n) = 0 & \text{if } t < 0, \end{cases}$$

where $(V_{n+1/2}, h_{n+1/2}, \psi_{n+1/2})$ is a suitable modified state to be specified in Proposition 4.7 so that $(U^a + V_{n+1/2}, h^a + h_{n+1/2}, \varphi^a + \psi_{n+1/2})$ satisfies (3.1)–(3.6), $\Psi_{n+1/2} := \chi(x_1)\psi_{n+1/2}$, and

$$(4.12) \quad \mathbb{B}'_{n+1/2} := \mathbb{B}'_{e^+}(U^a + V_{n+1/2}, h^a + h_{n+1/2}, \varphi^a + \psi_{n+1/2}),$$

$$(4.13) \quad \delta \dot{V}_n := \delta V_n - \frac{\partial_1(U^a + V_{n+1/2})}{\partial_1(\Phi^a + \Psi_{n+1/2})} \delta \Psi_n, \quad \delta \dot{h}_n := \delta h_n - \frac{\partial_1(h^a + h_{n+1/2})}{\partial_1(\Phi^a + \Psi_{n+1/2})} \delta \Psi_n.$$

Source terms f_n^\pm, g_n^\pm will be chosen via the accumulated error terms at step n .

Assumption (A-2). *Set $f_0^+ := \mathcal{S}_{\theta_0} f^a$ and $(e_0^\pm, \tilde{e}_0^\pm, f_0^-, g_0^\pm) := 0$ for $\theta_0 \geq 1$ sufficiently large. Let $(e_k^\pm, \tilde{e}_k^\pm, f_k^\pm, g_k^\pm)$ be given and vanish in the past for $k = 1, \dots, n - 1$.*

Under Assumptions (A-1)–(A-2), we set the accumulated error terms by

$$(4.14) \quad E_n^\pm := \sum_{k=0}^{n-1} e_k^\pm, \quad \tilde{E}_n^\pm := \sum_{k=0}^{n-1} \tilde{e}_k^\pm,$$

and compute the source terms f_n^\pm, g_n^\pm from

$$(4.15) \quad \sum_{k=0}^n f_k^+ + \mathcal{S}_{\theta_n} E_n^+ = \mathcal{S}_{\theta_n} f^a, \quad \sum_{k=0}^n f_k^- + \mathcal{S}_{\theta_n} E_n^- = 0, \quad \sum_{k=0}^n g_k^\pm + \mathcal{S}_{\theta_n} \tilde{E}_n^\pm = 0,$$

where \mathcal{S}_{θ_n} are the smoothing operators given in Proposition 4.2 with $\theta_n := (\theta_0^2 + n)^{1/2}$. Once f_n^\pm and g_n^\pm are specified, applying Theorem 3.1 to problem (4.11) can determine $(\delta \dot{V}_n, \delta \dot{h}_n, \delta \dot{\psi}_n)$. Then we obtain δV_n and δh_n from (4.13).

To define the error terms, we decompose

$$(4.16) \quad \begin{aligned} \mathcal{L}_+(V_{n+1}, \Psi_{n+1}) - \mathcal{L}_+(V_n, \Psi_n) &= \mathbb{L}'_+(U^a + V_n, \Phi^a + \Psi_n)(\delta V_n, \delta \Psi_n) + e'_{n+} \\ &= \mathbb{L}'_+(U^a + \mathcal{S}_{\theta_n} V_n, \Phi^a + \mathcal{S}_{\theta_n} \Psi_n)(\delta V_n, \delta \Psi_n) + e'_{n+} + e''_{n+} \\ &= \mathbb{L}'_+(U^a + V_{n+1/2}, \Phi^a + \Psi_{n+1/2})(\delta V_n, \delta \Psi_n) + e'_{n+} + e''_{n+} + e'''_{n+} \\ &= \mathbb{L}'_{e^+}(U^a + V_{n+1/2}, \Phi^a + \Psi_{n+1/2})\delta \dot{V}_n + e'_{n+} + e''_{n+} + e'''_{n+} + e^*_{n+}, \end{aligned}$$

$$(4.17) \quad \begin{aligned} \mathcal{L}_-(h_{n+1}, \Psi_{n+1}) - \mathcal{L}_-(h_n, \Psi_n) &= \mathbb{L}'_-(h^a + h_n, \Phi^a + \Psi_n)(\delta h_n, \delta \Psi_n) + e'_{n-} \\ &= \mathbb{L}'_-(h^a + \mathcal{S}_{\theta_n} h_n, \Phi^a + \mathcal{S}_{\theta_n} \Psi_n)(\delta h_n, \delta \Psi_n) + e'_{n-} + e''_{n-} \\ &= \mathbb{L}'_-(h^a + h_{n+1/2}, \Phi^a + \Psi_{n+1/2})(\delta h_n, \delta \Psi_n) + e'_{n-} + e''_{n-} + e'''_{n-} \\ &= L_-(\Phi^a + \Psi_{n+1/2})\delta \dot{h}_n + e'_{n-} + e''_{n-} + e'''_{n-} + e^*_{n-}, \end{aligned}$$

and

$$(4.18) \quad \begin{aligned} \mathcal{B}_+(V_{n+1}, h_{n+1}, \psi_{n+1}) - \mathcal{B}_+(V_n, h_n, \psi_n) &= \mathbb{B}'_+(U^a + V_n, h^a + h_n, \varphi^a + \psi_n)(\delta V_n, \delta h_n, \delta \psi_n) + \tilde{e}'_{n+} \\ &= \mathbb{B}'_+(U^a + \mathcal{S}_{\theta_n} V_n, h^a + \mathcal{S}_{\theta_n} h_n, \varphi^a + \mathcal{S}_{\theta_n} \psi_n)(\delta V_n, \delta h_n, \delta \psi_n) + \tilde{e}'_{n+} + \tilde{e}''_{n+} \\ &= \mathbb{B}'_{n+1/2}(\delta \dot{V}_n, \delta \dot{h}_n, \delta \psi_n) + \tilde{e}'_{n+} + \tilde{e}''_{n+} + \tilde{e}'''_{n+}, \end{aligned}$$

$$(4.19) \quad \begin{aligned} \mathcal{B}_-(h_{n+1}, \psi_{n+1}) - \mathcal{B}_-(h_n, \psi_n) &= \mathbb{B}'_-(h^a + h_n, \varphi^a + \psi_n)(\delta h_n, \delta \psi_n) + \tilde{e}'_{n-} \\ &= \mathbb{B}'_-(h^a + \mathcal{S}_{\theta_n} h_n, \varphi^a + \mathcal{S}_{\theta_n} \psi_n)(\delta h_n, \delta \psi_n) + \tilde{e}'_{n-} + \tilde{e}''_{n-} \\ &= \mathbb{B}'_{e^-}(h^a + h_{n+1/2}, \varphi^a + \psi_{n+1/2})(\delta \dot{h}_n, \delta \psi_n) + \tilde{e}'_{n-} + \tilde{e}''_{n-} + \tilde{e}'''_{n-}, \end{aligned}$$

where $\mathbb{L}'_\pm, \mathbb{L}'_{e^+}, \mathbb{B}'_\pm$, and $\mathbb{B}'_{n+1/2}$ are given in (3.8)–(3.9), (3.14a), (3.12)–(3.13), and (4.12), respectively. The description of the iteration scheme is completed by setting

$$(4.20) \quad e_n^\pm := e'_{n\pm} + e''_{n\pm} + e'''_{n\pm} + e^*_{n\pm}, \quad \tilde{e}_n^\pm := \tilde{e}'_{n\pm} + \tilde{e}''_{n\pm} + \tilde{e}'''_{n\pm}.$$

Let us formulate the inductive hypothesis. Set $m \in \mathbb{N}$ with $m \geq 13$ and $\tilde{\alpha} := m - 5$. The initial data (U_0, φ_0) are supposed to satisfy all the conditions of Lemma 4.1, which implies estimates (4.6)–(4.8). Moreover, Assumptions (A-1)–(A-2) are supposed to hold. For some integer $\alpha \in (6, \tilde{\alpha})$ and constant $\epsilon > 0$ to be chosen later on, our inductive hypothesis reads

$$(\mathbf{H}_{n-1}) \left\{ \begin{array}{l} \text{(a) } \|(\delta V_k, \delta h_k, \delta \psi_k)\|_s + \|\delta \Psi_k\|_{H^s(\Omega_T)} + \|D_{x'} \delta \psi_k\|_{H^s(\Sigma_T)} \leq \epsilon \theta_k^{s-\alpha-1} \Delta_k \\ \text{for all } k = 0, \dots, n-1 \text{ and } s = 6, \dots, \tilde{\alpha}; \\ \text{(b) } \max \{ \|\mathcal{L}_+(V_k, \Psi_k) - f^a\|_{H^s(\Omega_T^+)}, \|\mathcal{L}_-(h_k, \Psi_k)\|_{H^{s+1}(\Omega_T^-)} \} \leq 2\epsilon \theta_k^{s-\alpha-1} \\ \text{for all } k = 0, \dots, n-1 \text{ and } s = 6, \dots, \tilde{\alpha} - 2; \\ \text{(c) } \|(\mathcal{B}_+(V_k, h_k, \psi_k), \mathcal{B}_-(h_k, \psi_k))\|_{H^s \times H^{s+1}} \leq \epsilon \theta_k^{s-\alpha-1} \\ \text{for all } k = 0, \dots, n-1 \text{ and } s = 7, \dots, \alpha, \end{array} \right.$$

for $\Delta_k := \theta_{k+1} - \theta_k$ with $\theta_k := (\theta_0^2 + k)^{1/2}$, where $\|(U, h, \varphi)\|_s$ and $\|(g^+, g^-)\|_{H^s \times H^{s+1}}$ are defined by (3.17) and (3.18), respectively. Using hypothesis (\mathbf{H}_{n-1}) leads to the following result as in [7, Lemmas 6–7].

LEMMA 4.3. *If θ_0 is large enough, then*

$$\begin{aligned}
 \|(V_k, h_k, \psi_k)\|_s + \|\Psi_k\|_{H^s(\Omega_T)} &\leq \begin{cases} \epsilon \theta_k^{(s-\alpha)_+} & \text{if } s \neq \alpha, \\ \epsilon \log \theta_k & \text{if } s = \alpha, \end{cases} \\
 \|((I - \mathcal{S}_{\theta_k})V_k, (I - \mathcal{S}_{\theta_k})h_k, (I - \mathcal{S}_{\theta_k})\psi_k)\|_s + \|(I - \mathcal{S}_{\theta_k})\Psi_k\|_{H^s(\Omega_T)} &\lesssim \epsilon \theta_k^{s-\alpha}
 \end{aligned}$$

for $k = 0, \dots, n-1$ and $s = 6, \dots, \tilde{\alpha}$, where $\|\cdot\|_s$ is defined by (3.17). Moreover,

$$\|(\mathcal{S}_{\theta_k} V_k, \mathcal{S}_{\theta_k} h_k, \mathcal{S}_{\theta_k} \psi_k)\|_s + \|\mathcal{S}_{\theta_k} \Psi_k\|_{H^s(\Omega_T)} \lesssim \begin{cases} \epsilon \theta_k^{(s-\alpha)_+} & \text{if } s \neq \alpha, \\ \epsilon \log \theta_k & \text{if } s = \alpha \end{cases}$$

for $k = 0, \dots, n-1$ and $s = 6, \dots, \tilde{\alpha} + 6$.

4.3. Estimate of the error terms. This section is devoted to obtaining the estimate for the error terms E_k^\pm and \tilde{E}_k^\pm defined by (4.14) and (4.20). For this purpose, we need estimates for the second derivatives \mathbb{L}_\pm'' and \mathbb{B}_\pm'' of the operators \mathbb{L}_\pm and \mathbb{B}_\pm . In particular, for \mathbb{B}_\pm defined by (2.8), we have

$$\begin{aligned}
 &\mathbb{B}_+''(\tilde{\varphi})((V, h, \psi), (\tilde{V}, \tilde{h}, \tilde{\psi})) \\
 (4.21) \quad &= \begin{pmatrix} \tilde{v}' \cdot D_{x'} \psi + v' \cdot D_{x'} \tilde{\psi} \\ \mathfrak{s} D_{x'} \cdot \left(\frac{\tilde{\zeta} \cdot \tilde{\zeta}}{|\tilde{N}|^3} \zeta - \frac{\tilde{\zeta} \cdot \zeta}{|\tilde{N}|^3} \tilde{\zeta} - \frac{\tilde{\zeta} \cdot \zeta}{|\tilde{N}|^3} \tilde{\zeta} + \frac{3(\tilde{\zeta} \cdot \zeta)(\tilde{\zeta} \cdot \tilde{\zeta})}{|\tilde{N}|^5} \tilde{\zeta} \right) - h \cdot \tilde{h} \\ 0 \end{pmatrix},
 \end{aligned}$$

$$(4.22) \quad \mathbb{B}_-''((h, \psi), (\tilde{h}, \tilde{\psi})) = \begin{pmatrix} -\tilde{h}' \cdot D_{x'} \psi - h' \cdot D_{x'} \tilde{\psi} \\ 0 \end{pmatrix},$$

with $\zeta := D_{x'} \psi$, $\tilde{\zeta} := D_{x'} \tilde{\psi}$, and $\tilde{\zeta} := D_{x'} \tilde{\psi}$. We employ the embedding and Moser-type calculus inequalities to derive the following result (cf. [23, Proposition 23]).

PROPOSITION 4.4. *Let $T > 0$, $s \in \mathbb{N}$ with $s \geq 6$, and*

$$(V_i, h_i, \Psi_i, \psi_i) \in H_*^{s+2}(\Omega_T^+) \times H^{s+2}(\Omega_T^-) \times H^{s+2}(\Omega_T) \times H^{s+2}(\Sigma_T) \text{ for } i = 0, 1, 2.$$

If $\|(V_0, \Psi_0)\|_{H_*^6(\Omega_T^+)} + \|(h_0, \Psi_0)\|_{H^4(\Omega_T^-)} + \|\psi_0\|_{H^3(\Sigma_T)} \leq \tilde{K}$ for some $\tilde{K} > 0$, then

$$\begin{aligned} \|\mathbb{L}_+''(V_0, \Psi_0)((V_1, \Psi_1), (V_2, \Psi_2))\|_{H_*^s(\Omega_T^+)} &\lesssim_{\tilde{K}} \sum_{i+j=3} \|(V_i, \Psi_i)\|_{H_*^6(\Omega_T^+)} \\ &\quad \times \|(V_j, \Psi_j)\|_{H_*^{s+2}(\Omega_T^+)} + \|(V_0, \Psi_0)\|_{H_*^{s+2}(\Omega_T^+)} \prod_{i=1,2} \|(V_i, \Psi_i)\|_{H_*^6(\Omega_T^+)}, \\ \|\mathbb{L}_-''(h_0, \Psi_0)((h_1, \Psi_1), (h_2, \Psi_2))\|_{H^{s+1}(\Omega_T^-)} &\lesssim_{\tilde{K}} \sum_{i+j=3} \|(h_i, \Psi_i)\|_{H^4(\Omega_T^-)} \\ &\quad \times \|(h_j, \Psi_j)\|_{H^{s+2}(\Omega_T^-)} + \|(h_0, \Psi_0)\|_{H^{s+2}(\Omega_T^-)} \prod_{i=1,2} \|(h_i, \Psi_i)\|_{H^4(\Omega_T^-)}, \end{aligned}$$

and

$$\begin{aligned} &\|(\mathbb{B}_+''(\psi_0)((V_1, h_1, \psi_1), (V_2, h_2, \psi_2)), \mathbb{B}_-''((h_1, \psi_1), (h_2, \psi_2)))\|_{H^s \times H^{s+1}} \\ &\lesssim_{\tilde{K}} \|\psi_0\|_{H^{s+2}(\Sigma_T)} \prod_{i=1,2} \|\psi_i\|_{H^3(\Sigma_T)} + \sum_{i+j=3} \left\{ \|\psi_i\|_{H^3(\Sigma_T)} \|\psi_j\|_{H^{s+2}(\Sigma_T)} + \|h_i\|_{H^2(\Sigma_T)} \right. \\ &\quad \left. \times \|h_j\|_{H^s(\Sigma_T)} + \|(V_i, h_i)\|_{H^{s+1}(\Sigma_T)} \|\psi_j\|_{H^3(\Sigma_T)} + \|(V_i, h_i)\|_{H^2(\Sigma_T)} \|\psi_j\|_{H^{s+2}(\Sigma_T)} \right\}, \end{aligned}$$

where the norm $\|\cdot\|_{H^s \times H^{s+1}}$ is defined by (3.18).

We first employ Proposition 4.4 to estimate the quadratic error terms $e'_{k\pm}$ and $\tilde{e}'_{k\pm}$ defined in (4.16)–(4.19).

LEMMA 4.5. *If $\theta_0 \geq 1$ is large enough and $\epsilon > 0$ is sufficiently small, then*

$$\|e'_{k+}\|_{H_*^s(\Omega_T^+)} + \|e'_{k-}\|_{H^{s+1}(\Omega_T^-)} + \|(\tilde{e}'_{k+}, \tilde{e}'_{k-})\|_{H^s \times H^{s+1}} \lesssim \epsilon^2 \theta_k^{\varsigma_1(s)-1} \Delta_k,$$

with $\varsigma_1(s) := \max\{(s+2-\alpha)_+ + 10 - 2\alpha, s+6 - 2\alpha\}$, for all $k \in \{0, \dots, n-1\}$ and $s \in \{6, \dots, \tilde{\alpha} - 2\}$.

Proof. Rewriting the quadratic error term e'_{k-} as

$$e'_{k-} = \int_0^1 (1-\tau) \mathbb{L}_-''(h^a + h_k + \tau \delta h_k, \Phi^a + \Psi_k + \tau \delta \Psi_k)((\delta h_k, \delta \Psi_k), (\delta h_k, \delta \Psi_k)) d\tau,$$

we employ Proposition 4.4 and hypothesis (\mathbf{H}_{n-1}) to infer

$$\begin{aligned} \|e'_{k-}\|_{H^{s+1}(\Omega_T^-)} &\lesssim \|(\delta h_k, \delta \Psi_k)\|_{H^4(\Omega_T^-)} \|(\delta h_k, \delta \Psi_k)\|_{H^{s+2}(\Omega_T^-)} \\ &\quad + \|(\delta h_k, \delta \Psi_k)\|_{H^4(\Omega_T^-)}^2 \|(h^a, \Phi^a, h_k, \Psi_k, \delta h_k, \delta \Psi_k)\|_{H^{s+2}(\Omega_T^-)} \\ &\lesssim \epsilon^2 \theta_k^{s+6-2\alpha} \Delta_k^2 + \epsilon^2 \theta_k^{10-2\alpha} \Delta_k^2 (1 + \|(h_k, \Psi_k)\|_{H^{s+2}(\Omega_T^-)}) \end{aligned}$$

for all $s \in \{6, \dots, \tilde{\alpha} - 2\}$. In view of Lemma 4.3, we analyze the cases $s+2 \neq \alpha$ and $s+2 = \alpha$ separately to obtain the estimate for e'_{k-} . And the estimates for e'_{k+} and $\tilde{e}'_{k\pm}$ follow in an entirely similar way. \square

Then we obtain the following result concerning the estimate of the first substitution error terms $e''_{k\pm}$ and $\tilde{e}''_{k\pm}$ given in (4.16)–(4.19).

LEMMA 4.6. *If $\theta_0 \geq 1$ is large enough and $\epsilon > 0$ is sufficiently small, then*

$$\|e''_{k+}\|_{H_*^s(\Omega_T^+)} + \|e''_{k-}\|_{H^{s+1}(\Omega_T^-)} + \|(\tilde{e}''_{k+}, \tilde{e}''_{k-})\|_{H^s \times H^{s+1}} \lesssim \epsilon^2 \theta_k^{\varsigma_2(s)-1} \Delta_k,$$

with $\varsigma_2(s) := \max\{(s+2-\alpha)_+ + 12 - 2\alpha, s+8 - 2\alpha\}$, for all $k \in \{0, \dots, n-1\}$ and $s \in \{6, \dots, \tilde{\alpha} - 2\}$.

Proof. Applying Proposition 4.4 to the first substitution error term

$$e''_{k-} = \int_0^1 \mathbb{L}''_-(h^a + \mathcal{S}_{\theta_k} h_k + \tau(I - \mathcal{S}_{\theta_k})h_k, \Phi^a + \mathcal{S}_{\theta_k} \Psi_k + \tau(I - \mathcal{S}_{\theta_k})\Psi_k) ((\delta h_k, \delta \Psi_k), ((I - \mathcal{S}_{\theta_k})h_k, (I - \mathcal{S}_{\theta_k})\Psi_k)) \, d\tau,$$

we use hypothesis (\mathbf{H}_{n-1}) and Lemma 4.3 to deduce

$$\|e''_{k-}\|_{H^{s+1}(\Omega_T^-)} \lesssim \epsilon^2 \theta_k^{s+7-2\alpha} \Delta_k + \epsilon^2 \theta_k^{11-2\alpha} \Delta_k (1 + \|(\mathcal{S}_{\theta_k} h_k, \mathcal{S}_{\theta_k} \Psi_k)\|_{H^{s+2}(\Omega_T^-)})$$

for all $s \in \{6, \dots, \tilde{\alpha} - 2\}$. Then the estimate for e''_{k-} follows by using Lemma 4.3 again. A similar argument applies also for the terms e''_{k+} and $\tilde{e}''_{k\pm}$. \square

The construction and estimate of the modified state in [23, section 12.4] are independent of the second boundary condition in (2.5c) (cf. [30, 31]), which yields the following proposition for our problem.

PROPOSITION 4.7. *Let $\alpha \geq 8$. If $\theta_0 \geq 1$ is large enough and $\epsilon, T > 0$ are sufficiently small, then there exist functions $V_{n+1/2}$, $h_{n+1/2}$, and $\psi_{n+1/2}$ vanishing in the past, such that $(U^a + V_{n+1/2}, h^a + h_{n+1/2}, \varphi^a + \psi_{n+1/2})$ satisfies (3.1)–(3.6) for the approximate solution (U^a, h^a, φ^a) constructed in Lemma 4.1. Moreover,*

$$\begin{aligned} \psi_{n+1/2} &= \mathcal{S}_{\theta_n} \psi_n, \quad \|\mathcal{S}_{\theta_n} \Psi_n - \Psi_{n+1/2}\|_{H^s(\Omega_T)} \lesssim \epsilon \theta_n^{s-\alpha} \quad \text{for } s = 6, \dots, \tilde{\alpha} + 6, \\ v'_{n+1/2} &= \mathcal{S}_{\theta_n} v'_n, \quad \|\mathcal{S}_{\theta_n} V_n - V_{n+1/2}\|_{H^s_*(\Omega_T^+)} + \|\mathcal{S}_{\theta_n} h_n - h_{n+1/2}\|_{H^{s+1}(\Omega_T^-)} \lesssim \epsilon \theta_n^{s+2-\alpha} \end{aligned}$$

for $s = 6, \dots, \tilde{\alpha} + 4$ and $\Psi_{n+1/2} := \chi(x_1)\psi_{n+1/2}$.

We have the estimate for the second substitution error terms $e'''_{k\pm}$ and $\tilde{e}'''_{k\pm}$ defined in (4.16)–(4.19).

LEMMA 4.8. *Let $\alpha \geq 8$. If $\theta_0 \geq 1$ is large enough and $\epsilon, T > 0$ are sufficiently small, then*

$$\begin{aligned} \|e'''_{k-}\|_{H^s_*(\Omega_T^+)} + \|e'''_{k-}\|_{H^{s+1}(\Omega_T^-)} &\lesssim \epsilon^2 \theta_k^{\varsigma_3(s)-1} \Delta_k, \\ \|(\tilde{e}'''_{k+}, \tilde{e}'''_{k-})\|_{H^s \times H^{s+1}} &\lesssim \epsilon^2 \theta_k^{\varsigma_2(s)-1} \Delta_k, \end{aligned}$$

with $\varsigma_3(s) := \max\{(s + 2 - \alpha)_+ + 14 - 2\alpha, s + 10 - 2\alpha\}$, for all $k \in \{0, \dots, n - 1\}$ and $s \in \{6, \dots, \tilde{\alpha} - 2\}$.

Proof. First we infer from Proposition 4.7 that

$$\begin{aligned} \tilde{e}'''_{k+} &= \int_0^1 \mathbb{B}''_+(\varphi^a + \psi_{k+1/2})((\delta V_k, \delta h_k, \delta \psi_k), (\mathcal{S}_{\theta_k} V_k - V_{k+1/2}, \mathcal{S}_{\theta_k} h_k - h_{k+1/2}, 0)) \, d\tau, \\ \tilde{e}'''_{k-} &= \int_0^1 \mathbb{B}''_-(\delta h_k, \delta \psi_k), (\mathcal{S}_{\theta_k} h_k - h_{k+1/2}, 0) \, d\tau, \end{aligned}$$

which combined with (4.21)–(4.22) yield

$$\tilde{e}'''_{k+} = \begin{pmatrix} 0 \\ -\delta h_k \cdot (\mathcal{S}_{\theta_k} h_k - h_{k+1/2}) \\ 0 \end{pmatrix}, \quad \tilde{e}'''_{k-} = \begin{pmatrix} (h'_{k+1/2} - \mathcal{S}_{\theta_k} h'_k) \cdot D_{x'} \delta \psi_k \\ 0 \end{pmatrix}.$$

Applying the Moser-type calculus inequalities to the above identities, we use hypothesis (\mathbf{H}_{n-1}) , Lemma 4.3, and Proposition 4.7 to get the estimate for $\tilde{e}'''_{k\pm}$. Apply a similar argument to deduce the estimate for $e'''_{k\pm}$. \square

Using (3.8)–(3.9), we can rewrite the last error terms $e_{n\pm}^*$ in (4.16)–(4.17) as

$$e_{n+}^* = \frac{\partial_1 \mathbb{L}_+(U^a + V_{n+1/2}, \Phi^a + \Psi_{n+1/2})}{\partial_1(\Phi^a + \Psi_{n+1/2})} \delta \Psi_n,$$

$$e_{n-}^* = \frac{\partial_1 \mathbb{L}_-(h^a + h_{n+1/2}, \Phi^a + \Psi_{n+1/2})}{\partial_1(\Phi^a + \Psi_{n+1/2})} \delta \Psi_n.$$

As in [7, section 7.6] or [30, section 4], we can get the following result by use of the embedding and Moser-type calculus inequalities, hypothesis (\mathbf{H}_{n-1}) , and Proposition 4.7.

LEMMA 4.9. *Let $\tilde{\alpha} \geq \alpha + 2$ and $\alpha \geq 8$. If $\theta_0 \geq 1$ is large enough and $\epsilon, T > 0$ are sufficiently small, then*

$$\|e_{k+}^*\|_{H_*^s(\Omega_T^+)} + \|e_{k-}^*\|_{H^{s+1}(\Omega_T^-)} \lesssim \epsilon^2 \theta_k^{\varsigma_4(s)-1} \Delta_k,$$

with $\varsigma_4(s) := \max\{(s - \alpha)_+ + 18 - 2\alpha, s + 12 - 2\alpha\}$, for all $k \in \{0, \dots, n - 1\}$ and $s \in \{6, \dots, \tilde{\alpha} - 2\}$.

Similar to [30, Lemma 4.12], we utilize Lemmas 4.5–4.9 to obtain the following estimates for the accumulated error terms E_n^\pm and \tilde{E}_n^\pm defined by (4.14) and (4.20).

LEMMA 4.10. *Let $\tilde{\alpha} = \alpha + 3$ and $\alpha \geq 12$. If $\theta_0 \geq 1$ is large enough and $\epsilon, T > 0$ are sufficiently small, then*

$$\|E_n^+\|_{H_*^{\alpha+1}(\Omega_T^+)} + \|E_n^-\|_{H^{\alpha+2}(\Omega_T^-)} \lesssim \epsilon^2 \theta_n, \quad \|(\tilde{E}_n^+, \tilde{E}_n^-)\|_{H^{\alpha+1} \times H^{\alpha+2}} \lesssim \epsilon^2,$$

where the norm $\|\cdot\|_{H^s \times H^{s+1}}$ is defined by (3.18).

4.4. Proof of Theorem 2.3. Similar to [30, Lemma 4.13], we can deduce the following estimates for source terms f_n^\pm and g_n^\pm computed from (4.15).

LEMMA 4.11. *Let $\tilde{\alpha} = \alpha + 3$ and $\alpha \geq 12$. If $\theta_0 \geq 1$ is large enough and $\epsilon, T > 0$ are sufficiently small, then*

$$\|f_n^+\|_{H_*^s(\Omega_T^+)} + \|f_n^-\|_{H^{s+1}(\Omega_T^-)} \lesssim \Delta_n (\theta_n^{s-\alpha-1} \|f^a\|_{\alpha,*,T} + \epsilon^2 \theta_n^{s-\alpha-1} + \epsilon^2 \theta_n^{\varsigma_4(s)-1}),$$

$$\|(g_n^+, g_n^-)\|_{H^{s+1} \times H^{s+2}} \lesssim \epsilon^2 \Delta_n (\theta_n^{s-\alpha-1} + \theta_n^{\varsigma_2(s+1)-1}) \quad \text{for all } s \in \{6, \dots, \tilde{\alpha}\}.$$

Similar to [7, Lemma 16] and [27, Lemma 15], applying the tame estimate (3.20) to the problem (4.11), we can use Proposition 4.7 and Lemma 4.11 to derive the estimate (a) in hypothesis (\mathbf{H}_n) .

LEMMA 4.12. *Let $\tilde{\alpha} = \alpha + 3$ and $\alpha \geq 12$. If $\epsilon, T > 0$ and $\frac{1}{\epsilon} \|f^a\|_{H_*^\alpha(\Omega_T^+)}$ are sufficiently small, and $\theta_0 \geq 1$ is large enough, then*

$$\|(\delta V_n, \delta h_n, \delta \psi_n)\|_s + \|\delta \Psi_n\|_{H^s(\Omega_T)} + \|D_{x'} \delta \psi_n\|_{H^s(\Sigma_T)} \leq \epsilon \theta_n^{s-\alpha-1} \Delta_n$$

for all $s \in \{6, \dots, \tilde{\alpha}\}$.

The next lemma gives the other estimates in hypothesis (\mathbf{H}_n) , whose proof is similar to that of [27, Lemma 16].

LEMMA 4.13. *Let $\tilde{\alpha} = \alpha + 3$ and $\alpha \geq 12$. If $\epsilon, T > 0$ and $\frac{1}{\epsilon} \|f^a\|_{H_*^\alpha(\Omega_T^+)}$ are sufficiently small, and $\theta_0 \geq 1$ is large enough, then*

$$(4.23) \quad \|\mathcal{L}_+(V_n, \Psi_n) - f^a\|_{H_*^s(\Omega_T^+)} \leq 2\epsilon \theta_n^{s-\alpha-1}, \quad \|\mathcal{L}_-(h_n, \Psi_n)\|_{H^{s+1}(\Omega_T^-)} \leq 2\epsilon \theta_n^{s-\alpha-1}$$

for all $s \in \{6, \dots, \tilde{\alpha} - 2\}$, and

$$(4.24) \quad \|(\mathcal{B}_+(V_n, h_n, \psi_n), \mathcal{B}_-(h_n, \psi_n))\|_{H^s \times H^{s+1}} \leq \epsilon \theta_n^{s-\alpha-1} \quad \text{for all } s \in \{7, \dots, \alpha\}.$$

Thanks to Lemmas 4.12–4.13, hypothesis (\mathbf{H}_n) follows from (\mathbf{H}_{n-1}) provided that $\tilde{\alpha} = \alpha + 3$ and $\alpha \geq 12$ hold, $\epsilon, T > 0$ and $\frac{1}{\epsilon} \|f^a\|_{H_*^\alpha(\Omega_T^+)}$ are sufficiently small, and $\theta_0 \geq 1$ is large enough. Fixing the constants $\alpha \geq 12$, $\tilde{\alpha} = \alpha + 3$, $\epsilon > 0$, and $\theta_0 \geq 1$, we can show hypothesis (\mathbf{H}_0) as in [27, Lemma 17].

LEMMA 4.14. *If time $T > 0$ is small enough, then hypothesis (\mathbf{H}_0) is satisfied.*

We are ready to conclude the proof of our main result.

Proof of Theorem 2.3. Suppose that the initial data (U_0, φ_0) satisfy all the assumptions in Theorem 2.3. Set $\tilde{\alpha} = m - 5$ and $\alpha = \tilde{\alpha} - 3 \geq 12$. Then the initial data (U_0, φ_0) are compatible up to order $m = \tilde{\alpha} + 5$. Taking $\epsilon, T > 0$ small enough and $\theta_0 \geq 1$ suitably large can verify all the requirements of Lemmas 4.12–4.14 due to (4.6)–(4.8). Therefore, we can find some $T > 0$, such that hypothesis (\mathbf{H}_n) is satisfied for all $n \in \mathbb{N}$, implying

$$\sum_{n=0}^{\infty} \left(\|\delta V_n\|_{H_*^s(\Omega_T^+)} + \|\delta h_n\|_{H^s(\Omega_T^-)} + \|(\delta \psi_n, D_{x'} \delta \psi_n)\|_{H^s(\Sigma_T)} \right) \lesssim \sum_{n=0}^{\infty} \theta_n^{s-\alpha-2} < \infty$$

for all $s \in \{6, \dots, \alpha - 1\}$. So the sequence (V_n, h_n, ψ_n) converges to some limit (V, h, ψ) in $H_*^{\alpha-1}(\Omega_T^+) \times H^{\alpha-1}(\Omega_T^-) \times H^{\alpha-1}(\Sigma_T)$. Furthermore, $V_n \rightarrow V$ in $H^{[(\alpha-1)/2]}(\Omega_T^+)$ as $n \rightarrow \infty$ and $D_{x'} \psi \in H^{\alpha-1}(\Sigma_T)$. Passing to the limit in (4.23)–(4.24) for $s = m - 9$ implies (4.9), and hence $(U, \hat{h}, \varphi) = (U^a + V, h^a + h, \varphi^a + \psi)$ is a solution of problem (2.5) on $[0, T]$. The uniqueness of solutions to problem (2.5) can be achieved by a standard argument (see, e.g., [23, section 13] or [31, section 3.5]). The proof is complete. \square

Acknowledgments. The authors are grateful to the referees for insightful comments and suggestions that improved the quality of the paper.

REFERENCES

- [1] S. ALINHAC, *Existence d'ondes de raréfaction pour des systèmes quasi-linéaires hyperboliques multidimensionnels*, Commun. Partial Differential Equations, 14 (1989), pp. 173–230.
- [2] S. ALINHAC AND P. GÉRARD, *Pseudo-differential Operators and the Nash–Moser Theorem*, AMS, Providence, RI, 2007.
- [3] A. A. BLANK, K. O. FRIEDRICHS, AND H. GRAD, *Notes on Magneto-hydrodynamics. V: Theory of Maxwell's Equations Without Displacement Current*, Report NYO-6486-V, New York University, 1957.
- [4] J. CHAZARAIN AND A. PIRIOU, *Introduction to the Theory of Linear Partial Differential Equations*. North-Holland, Amsterdam, 1982.
- [5] G.-Q. CHEN AND Y.-G. WANG, *Existence and stability of compressible current-vortex sheets in three-dimensional magnetohydrodynamics*, Arch. Ration. Mech. Anal., 187 (2008), pp. 369–408.
- [6] S. CHEN, *Initial boundary value problems for quasilinear symmetric hyperbolic systems with characteristic boundary*, Chinese Ann. Math. 3 (1982), pp. 222–232 (in chinese); Front. Math. China 2 (2007), pp. 87–102 (in English).
- [7] J.-F. COULOMBEL AND P. SECCHI, *Nonlinear compressible vortex sheets in two space dimensions*, Ann. Sci. Éc. Norm. Supér. (4), 41 (2008), pp. 85–139.
- [8] D. COUTAND AND S. SHKOLLER, *Well-posedness of the free-surface incompressible Euler equations with or without surface tension*, J. Amer. Math. Soc., 20 (2007), pp. 829–930.
- [9] D. COUTAND, J. HOLE, AND S. SHKOLLER, *Well-posedness of the free-boundary compressible 3-D Euler equations with surface tension and the zero surface tension limit*, SIAM J. Math. Anal., 45 (2013), pp. 3690–3767.

- [10] J. M. DELHAYE, *Jump conditions and entropy sources in two-phase systems. Local instant formulation*, Int. J. Multiph. Flow, 1 (1974), pp. 395–409.
- [11] L. C. EVANS, *Partial Differential Equations*, 2nd ed., AMS, Providence, RI, 2010.
- [12] H. GOEDBLOED, R. KEPPENS, AND S. POEDTS, *Magnetohydrodynamics of Laboratory and Astrophysical Plasmas*, Cambridge University Press, Cambridge, UK, 2019.
- [13] L. HÖRMANDER, *The boundary problems of physical geodesy*, Arch. Ration. Mech. Anal., 62 (1976), pp. 1–52.
- [14] L. D. LANDAU AND E. M. LIFSHITZ, *Electrodynamics of Continuous Media*, 2nd ed., Pergamon Press, Oxford, UK, 1984.
- [15] H. LINDBLAD AND J. ZHANG, *Anisotropic Regularity of the Free-Boundary Problem in Compressible Ideal Magnetohydrodynamics*, preprint, arXiv:2106.12173, 2021.
- [16] A. MAJDA, *The existence of multidimensional shock fronts*, Mem. Amer. Math. Soc., 43, 1983.
- [17] A. MORANDO, P. SECCHI, AND P. TREBESCHI, *Regularity of solutions to characteristic initial-boundary value problems for symmetrizable systems*, J. Hyperbolic Differ. Equ., 6 (2009), pp. 753–808.
- [18] M. OHNO, Y. SHIZUTA, AND T. YANAGISAWA, *The trace theorem on anisotropic Sobolev spaces*, Tohoku Math. J., 46 (1994), pp. 393–401.
- [19] R. SAMULYAK, J. DU, J. GLIMM, AND Z. XU, *A numerical algorithm for MHD of free surface flows at low magnetic Reynolds numbers*, J. Comput. Phys., 226 (2007), pp. 1532–1549.
- [20] P. SECCHI, *Well-posedness of characteristic symmetric hyperbolic systems*, Arch. Ration. Mech. Anal., 134 (1996), pp. 155–197.
- [21] P. SECCHI, *On the Nash-Moser iteration technique*, in Recent Developments of Mathematical Fluid Mechanics, Amann, H., Giga, Y., Kozono, H., Okamoto, H., Yamazaki, M., eds., pp. 443–457, Birkhäuser, Basel (2016).
- [22] P. SECCHI AND Y. TRAKHININ, *Well-posedness of the linearized plasma-vacuum interface problem*, Interfaces Free Bound., 15 (2013), pp. 323–357.
- [23] P. SECCHI AND Y. TRAKHININ, *Well-posedness of the plasma-vacuum interface problem*, Nonlinearity, 27 (2014), pp. 105–169.
- [24] J. SHATAH AND C. ZENG, *Geometry and a priori estimates for free boundary problems of the Euler equation*, Comm. Pure Appl. Math., 61 (2008), pp. 698–744.
- [25] J. SHATAH AND C. ZENG, *Local well-posedness for fluid interface problems*, Arch. Ration. Mech. Anal., 199 (2011), pp. 653–705.
- [26] Y. TRAKHININ, *Existence of compressible current-vortex sheets: variable coefficients linear analysis*, Arch. Ration. Mech. Anal., 177 (2005), pp. 331–366.
- [27] Y. TRAKHININ, *The existence of current-vortex sheets in ideal compressible magnetohydrodynamics*, Arch. Ration. Mech. Anal., 191 (2009), pp. 245–310.
- [28] Y. TRAKHININ, *Local existence for the free boundary problem for nonrelativistic and relativistic compressible Euler equations with a vacuum boundary condition*, Comm. Pure Appl. Math., 62 (2009), pp. 1551–1594.
- [29] Y. TRAKHININ, *On well-posedness of the plasma-vacuum interface problem: The case of non-elliptic interface symbol*, Commun. Pure Appl. Anal., 15 (2016), pp. 1371–1399.
- [30] Y. TRAKHININ AND T. WANG, *Well-posedness of free boundary problem in non-relativistic and relativistic ideal compressible magnetohydrodynamics*, Arch. Ration. Mech. Anal., 239 (2021), pp. 1131–1176.
- [31] Y. TRAKHININ AND T. WANG, *Well-posedness for the free-boundary ideal compressible magnetohydrodynamic equations with surface tension*, Math. Ann., 383 (2022), pp. 761–808.
- [32] Y. TRAKHININ AND T. WANG, *Nonlinear stability of MHD contact discontinuities with surface tension*, Arch. Ration. Mech. Anal., 243 (2022), pp. 1091–1149.