



香港城市大學
City University of Hong Kong

專業 創新 胸懷全球
Professional · Creative
For The World

CityU Scholars

Genome assembly of the milky mangrove *Excoecaria agallocha*

Hong Kong Biodiversity Genomics Consortium, 18 authors, including; Chan, Lai Leo; Cheung, Siu Gin

Published in:
GigaByte

Published: 01/01/2024

Document Version:
Final Published version, also known as Publisher's PDF, Publisher's Final version or Version of Record

License:
CC BY

Publication record in CityU Scholars:
[Go to record](#)

Published version (DOI):
[10.46471/gigabyte.119](https://doi.org/10.46471/gigabyte.119)

Publication details:
Hong Kong Biodiversity Genomics Consortium, 18 authors, including, Chan, L. L., & Cheung, S. G. (2024). Genome assembly of the milky mangrove *Excoecaria agallocha*. *GigaByte*, 2024.
<https://doi.org/10.46471/gigabyte.119>

Citing this paper

Please note that where the full-text provided on CityU Scholars is the Post-print version (also known as Accepted Author Manuscript, Peer-reviewed or Author Final version), it may differ from the Final Published version. When citing, ensure that you check and use the publisher's definitive version for pagination and other details.

General rights

Copyright for the publications made accessible via the CityU Scholars portal is retained by the author(s) and/or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights. Users may not further distribute the material or use it for any profit-making activity or commercial gain.

Publisher permission

Permission for previously published items are in accordance with publisher's copyright policies sourced from the SHERPA RoMEO database. Links to full text versions (either Published or Post-print) are only available if corresponding publishers allow open access.

Take down policy

Contact lbscholars@cityu.edu.hk if you believe that this document breaches copyright and provide us with details. We will remove access to the work immediately and investigate your claim.

Genome assembly of the milky mangrove *Excoecaria agallocha*

Hong Kong Biodiversity Genomics Consortium^{*,†}

ABSTRACT

The milky mangrove *Excoecaria agallocha* is a latex-secreting mangrove that are distributed in tropical and subtropical regions. While its poisonous latex is regarded as a potential source of phytochemicals for biomedical applications, the genomic resources of *E. agallocha* remains limited. Here, we present a chromosomal level genome of *E. agallocha*, assembled from the combination of PacBio long-read sequencing and Omni-C data. The resulting assembly size is 1,332.45 Mb and has high contiguity and completeness with a scaffold N50 of 58.9 Mb and a BUSCO score of 98.4%, with 86.08% of sequences anchored to 18 pseudomolecules. 73,740 protein-coding genes were also predicted. The milky mangrove genome provides a useful resource for further understanding the biosynthesis of phytochemical compounds in *E. agallocha*.

Subjects Genetics and Genomics, Plant Genetics, Botany

INTRODUCTION

The milky mangrove *Excoecaria agallocha* (Euphorbiaceae) (Figure 1A), also known as blind-your-eye mangrove due to its toxic properties causing blindness when its milky latex in contact with eyes, can be found in the brackish water in tropical mangrove forests. In documented human history, this plant has traditionally been used to treat pains and stings from marine organisms, ulcers, as well as leprosy [1, 2]; and is also rich in phytoconstituents and potential source of bioactive compounds, such as polyphenols and terpenoids, for biomedical applications [3, 4]. *E. agallocha* is dioecious, and contrary to typical mangrove species, it does not exhibit specialized aerial roots for gas exchange [5, 6]. It has a relatively wide distribution globally, including Australia, Bangladesh, India, and Hong Kong. While it has important ecological values in mangroves, such as being the food sources of jewel bugs, genome of this ecologically important species is lacking.

CONTEXT

To date, a few molecular and genomic studies have been conducted on *E. agallocha*. These include a transcriptomic study on the flower sex determination of this dioecious species [7] and the assembly of its chloroplast genome [8]. However, the genome of this mangrove species remained missing. Previous studies have reported different karyotypes of *E. agallocha*, including $2n = 108$ [9], $2n = 130$ [10] and $2n = 140$ [11]. Its reported chromosome numbers were remarkably different to other species in the same genus, such as *Excoecaria acerifolia* Didr. $2n = 24$ [12].

Here, *E. agallocha* (NCBI:txid241838) has been selected as one of the species to be sequenced under the Hong Kong Biodiversity Genomics Consortium (a.k.a. EarthBioGenome Project Hong Kong), formed by researchers from 8 publicly funded universities in Hong Kong, in light to provide a useful resource for further understanding of its biology, ecology, evolution, and to set a foundation to carry out any necessary conservation measures.

Submitted: 11 January 2024
Accepted: 13 April 2024
Published: 25 April 2024

* Correspondence on behalf of the consortium. E-mail: jeromehui@cuhk.edu.hk

† Collaborative Authors: Entomological experts who validated the dataset and their affiliations appears at the end of the document

Published by GigaScience Press.

Preprint submitted at <https://doi.org/10.1101/2024.01.13.575302>

Included in the series: **Hong Kong Biodiversity Genomics** (https://doi.org/10.46471/GIGABYTE_SERIES_0006)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Gigabyte, 2024, 1–10

PacBio.hifiasm.asm.bp.p_ctg.gfa.fasta.dedupe.purged.fa.blobtools_pacbio.blobDB.json.bestsum.phylum.p8.span.100.blobplot.bam0




Figure 1. Genome assembly QC and contaminant/cobiont detection.

METHODS

Sample collection

Leaf tissues of a male individual of *E. agallocha* were collected at a mangrove sandy shore at Wu Kai Sha, New Territories, Hong Kong (22°25'51.6"N, 114°14'17.3"E) in February 2023. The sample was snap-frozen with liquid nitrogen and stored at -80 °C until DNA extraction.

High molecular weight DNA extraction

High molecular weight (HMW) genomic DNA isolation was started from grinding 1 g of leaf tissues with liquid nitrogen and performed using NucleoBond HMW DNA kit (Macherey Nagel Item No. 740160.20) with prior CTAB treatment. In brief, around 0.8 g of sample was digested in 5 mL CTAB [13] with addition of 1% PVP for 1 h. After RNase A treatment, 1/3 volume (~1.6 mL) of 3M potassium acetate was added for contaminant precipitation, followed by two washes of chloroform:IAA (24:1). The resulting supernatant (~4.2 mL) was topped up to 6 ml by adding H1 buffer from NucleoBond HMW DNA kit and continued with the manufacturer's protocol. The resulting DNA was eluted with 80 µL elution buffer (PacBio Ref. No. 101-633-500) and was subject to quality check using the NanoDrop™ One/OneC Microvolume UV-Vis Spectrophotometer, Qubit® Fluorometer, and overnight pulse-field gel electrophoresis.

Pacbio library preparation and sequencing

Prior to library preparation, DNA shearing was performed. Briefly, a dilution of 5 µg HMW DNA in 120 µL elution buffer was transferred to a g-tube (Covaris Part No. 520079) for 6 passes of centrifugation with $1,990 \times g$ for 2 min, followed by DNA purification with SMRTbell® cleanup beads (PacBio Ref. No. 102158-300). 2 µL sheared DNA was used to perform overnight pulse-field gel electrophoresis while the remaining sheared DNA was stored in a 2 mL DNA LoBind® Tube (Eppendorf Cat. No. 022431048) at 4 °C overnight. Subsequently, a SMRTbell library was constructed using the SMRTbell® prep kit 3.0 (PacBio Ref. No. 102-141-700), following the manufacturer's protocol. In brief, the sheared DNA was processed with DNA repair and then each DNA strand was polished at both ends and tailed with an A-overhang, followed by ligation of T-overhand SMRTbell adapters. The SMRTbell library was purified using SMRTbell® cleanup beads and 2 µL of eluted sample was subject to quantity assessment using Qubit® Fluorometer and fragment size examination with overnight pulse-field gel electrophoresis. After that, a nuclease treatment was processed to eliminate non-SMRTbell structures and a final size-selection step with 35% AMPure PB beads was performed to remove short fragments in the library.

The final library preparation for sequencing was performed with The Sequel® II binding kit 3.2 (PacBio Ref. No. 102-194-100). In brief, the SMRT bell structures were annealed and bound with Sequel II® primer 3.2 and Sequel II® DNA polymerase 2.2, respectively. A final cleanup was processed with SMRTbell® cleanup beads, followed by an addition of serial diluted Sequel II® DNA Internal Control Complex. The library was loaded with the diffusion loading mode at an on-plate concentration of 90 pM. The sequencing was performed on the Pacific Biosciences SEQUEL IIe System running for 30-hour movies with 120 min pre-extension to generate HiFi reads. In total, two SMRT cells were used for the sequencing. Details of the resulting sequencing data are listed in Table 1.

Table 1. Summary of genomic sequencing data.

Library	Reads	Bases	Coverage (X)	Accession number
PacBio HiFi	3,503,202	33,204,508,502	25	SRR24631716
Omnic	500,462,964	75,069,444,600	56	SRR26908863

Omnic-C library preparation and sequencing

A nuclei isolation procedure was performed from 2 g ground leaf tissues, following the modification of Workman *et al.* [14]. The resulting nuclei pellet was used to construct an Omni-C library using the Dovetail® Omni-C® Library Preparation Kit (Dovetail Cat. No. 21005) by following the manufacturer's instructions. In brief, the nuclei pellet was resuspended in 4 mL 1× PBS, followed by crosslinking with formaldehyde and DNA digestion with endonuclease DNase I. The concentration and fragment size of the digested lysate was quantified using Qubit® Fluorometer and TapeStation D5000 HS ScreenTape, respectively. Subsequently, both ends of DNA were polished and ligation of biotinylated bridge adaptors were proceeded at 22 °C for 30 min. Proximity ligation between crosslinked DNA fragments was conducted at 22 °C for 1 h, followed by crosslink reversal of DNA and then purification with SPRIselect™ Beads (Beckman Coulter Product No. B23317).

End repair and adapter ligation were conducted using the Dovetail™ Library Module for Illumina (Dovetail Cat. No. 21004). In brief, DNA was tailed with an A-overhang and then ligated with Illumina-compatible adapters at 20 °C for 15 min. The Omni-C library was sheared using USER Enzyme Mix and purified with SPRIselect™ Beads. Afterwards, the DNA fragments were isolated using Streptavidin Beads. The DNA library was amplified with Universal and Index PCR Primers from the Dovetail™ Primer Set for Illumina (Dovetail Cat. No. 25005). A final size selection step was done with SPRIselect™ Beads to retain DNA fragments ranging between 350 bp and 1000 bp only. The concentration and fragment size of the library was validated by Qubit® Fluorometer and TapeStation D5000 HS ScreenTape, respectively. The qualified library was eventually sequenced on an Illumina HiSeq-PE150 platform. Details of the resulting sequencing data are listed in Table 1.

Genome assembly and gene model prediction

De novo genome assembly was performed using Hifiasm (version 0.16.1-r375) [15] with default parameters, which was then searched against the NT database using BLAST for the input for BlobTools (v1.1.1) [16] with default parameters to identify and remove any possible contaminations (Figure 1). Haplotypic duplications were removed using “purge_dups” based on the depth of HiFi reads [17] with default parameters. Furthermore, proximity ligation data sequenced from the Omni-C library were employed to scaffold the assembly with YaHS (version 1.2a.2) [18] with default parameters.

Gene model prediction was run by funannotate (version 1.8.15) [19] using the following parameters “--repeats2evm --protein_evidence uniprot_sprot.fasta --genemark_mode ES --optimize_augustus --organism other --max_intronlen 350000”. The predicted gene models from various prediction sources including GeneMark, high-quality Augustus predictions (HiQ), pasa, Augustus, GlimmerHM and snap were combined and processed with Evidence Modeler to produce the annotation files.




Figure 2. Genomic information of *Excoecaria agallocha*. (A) Picture of a female *Excoecaria agallocha*; (B) Genome statistics; (C) Omni-C contact map of the assembly; (D) Pie chart and repeat landscape plot of repetitive elements in the assembled genome annotated by Earl Grey.

Repeat annotation

Annotation of transposable elements (TEs) were conducted using the Earl Grey TE annotation workflow pipeline (version 1.2) [20] as well as the Extensive de novo TE Annotator (EDTA) (v2.2.0) [21].

RESULTS AND DISCUSSION

Genome assembly of *Excoecaria agallocha*

A total of 33.20 Gb of HiFi reads from the whole genome of milky mangrove *Excoecaria agallocha* were generated by PacBio sequencing. After scaffolding with 75 Gb Omni-C data, 86.08% of the sequences were assembled into 18 pseudochromosomes (Figure 2B). The assembled genome size was 1,332.45 Mb, with 1,402 scaffolds and a scaffold N50 of 58.95 Mb. The complete BUSCO value was estimated to be 98.4% (viridiplantae_odb10) (Figure 2C; Table 2). The GC content was 32.17%. A total of 73,740 protein-coding genes were predicted, with a mean coding sequence length of 288 amino acids (AA) and a BUSCO score of 82.1% (Figure 2C).

Repeat content analyses revealed that transposable elements (TEs) account for approximately 40%–60% of the milky mangrove assembly by annotation tools EDTA and Earl Grey, respectively. The results and classifications of TEs are summarized in Figure 2D and Table 3.

Table 2. Genome statistic and sequencing information.

<i>Excoecaria agallocha</i>	
Total length (bp)	1,332,447,397
Number	1,402
Mean length (bp)	950,390
Longest	126,146,454
Shortest	1,000
N_count	668,400
Gaps	3,342
N50	58,948,931
N50n	8
N70	43,065,745
N70n	13
N90	1,110,898
N90n	41
BUSCO (Genome)	C:98.4% [S:39.3%, D:59.1%], F:0.5%, M:1.1%, n:425
BUSCO (Proteome)	C:82.1% [S:45.9%, D:36.2%], F:12.0%, M:5.9%, n:425
HiFi Reads	3,503,202
HiFi Bases	33,204,508,502
HiFi Q30%	3
HiFi Q20%	4
HiFi GC%	32
HiFi Nppm	0
HiFi Ave_len	9,478
HiFi Min_len	153
HiFi Max_len	33,856

Table 3. Summary of transposable element annotation by Earl Grey and EDTA.

Repeat analysis by EarlGrey				
Classification	Total length (bp)	Count	Proportion (%)	No. of distinct classifications
DNA	78,065,183	77,115	5.86	7,410
LINE	10,652,399	10,771	0.80	3,831
LTR	299,391,089	124,737	22.47	7,829
Other (Simple Repeat, Microsatellite, RNA)	1,141,160	932	0.09	349
Penelope	37,879	151	0.00	137
Rolling Circle	1,147,084	1,803	0.09	1,028
SINE	25,619	107	0.00	93
Unclassified	425,523,922	338,107	31.94	8,243
SUM	815,984,335	553,723	61.24	28,920
Repeat analysis by EDTA				
Classification		bp masked	Count	% masked
LINE	L1	15,082,662	47,152	1.13
	L2	328,592	714	0.02
	RTE	398,335	1,397	0.03
LTR	Copia	65,202,479	48,709	4.90
	Gypsy	41,881,198	47,938	3.14
	Unknown	2,526	20	0.00
TIR	CACTA	48,274,154	150,680	3.62
	Mutator	96,860,270	244,712	7.27
	PIF Harbinger	51,260,554	123,517	3.85
	Tc1 Mariner	39,900,854	48,297	3.00
	hAT	29,307,586	70,668	2.20
Non-LTR	Penelope	10,803,234	38,103	0.81
Non-TIR	helitron	74,727,754	205,818	5.61
Repeat region		57,365,013	206,847	4.31
Total interspersed		531,395,211	1,234,572	39.90

Table 4. Information of 18 pseudochromosomes and BUSCO result.

Chr Number	scaffold_length	scaffold_id	% of whole genome
1	126,146,454	scaffold_2_1	9.47%
2	125,683,813	scaffold_3_1	9.43%
3	115,235,106	scaffold_1_1	8.65%
4	75,925,046	scaffold_4_1	5.70%
5	71,251,965	scaffold_5_1	5.35%
6	70,346,551	scaffold_6_1	5.28%
7	59,451,589	scaffold_7_1	4.46%
8	58,948,931	scaffold_8_1	4.42%
9	57,619,229	scaffold_9_1	4.32%
10	52,150,882	scaffold_10_1	3.91%
11	48,057,327	scaffold_11_1	3.61%
12	46,431,560	scaffold_12_1	3.48%
13	43,065,745	scaffold_13_1	3.23%
14	41,974,720	scaffold_14_1	3.15%
15	41,514,226	scaffold_15_1	3.12%
16	40,341,847	scaffold_16_1	3.03%
17	37,296,423	scaffold_17_1	2.80%
18	35,487,587	scaffold_18_1	2.66%
SUM	1,146,929,001		86.08%
BUSCO		C:92.7% [S:35.3%, D:57.1%]	

CONCLUSION AND FUTURE PERSPECTIVE

The genome assembly of *E. agallocha* presented in this study is the first genomic resource for this mangrove species, which provides a valuable resource for further investigation in the biosynthesis of phytochemical compounds in its milky latex and for the understanding of biology and evolution in genome architecture in the Euphorbiaceae family.

DATA VALIDATION AND QUALITY CONTROL

During HMW DNA extraction and Pacbio library preparation, quality control of the sample or library was assessed with the NanoDrop™ One/OneC Microvolume UV–Vis Spectrophotometer, Qubit® Fluorometer, and overnight pulse-field gel electrophoresis. The Omni-C library was validated by Qubit® Fluorometer and TapeStation D5000 HS ScreenTape.

For the genome assembly, BlobTools (v1.1.1) [16] was used to identify and remove any possible contaminations (Figure 1). Benchmarking Universal Single-Copy Orthologs (BUSCO, v5.5.0) [22] was run with a collection of single-copy orthologs dataset for the Viridiplantae (Viridiplantae Odb10) to validate the completeness of the genome assembly and gene annotation (Table 2; Table 4).

DISCLAIMER

The genomic data generated in this study was not assessed for the potential level of polyploidy.

DATA AVAILABILITY

The raw reads generated in this study were deposited in the NCBI database under the SRA accession [SRR24631716](#) and [SRR26908863](#). The genome, genome annotation and repeat annotation files were made publicly available in Figshare [23].

ABBREVIATIONS

BUSCO: Benchmarking Universal Single-Copy Orthologs; HMW: High Molecular Weight; TE: Transposable Element.

DECLARATIONS

Ethics approval and consent to participate

The authors declare that ethical approval was not required for this type of research.

Competing interests

The authors declare that they do not have competing interests.

Authors' contribution

JHLH, TFC, LLC, SGC, CCC, JKHF, JDG, SCKL, YHS, CKCW, KYLY and YW conceived and supervised the study; STSL, DTWL and JSYL collected the samples; STSL and WLS carried out DNA extraction, library preparation and genome sequencing; HYY arranged the logistics of samples; WN performed genome assembly and gene model prediction.

Funding

This work was funded and supported by the Hong Kong Research Grant Council Collaborative Research Fund (C4015-20EF), CUHK Strategic Seed Funding for Collaborative Research Scheme (3133356) and CUHK Group Research Scheme (3110154).

REFERENCES

- 1 **Vadlapudi V, Bobbarala V, Penumajji S et al.** *Excoecaria agallocha* L. antimicrobial properties against important pathogenic microorganisms. *Int. J. ChemTech Res.*, 2009; **1**(4): 865–867.
- 2 **Kaliampurthi S, Selvaraj G.** Insight on *Excoecaria agallocha*: an overview. *Nat. Prod. Chem. Res.*, 2016; **4**(2): 1000203. doi:10.4172/2329-6836.1000203.
- 3 **Raghavanpillai Sabu K, Sugathan S, Idhayadhulla A et al.** Antibacterial, antifungal, and cytotoxic activity of *Excoecaria agallocha* leaf extract. *J. Exp. Pharmacol.*, 2022; **14**: 17–26. doi:10.2147/JEP.S339383.
- 4 **Sadeer NB, Zengin G, Mahomoodally MF.** Biotechnological applications of mangrove plants and their isolated compounds in medicine—a mechanistic overview. *Crit. Rev. Biotechnol.*, 2023; **43**(3): 393–414. doi:10.1080/07388551.2022.2033682.
- 5 **Mondal S, Ghosh D, Ramakrishna K.** A complete profile on blind-your-eye mangrove *Excoecaria agallocha* L. (*Euphorbiaceae*): ethnobotany, phytochemistry, and pharmacological aspects. *Pharmacogn. Rev.*, 2016; **10**(20): 123–138. doi:10.4103/0973-7847.194049.
- 6 **Kader A, Sinha SN.** Sex-related differences of *Excoecaria agallocha* L. with a view to defence and growth. *Trop. Life Sci. Res.*, 2022; **33**(2): 55–74. doi:10.21315/tlsr2022.33.2.4.
- 7 **Zhou Y, Hao L, Huang L et al.** De novo transcriptome assembly reveals characteristics of flower sex determination of *Excoecaria agallocha*. *Ann. Forest Sci.*, 2022; **79**: 36. doi:10.1186/s13595-022-01156-6.
- 8 **Shi C, Han K, Li L et al.** Complete chloroplast genomes of 14 mangroves: phylogenetic and comparative genomic analyses. *Biomed. Res. Int.*, 2020; **2020**: 8731857. doi:10.1155/2020/8731857.
- 9 **Das A.** Genetic erosion of wetland biodiversity in Bhitarkanika forest of Orissa, India Anath Bandhu DAs, Uday Chand BAsAk & Premananda DAs. *Biologia, Bratislava*, 1999; **54**(4): 15–22.
- 10 **Das AB, Jena S, Pradhan C et al.** Genetic variability among male populations of a minor mangrove *Excoecaria agallocha* L. as evident by chromosome morphology and DNA markers. *Nucleus*, 2011; **54**: 39–47. doi:10.1007/s13237-011-0027-z.
- 11 **Sidhu S.** Vegetation of east Godavari. In: Proceedings of 48th Indian Science Congress. vol. 3, 1961; pp. 358–359.

- 12 **Perry BA.** Chromosome number and phylogenetic relationships in the Euphorbiaceae. *Am. J. Bot.*, 1943; **30**: 527–543. doi:10.1002/j.1537-2197.1943.tb14796.x.
- 13 **Doyle JJ, Doyle JL.** A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.*, 1987; **19**: 11–15.
- 14 PacBio: Isolating nuclei from plant tissue using TissueRuptor disruption. 2022; <https://www.pacb.com/wp-content/uploads/Procedure-checklist-Isolating-nuclei-from-plant-tissue-using-TissueRuptor-disruption.pdf>.
- 15 **Cheng H, Concepcion GT, Feng X et al.** Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. *Nat. Methods*, 2021; **18**(2): 170–175. doi:10.1038/s41592-020-01056-5.
- 16 **Laetsch DR, Blaxter ML.** BlobTools: interrogation of genome assemblies. *F1000Research*, 2017; **6**: 1287. doi:10.12688/f1000research.12232.1.
- 17 **Guan D, Guan D, McCarthy SA et al.** Identifying and removing haplotypic duplication in primary genome assemblies. *Bioinformatics*, 2020; **36**(9): 2896–2898. doi:10.1093/bioinformatics/btaa025.
- 18 **Zhou C, McCarthy SA, Durbin R.** YaHS: yet another Hi-C scaffolding tool. *Bioinformatics*, 2023; **39**(1): btac808. doi:10.1093/bioinformatics/btac808.
- 19 **Palmer JM, Stajich J.** Funannotate v1.8.1: eukaryotic genome annotation. Zenodo. 2020; <https://doi.org/10.5281/ZENODO.4054262>.
- 20 **Baril T, Galbraith J, Hayward A.** Earl Grey: a fully automated user-friendly transposable element annotation and analysis pipeline. *Mol. Biol. Evol.*, 2024; **41**: msae068. doi:10.1093/molbev/msae068.
- 21 **Ou S, Su W, Liao Y et al.** Benchmarking transposable element annotation methods for creation of a streamlined, comprehensive pipeline. *Genome Biol.*, 2019; **20**(1): 275. doi:10.1186/s13059-019-1905-y.
- 22 **Manni M, Berkeley MR, Seppey M et al.** BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Mol. Biol. Evol.*, 2021; **38**(10): 4647–4654. doi:10.1093/molbev/msab199.
- 23 **Hong Kong Biodiversity Genomics Consortium.** Genome assembly of the milky mangrove *Excoecaria agallocha*. figshare. [Dataset]. 2024; <https://doi.org/10.6084/m9.figshare.24923781>.

DETAILS OF COLLABORATIVE AUTHORS

• List of authors in Hong Kong Biodiversity Genomics Consortium

Jerome H. L. Hui,¹ Ting Fung Chan,² Leo Lai Chan,³ Siu Gin Cheung,⁴ Chi Chiu Cheang,^{5,6} James Kar-Hei Fang,⁷ Juan Diego Gaitan-Espitia,⁸ Stanley Chun Kwan Lau,⁹ Yik Hei Sung,^{10,11} Chris Kong Chu Wong,¹² Kevin Yuk-Lap Yip,^{13,14} Yingying Wei,¹⁵ Sean Tsz Sum Law,¹ Wai Lok So,¹ Wenyan Nong,¹ David Tai Wai Lau,¹⁶ Shing Yip Lee,¹⁷ Ho Yin Yip¹

¹School of Life Sciences, Simon F.S. Li Marine Science Laboratory, State Key Laboratory of Agrobiotechnology, Institute of Environment, Energy and Sustainability, The Chinese University of Hong Kong, Hong Kong, China

²School of Life Sciences, State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong, Hong Kong SAR, China

³State Key Laboratory of Marine Pollution and Department of Biomedical Sciences, City University of Hong Kong, Hong Kong SAR, China

⁴State Key Laboratory of Marine Pollution and Department of Chemistry, City University of Hong Kong, Hong Kong SAR, China

⁵Department of Science and Environmental Studies, The Education University of Hong Kong, Hong Kong SAR, China

⁶EcoEdu PEI, Charlottetown, PE, C1A 4B7, Canada

⁷Department of Food Science and Nutrition, Research Institute for Future Food, and State Key Laboratory of Marine Pollution, The Hong Kong Polytechnic University, Hong Kong SAR, China

⁸The Swire Institute of Marine Science and School of Biological Sciences, The University of Hong Kong, Hong Kong SAR, China

⁹Department of Ocean Science, The Hong Kong University of Science and Technology, Hong Kong SAR, China

¹⁰Science Unit, Lingnan University, Hong Kong SAR, China

¹¹School of Allied Health Sciences, University of Suffolk, Ipswich, IP4 1QJ, UK

¹²Croucher Institute for Environmental Sciences, and Department of Biology, Hong Kong Baptist University, Hong Kong SAR, China

¹³Department of Computer Science and Engineering, The Chinese University of Hong Kong, Hong Kong SAR, China

¹⁴Sanford Burnham Prebys Medical Discovery Institute, La Jolla, CA, USA

¹⁵Department of Statistics, The Chinese University of Hong Kong, Hong Kong SAR, China

¹⁶Shiu-Ying Hu Herbarium, School of Life Sciences, The Chinese University of Hong Kong, Hong Kong SAR, China

¹⁷Simon F.S. Li Marine Science Laboratory, The Chinese University of Hong Kong, Hong Kong SAR, China