



香港城市大學
City University of Hong Kong

專業 創新 胸懷全球
Professional · Creative
For The World

CityU Scholars

The concordance between upper and lower respiratory microbiota in children with *Mycoplasma pneumoniae* pneumonia

Dai, Wenkui; Wang, Heping; Zhou, Qian; Feng, Xin; Lu, Zhiwei; Li, Dongfang; Yang, Zhenyu; Liu, Yanhong; Li, Yinhu; Xie, Gan; Shen, Kunling ; Yang, Yonghong ; Zheng, Yuejie; Li, Shuaicheng

Published in:
Emerging Microbes and Infections

Published: 01/01/2018

Document Version:
Final Published version, also known as Publisher's PDF, Publisher's Final version or Version of Record

License:
CC BY

Publication record in CityU Scholars:
[Go to record](#)

Published version (DOI):
[10.1038/s41426-018-0097-y](https://doi.org/10.1038/s41426-018-0097-y)

Publication details:
Dai, W., Wang, H., Zhou, Q., Feng, X., Lu, Z., Li, D., Yang, Z., Liu, Y., Li, Y., Xie, G., Shen, K., Yang, Y., Zheng, Y., & Li, S. (2018). The concordance between upper and lower respiratory microbiota in children with *Mycoplasma pneumoniae* pneumonia. *Emerging Microbes and Infections*, 7, Article 92.
<https://doi.org/10.1038/s41426-018-0097-y>

Citing this paper

Please note that where the full-text provided on CityU Scholars is the Post-print version (also known as Accepted Author Manuscript, Peer-reviewed or Author Final version), it may differ from the Final Published version. When citing, ensure that you check and use the publisher's definitive version for pagination and other details.

General rights

Copyright for the publications made accessible via the CityU Scholars portal is retained by the author(s) and/or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights. Users may not further distribute the material or use it for any profit-making activity or commercial gain.

Publisher permission

Permission for previously published items are in accordance with publisher's copyright policies sourced from the SHERPA RoMEO database. Links to full text versions (either Published or Post-print) are only available if corresponding publishers allow open access.

Take down policy

Contact lbscholars@cityu.edu.hk if you believe that this document breaches copyright and provide us with details. We will remove access to the work immediately and investigate your claim.

ARTICLE

Open Access

The concordance between upper and lower respiratory microbiota in children with *Mycoplasma pneumoniae* pneumonia

Wenkui Dai¹, Heping Wang², Qian Zhou³, Xin Feng³, Zhiwei Lu², Dongfang Li^{3,4}, Zhenyu Yang³, Yanhong Liu³, Yinhu Li³, Gan Xie², Kunling Shen⁵, Yonghong Yang², Yuejie Zheng² and Shuaicheng Li¹

Abstract

In recent years, the morbidity of *Mycoplasma pneumoniae* pneumonia (MPP) has dramatically increased in China. An increasing number of studies indicate that an imbalance in the respiratory microbiota is associated with respiratory infection. We selected 28 hospitalized patients infected with *M. pneumoniae* and 32 healthy children. Nasopharyngeal (NP) and oropharyngeal (OP) swabs were collected from healthy children, whereas NP, OP and bronchoalveolar lavage (BAL) specimens were collected from patients. Microbiota analysis was performed on all microbial samples using 16 S ribosomal RNA (16 S rRNA) sequencing. The NP microbial samples in healthy children were divided into two groups, which were dominated by either *Staphylococcus* or mixed microbial components. The respiratory microbiota in pneumonia patients harbored a lower microbial diversity compared to healthy children, and both the NP and OP microbiota of patients differed significantly from that of healthy children. Hospitalized MPP children with a higher abundance of *Mycoplasma* in the BAL fluid (BALF) microbiota tended to suffer longer hospitalization lengths and higher peak fevers and serum C-reactive protein levels. Concordance analysis explained the succession of imbalanced NP microbiota to the OP and lung in diseased children. However, the association of the abundance of *Mycoplasma* in BALF microbiota with that in NP or OP microbiota varied among individuals, which suggested the sensitivity of BALF in MPP diagnostics, mirroring MPP severity.

Introduction

The respiratory tract is home to a variety of microbial commensals that cultivate the immune system and confer resistance to colonization of pathogens^{1, 2}. Increasing reports have demonstrated the association of altered respiratory microbiota with disease severity in acute respiratory infections (ARIs), such as bronchiolitis and pneumonia^{3–5}. *Mycoplasma pneumoniae* pneumonia (MPP) is resulting in an increasingly high morbidity in Chinese children⁶. Previous studies have also

demonstrated the alterations of the nasopharynx (NP) and oropharynx (OP) microbiota in MPP, as well as microbiota transmission between the NP and OP^{5, 7, 8}.

However, imbalanced lung microbiota in MPP remains unexplored. Meyer Sauter PM *et al.* showed that the clearance of *M. pneumoniae* in the lung differed significantly from that in the NP⁹, which suggested different alterations of lung microbiota in MPP compared to the NP/OP. In addition, the use of bronchoalveolar lavage fluid (BALF) is a more sensitive measure for the establishment of microbial diagnosis¹⁰ and assessment of disease severity compared with NP and OP swabs^{11, 12}. Given that BALF collection is restricted to cases of severe disease, it is imperative to explore the possibility of predicting lung microbiota alterations via NP or OP microbiota analysis.

Correspondence: Yuejie Zheng (shine1990@sina.com) or Shuaicheng Li (shuaicli@cityu.edu.hk)

¹Department of Computer Science, City University of Hong Kong, 999077 Hong Kong, China

²Department of Respiratory Diseases, Shenzhen Children's Hospital, No. 7019, Yitian Road, 518026 Futian District, Shenzhen, China

Full list of author information is available at the end of the article.

© The Author(s) 2018



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

In this study, we enrolled 28 MPP and 32 healthy children to analyze respiratory microbiota via 16 S rRNA analysis. We intended to answer the following two questions: 1) how the imbalanced NP/OP microbiota is associated with lung microbiota in MPP and 2) how altered respiratory microbiota is associated with MPP severity.

Results

Sample characteristics

Both healthy and diseased children were not exposed to antibiotics for at least one month prior to sampling to ensure no confounding effect of antibiotics on the respiratory microbiota. None of the recruited healthy children suffered any chronic respiratory diseases or acute respiratory infection during the one month prior to sampling and for one week after sampling. Selected patients had chest radiographic abnormalities consistent with pneumonia (Table 1). Notably, the average hospitalization length of MPP children with a history of pneumonia was longer than that of other patients (Table S1).

Data output and confounder assessment

A total of 8,177,030 high-quality tags were produced, averaging 41,846 (27,666–56,342), 30,848 (15,558–53,235), 67,467 (25,505–77,144), 73,856 (45,551–77,154) and 67,634 (19,862–77,454) for the NP-healthy (NP-H), OP-healthy (OP-H), NP-pneumonia (NP-P), OP-pneumonia (OP-P), and BALF groups, respectively. The average OTU numbers in the NP-H, OP-H, NP-P, OP-P, and BALF groups were 460, 343, 242, 134, and 763, respectively. Association analysis indicated that the onset of pneumonia most significantly explained the variations in microbial samples (p -value < 0.001) (Table S2). The concentration of extracted DNA in the unused sampling swabs and DNA extraction kits was lower than 0.01 ng/ μ l, whereas it was higher than 80 ng/ μ l in the sampling swabs and BALF. In addition, 16 S rRNA gene amplification on the extracted DNA indicated less than 0.01 nmol/l bacterial DNA in the enveloped sampling or extraction materials.

The alterations of NP and OP microbiota in MPP patients

There was no difference between bacterial diversity in the OP microbiota of MPP patients and that of healthy children (Fig. 1a). NP microbial samples in healthy children were divided into two clusters based on principal component analysis (PCA) (Fig. 1b), whereas OP microbial samples in healthy and diseased children were clearly separated (Fig. 1b). In parallel, the bacterial richness and evenness in the NP microbiota of the two clusters differed significantly in healthy children (Fig. 1a).


Compared to the other twenty-five NP-H samples (NP-H1), we identified a lower bacterial diversity in seven NP-H samples (NP-H2) (Fig. 1a), which were clustered in the

Table 1 Sample information

Characteristics	Pneumonia Patients (n = 28)	Healthy Children (n = 32)
Gender		
Female	12	16
Male	16	16
Age (years)	4.5(0.7–10.5)	4.0(0.3–9.9)
Delivery mode		
Cesarean section	12	9
Vaginally born	16	23
Feed pattern		
Breast feed	17	11
Breast feedmilk feed	5	15
Milk feed	6	6
Family history of allergy	0	0
History of pneumonia	4	0
Asthma	0	0
Clinical records		
Lung consolidation, atelectasis, infiltration	28	NA
Hospitalization time (days)	9.5(4–37)	–
Fever	20	–
Cough	25	–
Wheezing	2	–
CRP (<0.499mg/l)	8	NA
PCT (<0.5ng/ml)	28	NA
Eosinophil (0.5–5%)	15	NA

"–" represents not detected, NA represents not available, CRP C-response protein, PCT procalcitonin

NP-P group (Fig. 1b, p -value < 0.01). Further analysis indicated that *Staphylococcus* dominated in the NP-H2 (81.87%) and NP-P (62.28%) microbiota, whereas NP-H1 microbiota mainly comprised *Moraxella* (21.66%), *Streptococcus* (11.97%), *Corynebacterium* (11.44%) and *Dolosigranulum* (10.22%) (Fig. S1, Table S3). Conversely, the OP microbiota structure was similar in all healthy children (Table S3), comprising highly abundant *Streptococcus* (35.94%, vs. 31.32% in the OP-P microbiota, q -value = 0.93) and *Prevotella* (12.43%, vs. 6.82% in the OP-P microbiota, q -value = 0.09) (Fig. 2). In addition, *Staphylococcus*, *Corynebacterium* and *Mycoplasma* were significant components in the OP-P microbiota (Fig. 2).



Significant association of lung microbiota with disease severity in MPP patients

The hierarchical cluster analysis (Fig. 3) showed that one cluster was composed of only 12 BALF samples (BALF-1) (Fig. 3), with *Mycoplasma* being dominant (61.40%, 32.91–93.69%) in the lung microbiota (Table S4). Significantly decreased levels of *Mycoplasma* (4.08%, 0.12–32.24%) were verified in the lung microbiota in the remaining 16 BALF samples, which were identified in the NP or OP clusters (BALF-2) and featured *Haemophilus*, *Staphylococcus* or novel microbial genera as the dominant isolates in the lung microbiota (Fig. 3). The bacterial diversity in BALF-1 is higher than that in BALF-2, but the difference is not significant (Figure S2).


The peak fever was higher in BALF-1 patients than in BALF-2 patients (40.2 °C vs. 39.4 °C, p -value < 0.05) (Table S1). The length of hospitalization and length of persistent fever were also longer in BALF-1 patients (average 12.8 days and 14.4 days) compared with the BALF-2 (average 9.8 days and 7.1 days) group (p -value = 0.267 and < 0.05). In addition, we identified a higher abnormal CRP ratio in the BALF-1 group (10/12 in BALF-1, 6/16 in BALF-2) (Table S1). Subject S10, who previously had pneumonia, suffered the longest hospitalization period (37 days) and had an extremely high level of serum CRP (121.2 mg/l) during hospitalization, which may be caused by bilateral pulmonary effusion and necrotizing lung consolidation (Table S1). This can be attributed to severe

M. pneumoniae infection, with 76.39% of the lung microbiota being *Mycoplasma* (Table S4), and exposure to various empirical antibiotics (Table S1).

Similarity of microbial communities between respiratory niches in MPP patients

The similarity between upper respiratory tract (URT) and lower respiratory tract (LRT) microbiota was assessed to determine whether URT microbiota could mirror or predict microbiota imbalances in the lung. A non-metric multidimensional scaling (nMDS) analysis showed that the lung microbiota was more similar to that of the NP than the OP (Fig. 4). Additionally, NP microbial samples were clustered between BALF and OP samples (Fig. 4), which indicated the transmission of NP microbiota to both the lung and OP.

Of the top fifteen genera in URT and LRT microbiota, the abundances of *Staphylococcus*, *Corynebacterium* and *Haemophilus* were not different between the OP and lung microbiota (Table 2, Fig. S3). By comparison, we observed a decrease in *Streptococcus* and *Prevotella* in both the NP and lung microbiota (Table 2). *Mycoplasma* dominated the BALF microbiota (28.64%), which, however, was in low abundance in the URT (1.96% in NP and 0.60% in OP microbiota) (Table 2). *Ralstonia* and *Atopobium* represented $\leq 0.05\%$ of the lung microbiota, whereas *Acinetobacter* and *Buchnera* were not significantly different among the NP, OP and lung (Table 2).



Individual-specific NP and OP microbiota compared to the BALF

Bray–Curtis similarity measures were applied to assess the concordance between the NP, OP and BALF microbiota in hospitalized children. We observed variable intra-individual concordance between NP/OP and BALF, and the similarity between the NP and BALF microbiota was higher than that between the OP and BALF microbiota (Fig. 5). In the BALF-1 group, we identified a higher similarity between the NP and BALF microbiota and the OP and BALF microbiota compared with the BALF-2 group (Fig. 5). In parallel, the abundance of *Mycoplasma* was higher in the NP and OP microbiota of patients in the BALF-1 group compared to the BALF-2 group (Fig. S2, Table S4). For two individuals, S5 and S11 in the BALF-1 group, we identified *Staphylococcus* as being dominant (94.50% and 94.81%, respectively) in the lung microbiota, as well as a high similarity between the lung and NP microbiota (Fig. 5, Table S4). The present clinical tests identified *M. pneumoniae* in BALF samples even with 0.12% *Mycoplasma* (patient S4) in the lung microbiota

(Table S4), which suggested the accuracy and sensitivity of the current MPP diagnostics based on BALF. However, it is difficult to establish an *M. pneumoniae* diagnosis by NP or OP swab, given that *M. pneumoniae* represented less than 0.05% of the URT microbiota in individuals with 1.35–54.52% *M. pneumoniae* in the lung microbiota (Table S4).

Discussion

Emerging reports have demonstrated the pivotal role of respiratory microbiota in immune system education and resistance to colonization by pathogenic organisms^{2, 13}. Several studies revealed various patterns of the NP microbiota in healthy children, with different susceptibilities to ARI^{14–16}. Our study also identified the divergence of NP microbiota in healthy children, which was dominated by either *Staphylococcus* or mixed microbial components. A *Staphylococcus*-dominated NP microbiota pattern was reported to be associated with high severity of disease in ARIs, whereas a mixed NP microbiota pattern suggested a lower ARI risk^{3, 17, 18}. These differences may explain the clustering of the seven microbial samples with a *Staphylococcus*-dominated NP microbiota in the NP-P group, and it may also guide the comparison of NP microbiota structures between the two healthy groups (NP-H1 and NP-H2) and the patient group (NP-P). With only a one-week follow-up after sampling, we identified no differences in the clinical characteristics between the NP-H1 and NP-H2 groups. In contrast to the two NP groups identified, the OP microbiota in healthy children showed no significant difference. The respiratory commensals decreased significantly in MPP, including a reduced amount of the lactic acid-producing bacterium *Lactobacillus* and the short-chain fatty acid producer *Porphyromonas*^{19, 20}. *Prevotella*, which antagonizes lipid polysaccharides and inhibits mucosal inflammation, was also decreased²¹. This may provide clues about the microbial etiology in MPP at the level of the microbiota.

Food ingestion and esophageal reflux affected the oropharyngeal microbial composition, which signifies a higher bacterial richness in the OP than in NP and lung¹¹. Prevaes *et al.* demonstrated that BALF microbial samples clustered between the NP and OP, showing a higher similarity between the lung and NP microbiota than between the lung and OP microbiota in children with cystic fibrosis²². A prospective study of 112 infants indicated that there was exchange between the NP and OP microbiota in children with high ARI risk in early life²³. Our study identified the clustering of NP microbial samples between BALF and OP samples, which suggested the transmission of NP microbiota to both the OP and the lung. This may be due to nasal infection and robust pathogenic intrusion of *M. pneumoniae*²⁴, which also induced the disequilibrium of OP microbiota in diseased






Table 2 Comparison of top 15 genera between the URT and LRT in MPP patients

Genus	q-value	Nemenyi post hoc test p-value		Relative abundance %		
		NP-lung	OP-lung	lung	NP	OP
	Kruskal test of 3 sites					
<i>Staphylococcus</i>	<0.001	<0.001	0.276	12.26	62.28	10.71
<i>Corynebacterium</i>	0.005	0.008	0.871	0.43	1.28	2.11
<i>Streptococcus</i>	<0.001	0.257	<0.001	6.13	9.74	31.32
<i>Mycoplasma</i>	<0.001	<0.001	<0.001	28.64	1.96	0.60
<i>Acinetobacter</i>	0.684	0.977	0.823	3.44	1.02	5.57
<i>Actinomyces</i>	<0.001	0.026	<0.001	0.44	0.73	8.41
<i>Ralstonia</i>	<0.001	<0.001	<0.001	0.03	7.13	1.16
<i>Prevotella</i>	<0.001	0.937	<0.001	0.43	0.42	6.82
<i>Veillonella</i>	<0.001	<0.001	<0.001	0.19	0.86	6.25
<i>Haemophilus</i>	0.007	0.014	0.969	4.92	0.56	1.17
<i>Bacillus</i>	<0.001	<0.001	<0.001	5.52	0.14	0.13
<i>Rothia</i>	<0.001	0.018	<0.001	0.39	0.54	3.27
<i>Atopobium</i>	<0.001	<0.001	<0.001	0.02	0.31	2.40
<i>Neisseria</i>	0.009	0.726	0.090	1.09	0.16	1.33
<i>Buchnera</i>	0.496	0.566	1.000	0.09	2.37	0.07

p < 0.05 or q < 0.05 are shown in bold





children. Correspondingly, the NP-lung axis should be explored further to identify *M. pneumoniae* infections and the associated microbiota imbalance, as well as host responses.

Notably, MPP patients can be divided into two groups that feature different levels of *Mycoplasma* abundance in the lung microbiota. Consistent with previous studies that indicated that the abundance of either *Lactobacilli*, *Rothia* or *S. pneumoniae* was related to the pneumonia severity index (PSI)⁴, children with a higher load of *Mycoplasma* in the lung microbiota tended to suffer longer hospital stays, higher peak fevers and abnormal CRP ratios. Compared to patients with a low abundance of *Mycoplasma* in the lung microbiota, the average proportion of *Mycoplasma* in NP and OP microbiota was also greater in patients with a high abundance of *Mycoplasma* in the lung. Nonetheless, the *Mycoplasma* load in the NP and OP microbiota was not proportional to that in the lung microbiota at the individual level, suggesting the limitations of using URT microbiota to predict MPP severity. This could be explained by the differences in clearance of *M. pneumoniae* in the lung compared to the URT⁹. The complex etiology in pneumonia, such as co-infection with unknown pathogens, may explain the long hospitalization lengths and fever durations in patients with a low abundance of *M. pneumoniae* in the BALF.

Several limitations of this study should also be considered. The short-term follow-up for healthy children made it difficult to assess the susceptibility of different URT microbiota to ARI. The small sample size could impose restrictions on determining the association of respiratory microbiota with disease severity. Moreover, we could neither precisely identify the pathogens at the

species level nor reveal functional dysbiosis in the respiratory microbiota in MPP via 16 S rRNA analysis.

In conclusion, we provide significant evidence for an imbalanced URT microbiota in MPP and varied URT microbiota patterns in healthy children. More importantly, we explored the transmission of URT and LRT microbiota, which suggested an association of respiratory microbiota imbalance with severity of disease in MPP.

Materials and methods

Ethical approval

The study was approved by the Ethical Committee of Shenzhen Children's Hospital under registration number 2016013. The guardians of the recruited children provided their informed consent. All procedures performed in this study were in accordance with the ethical standards of the institutional and/or national research committee, as well as with the 1964 Helsinki declaration and its subsequent amendments or comparable ethical standards.

Sample inclusion and pathogen examination

All children with pneumonia were enrolled from inpatients at Shenzhen Children's Hospital. The inclusion criteria for patients were as follows: 1) no asthma; 2) a diagnosed *M. pneumoniae* infection via BALF detection; 3) no antibiotic exposure for at least 1 month before sampling; 4) no admission to a pediatric intensive care unit (PICU); and 5) no mechanical ventilation during hospitalization. Age-matched healthy children were recruited after physical examination in Shenzhen Children's Hospital according to the following criteria: 1) no asthma or family history of allergies; 2) no history of pneumonia; 3) no wheezing, fever, cough or other respiratory/allergic symptoms at sampling one month

prior to the study; 4) no antibiotic exposure within 1 month; and 5) no disease symptoms one week after sampling.

URT and LRT sampling was conducted 3 to 11 days after hospitalization when the diseased children maintained a high fever or had no significant disease remission (Table S1). URT samples were collected by NP (25-800-A-50, Puritan, Guilford, North Carolina, USA) and OP (155 C, COPAN, Murrieta, California, USA) swabs, and BALF was collected by fiberoptic bronchoscopy. The same BALF sample was used for both clinical pathogen detection and 16 S rRNA gene sequencing. All samples were frozen at -80°C within 10 minutes of sample collection.

Common pathogens were detected by the following methods: bacterial culturing was conducted to detect clinical common bacterial pathogens²⁵, and the nucleic acid testing (NAT) method was applied to identify the viral or atypical pathogens as described previously²⁶. Unused swabs and DNA extraction kits were utilized as negative controls to assess experimental contamination²⁶.

DNA extraction, library construction, and sequencing

The genomic DNA of the microbiota was extracted using the Power Soil DNA Isolation Kit (Mo Bio Laboratories, Carlsbad). A PCR amplicon library was constructed using the V3-V4 hypervariable region of the 16 S rRNA gene¹⁴. The qualified libraries were then sequenced with the Illumina MiSeq sequencing platform (Illumina, San Diego, USA)²⁷.

Data processing, statistical analysis, and visualization

Raw sequencing data were processed with the QIIME pipeline²⁸ and in-house scripts²⁶. Low-quality data filtration, OTU clustering, representative taxon classification and community diversity calculation were conducted as described previously^{26, 29}. PERMANOVA was used to evaluate the confounding effect of subject characteristics on microbiota composition^{30, 31}. An nMDS plot was used to visualize the concordance of the overall microbial community composition between niches. Inter-group comparisons were performed via the Wilcoxon rank-sum test. Hierarchical clustering of NP, OP and BALF microbial samples in MPP patients was conducted, and the sample backgrounds in different sub-clusters were summarized. Differences in the top fifteen genera in the NP, OP and BALF microbiota were calculated by Kruskal–Wallis tests, and all multiple test results were adjusted by the FDR method. The intra-individual distance between the BALF and NP/OP microbiota was calculated by the Bray-Curtis dissimilarity. All graphs were produced by the package ‘ggplot2’ of the R software (v3.2.3).

Data availability

All the sequencing data have been deposited in GenBank under the accession numbers SRP090593 (healthy children) and SRP130820 (children with pneumonia).

Acknowledgements

This study was supported by the Key Medical Disciplines Building Project of Shenzhen (201506053), Key Medical Disciplines Building Project of Shenzhen (SZXJ2017005), Guangdong Medical Research Fund (A2016501), Sanming Project of Medicine in Shenzhen [SZSM201512030], Shenzhen public service platform for clinical drug trials (20151964) and Shenzhen Science and Technology Project (JCYJ20170303155012371). We also would like to thank Mr. Xiaofeng Lin from EasyPub for polishing the language when preparing the first submission.

Author details

¹Department of Computer Science, City University of Hong Kong, 999077 Hong Kong, China. ²Department of Respiratory Diseases, Shenzhen Children's Hospital, No. 7019, Yitian Road, 518026 Futian District, Shenzhen, China. ³Department of Microbial Research, WeHealthGene Institute, 3C19, No. 19 Building, 518000 Dayun Software Town, Shenzhen, China. ⁴Institute of Statistics, NanKai University, No. 94 Weijin Road, 300071 Tianjin, China. ⁵Department of Respiratory Diseases, Beijing Children's Hospital, 100045 Beijing, China

Author contributions

Y.Z. and S.L. managed the project. Z.L., H.W., and G.X. enrolled the children and performed the specimen sampling and common pathogen detection. X.F. and D.L. performed the bioinformatics analysis. Q.Z. and W.D. interpreted the analysis results and wrote the paper. Y.L. and Z.Y. optimized the graphs. Y.L. polished the article. K.S. and Y.Y. provided professional suggestions and revised the manuscript. All authors reviewed this manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

Supplementary Information accompanies this paper at (<https://doi.org/10.1038/s41426-018-0097-y>).

Received: 25 January 2018 Revised: 16 April 2018 Accepted: 17 April 2018
Published online: 23 May 2018

References

- Zhou, Y. et al. Biogeography of the ecosystems of the healthy human body. *Genome Biol.* **14**, R1 (2013).
- Human Microbiome Project C. Structure, function and diversity of the healthy human microbiome. *Nature* **486**, 207–214 (2012).
- Hasegawa, K. et al. Association of nasopharyngeal microbiota profiles with bronchiolitis severity in infants hospitalised for bronchiolitis. *Eur. Respir. J.* **48**, 1329–1339 (2016).
- de Steenhuijsen Piters, W. A. et al. Dysbiosis of upper respiratory tract microbiota in elderly pneumonia patients. *ISME J.* **10**, 97–108 (2016).
- Hasegawa, K. et al. Nasal airway microbiota profile and severe bronchiolitis in infants: A case-control study. *Pediatr. Infect. Dis. J.* **36**, 1044–1051 (2017).
- Qin, Q., Xu, Baoping, Liu, X. & Shen, K. Status of *Mycoplasma pneumoniae* pneumonia in chinese children: A systematic review. *Adv. Microbiol.* **4**, 704–711 (2014).
- Bassi, C. M. et al. Analysis of the upper respiratory tract microbiotas as the source of the lung and gastric microbiotas in healthy individuals. *mBio* **6**, e00037 (2015).
- Charlson, E. S. et al. Topographical continuity of bacterial populations in the healthy human respiratory tract. *Am. J. Respir. Crit. Care Med.* **184**, 957–963 (2011).
- Meyer Sauter, P. M. et al. The role of B cells in carriage and clearance of *Mycoplasma pneumoniae* from the respiratory tract of mice. *J. Infect. Dis.* **217**, 298–309 (2018).

10. Brennan, S., Gangell, C., Wainwright, C. & Sly, P. D. Disease surveillance using bronchoalveolar lavage. *Paediatr. Respir. Rev.* **9**, 151–159 (2008).
11. Avital, A., Uwyed, K., Picard, E., Godfrey, S. & Springer, C. Sensitivity and specificity of oropharyngeal suction versus bronchoalveolar lavage in identifying respiratory tract pathogens in children with chronic pulmonary infection. *Pediatr. Pulmonol.* **20**, 40–43 (1995).
12. Huang, Y. J. Nasopharyngeal microbiota: Gatekeepers or fortune tellers of susceptibility to respiratory tract infections? *Am. J. Respir. Crit. Care Med.* **196**, 1504–1505 (2017).
13. Esposito, S. & Principi, N. Impact of nasopharyngeal microbiota on the development of respiratory tract diseases. *Eur. J. Clin. Microbiol. Infect. Dis.* **37**, 1–7 (2018).
14. Teo, S. M. et al. The infant nasopharyngeal microbiome impacts severity of lower respiratory infection and risk of asthma development. *Cell Host Microbe* **17**, 704–715 (2015).
15. Tano, K., von Essen, R., Eriksson, P. O. & Sjöstedt, A. *Alloiococcus otitidis*—otitis media pathogen or normal bacterial flora? *APMIS* **116**, 785–790 (2008).
16. Biesbroek, G. et al. Seven-valent pneumococcal conjugate vaccine and nasopharyngeal microbiota in healthy children. *Emerg. Infect. Dis.* **20**, 201–210 (2014).
17. Hasegawa, K. et al. Serum cathelicidin, nasopharyngeal microbiota, and disease severity among infants hospitalized with bronchiolitis. *J. Allergy Clin. Immunol.* **139**, 1383–1386 (2017).
18. Bogaert, D. et al. Variability and diversity of nasopharyngeal microbiota in children: A metagenomic analysis. *PLoS One.* **6**, e17035 (2011).
19. Zauwiler, T., Eichert, M., Richter, H. & Uden, G. Variations in the energy metabolism of biotechnologically relevant heterofermentative lactic acid bacteria during growth on sugars and organic acids. *Appl. Microbiol. Biotechnol.* **72**, 421–429 (2006).
20. Sato, M. et al. Three CoA transferases involved in the production of short chain fatty acids in *Porphyromonas gingivalis*. *Front. Microbiol.* **7**, 1146 (2016).
21. Miller, S. I., Ernst, R. K. & Bader, M. W. LPS, TLR4 and infectious disease diversity. *Nat. Rev. Microbiol.* **3**, 36–46 (2005).
22. Prevaes S. M., et al. Concordance between upper and lower airway microbiota in infants with cystic fibrosis. *Eur. Respir. J.* **49**, 1602235 (2017).
23. AATM, Bosch et al. . Maturation of the infant respiratory microbiota, environmental drivers and health consequences a prospective cohort study. *Am. J. Respir. Crit. Care Med.* **196**, 1582–1590 (2017).
24. Al Masalma, M. et al. The expansion of the microbiological spectrum of brain abscesses with use of multiple 16s ribosomal DNA sequencing. *Clin. Infect. Dis.* **48**, 1169–1178 (2009).
25. Pettigrew, M. M. et al. Association of sputum microbiota profiles with severity of community-acquired pneumonia in children. *BMC Infect. Dis.* **16**, 317 (2016).
26. Lu Z., et al. The alteration of nasopharyngeal and oropharyngeal microbiota in children with MPP and non-MPP. *Genes* **8**, E380 (2017).
27. Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K. & Schloss, P. D. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the Miseq Illumina sequencing platform. *Appl. Environ. Microbiol.* **79**, 5112–5120 (2013).
28. Caporaso, J. G. et al. Qiime allows analysis of high-throughput community sequencing data. *Nat. Methods* **7**, 335–336 (2010).
29. Wang, H. et al. *Mycoplasma pneumoniae* and *Streptococcus pneumoniae* caused different microbial structure and correlation network in lung microbiota. *J. Thorac. Dis.* **8**, 1316–1322 (2016).
30. Kelly, B. J. et al. Composition and dynamics of the respiratory tract microbiome in intubated patients. *Microbiome* **4**, 7 (2016).
31. Yilmaz, P. et al. The Silva and “All-species Living Tree Project (LTP)” taxonomic frameworks. *Nucleic Acids Res.* **42**, D643–D648 (2014).