



香港城市大學
City University of Hong Kong

專業 創新 胸懷全球
Professional · Creative
For The World

CityU Scholars

FoldAtlas

a repository for genome-wide RNA structure probing data

Norris, Matthew; Kwok, Chun Kit; Cheema, Jitender; Hartley, Matthew; Morris, Richard J.; Aviran, Sharon; Ding, Yiliang

Published in:
Bioinformatics

Published: 15/01/2017

Document Version:

Final Published version, also known as Publisher's PDF, Publisher's Final version or Version of Record

License:
CC BY-NC

Publication record in CityU Scholars:
[Go to record](#)

Published version (DOI):

[10.1093/bioinformatics/btw611](https://doi.org/10.1093/bioinformatics/btw611)

Publication details:

Norris, M., Kwok, C. K., Cheema, J., Hartley, M., Morris, R. J., Aviran, S., & Ding, Y. (2017). FoldAtlas: a repository for genome-wide RNA structure probing data. *Bioinformatics*, 33(2), 306-308.
<https://doi.org/10.1093/bioinformatics/btw611>

Citing this paper

Please note that where the full-text provided on CityU Scholars is the Post-print version (also known as Accepted Author Manuscript, Peer-reviewed or Author Final version), it may differ from the Final Published version. When citing, ensure that you check and use the publisher's definitive version for pagination and other details.

General rights

Copyright for the publications made accessible via the CityU Scholars portal is retained by the author(s) and/or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights. Users may not further distribute the material or use it for any profit-making activity or commercial gain.

Publisher permission

Permission for previously published items are in accordance with publisher's copyright policies sourced from the SHERPA RoMEO database. Links to full text versions (either Published or Post-print) are only available if corresponding publishers allow open access.

Take down policy

Contact lbscholars@cityu.edu.hk if you believe that this document breaches copyright and provide us with details. We will remove access to the work immediately and investigate your claim.

Databases and ontologies

FoldAtlas: a repository for genome-wide RNA structure probing data

Matthew Norris^{1,*}, Chun Kit Kwok², Jitender Cheema¹,
Matthew Hartley¹, Richard J. Morris¹, Sharon Aviran³ and
Yiliang Ding^{1,*}

¹John Innes Centre, Norwich Research Park, Norwich, UK, ²Department of Biology and Chemistry, City University of Hong Kong, Kowloon Tong, Hong Kong SAR, China and ³Department of Biomedical Engineering and Genome Center, UC Davis, Davis, CA, USA

*To whom correspondence should be addressed.

Associate Editor: Ivo Hofacker

Received on December 21, 2015; revised on August 5, 2016; accepted on September 20, 2016

Abstract

Summary: Most RNA molecules form internal base pairs, leading to a folded secondary structure. Some of these structures have been demonstrated to be functionally significant. High-throughput RNA structure chemical probing methods generate millions of sequencing reads to provide structural constraints for RNA secondary structure prediction. At present, processed data from these experiments are difficult to access without computational expertise. Here we present *FoldAtlas*, a web interface for accessing raw and processed structural data across thousands of transcripts. *FoldAtlas* allows a researcher to easily locate, view, and retrieve probing data for a given RNA molecule. We also provide *in silico* and *in vivo* secondary structure predictions for comparison, visualized in the browser as circle plots and topology diagrams. Data currently integrated into *FoldAtlas* are from a new high-depth Structure-seq data analysis in *Arabidopsis thaliana*, released with this work.

Availability and Implementation: The *FoldAtlas* website can be accessed at www.foldatlas.com. Source code is freely available at github.com/mnori/foldatlas under the MIT license. Raw reads data are available under the NCBI SRA accession SRP066985.

Contact: yiliang.ding@jic.ac.uk or matthew.norris@jic.ac.uk.

Supplementary information: [Supplementary data](#) are available at *Bioinformatics* online.

1 Introduction

RNA structure plays an important role in all steps of gene expression and regulation (Mortimer *et al.*, 2014; Sharp, 2009). Earlier studies inferred the secondary structures of individual RNA sequences using low throughput *in vitro* probing or *in silico* prediction approaches. More recently, genome-wide *in vivo* structure probing methods have emerged, allowing structures to be determined across the transcriptomes of living cells (Ding *et al.*, 2014; Rouskin *et al.*, 2014; Spitale *et al.*, 2015; Talkish *et al.*, 2014; Tang *et al.*, 2015).

Chemical probing methods can be used to determine RNA secondary structure in living cells (Kwok *et al.*, 2013; McGinnis and

Weeks, 2014; Spitale *et al.*, 2013). These methods include dimethyl sulfate (DMS) probing and selective 2'-hydroxyl acylation analyzed by primer extension (SHAPE). In DMS probing, the N1 position of adenine and the N3 position of cytosine are methylated when the base is not involved in Watson–Crick base pairing. In SHAPE, all unpaired bases are modified.

Chemically modified bases lead to stalling of reverse transcriptase. With reverse transcription, PCR, deep sequencing and normalization, reactivities can be assigned to individual RNA sequence positions. These reactivities describe the extent of exposure of a nucleotide to solution, and can be exploited as pseudo-free energy constraints for




Fig. 1 (A) d3nome genome browser, indicating splice isoforms. (B) Normalized reactivities. The sequence position is shown on the x axis, whilst the y axis provides the normalized reactivity value. (C) Principal components analysis, indicating structural similarity. Each dot represents a single RNA structure prediction, with red dots indicating the lowest free energies. Structures with similar base pair configurations are plotted in close proximity to each other. (D) Circle plot describing a single RNA structure prediction. Sequence positions are indicated around the edge of the plot, with lines between positions indicating base pairs. Green lines indicate high base pair probability. (E) RNA fold prediction diagram. Bases with high reactivities are in red, whilst green indicates little or no reactivity

RNA secondary structure prediction. At present, raw and processed reactivity data are hard to access without computational expertise.

Here we introduce *FoldAtlas*, a repository and web interface for accessing genome-scale RNA structure probing data. We also provide visualization of data-constrained RNA structures across the genome. The data included with *FoldAtlas* are from a new high-depth Structure-seq DMS analysis in *Arabidopsis thaliana*, covering over 11 000 transcripts.

2 Results

FoldAtlas allows a researcher without computational expertise to select a transcript of interest and retrieve its corresponding raw and processed structure probing data, along with pre-generated RNA structure predictions. *FoldAtlas* is the first tool that provides this functionality across the genome. When a transcript is selected and loaded, the *d3nome* genome browser (Fig. 1A) released with this work, displays the splicing configuration of the selected transcript, along with alternative splice isoforms, where relevant. An overview of the normalized chemical reactivities (Fig. 1B) is also shown, which can be expanded to show detailed nucleotide-resolution chemical reactivities. The reactivities are generated as described in the [Supplementary Results](#) section of the [Supplementary Material](#). Tab delimited text files containing normalized chemical reactivities are available for download. We also provide corresponding raw read termination counts from 3 independent biological replicates, allowing the significance of structure probing data to be estimated by assigning errors to reactivities.

For each transcript, we include the 20 lowest free energy unconstrained *in silico* and data-constrained *in vivo* structures generated by using the *Fold* program, from version 5.7 of the *RNAstructure* package (Reuter and Mathews, 2010), with default slope and intercept parameters of 1.8 and -0.6 kcal/mol respectively. The structure prediction set includes the MFE structure alongside suboptimal low free energy structures. Differences and similarities between these structures

are visualized using a Principal Components Analysis (PCA) view (Fig. 1C). PCA plots were generated using a previously described method (Halvorsen *et al.*, 2010). Each structure can also be visualized using both circle plots (Fig. 1D) and structure diagrams (Fig. 1E) generated using the *ViennaRNA* package (Hofacker, 2013; Kerpedjiev *et al.*, 2015; Lorenz *et al.*, 2011). The corresponding MFE structures can be downloaded as tab-delimited text files.

The *FoldAtlas* chemical reactivity data are from a DMS chemical modification experiment in *Arabidopsis thaliana*. These data were generated by using a previously established Structure-seq method (Ding *et al.*, 2014, 2015), but with two rounds of poly-A selection to enrich the proportion of mRNA. Detailed analysis of this experiment is provided in the [Supplementary Results](#) section of the [Supplementary Material](#).

3 Conclusions and future work

FoldAtlas provides convenient access to *in vivo* RNA structure probing data across thousands of transcripts. The current release, 1.1, includes data from a high depth genome-scale probing experiment in *Arabidopsis thaliana*. To predict structure for a transcript, we generated up to 20 secondary structures using the *RNAstructure Fold* tool, and visualized the structure ensemble using PCA plots. In this work, our preference to use *RNAstructure* is due to the ability to specify experimental constraints, and is consistent with the approach taken in our earlier work (Ding *et al.*, 2014). In future versions of *FoldAtlas*, we plan to also provide options to visualize structure predictions made using other methods, including *SeqFold* (Ouyang *et al.*, 2013), and *Vienna RNAfold*, which now allows experimental constraints (Lorenz *et al.*, 2011, 2015). We are also considering including SHAPE probing data, *in vitro* data, reactivities calculated using alternative normalization methods, data from other organisms, and data from other studies.

Acknowledgements

This research was supported in part by the NBIP Computing infrastructure for Science (CiS) group through the provision of a High Performance Computing Cluster.

Funding

This work was supported by the Biotechnology and Biological Sciences Research Council (BBSRC) [BB/L025000/1 to M.N. and Y.D. and BB/N022572/1 to M.N. Y.D and S.A.]; and the National Institutes of Health [HG006860 to S.A.].

Conflict of Interest: none declared.

References

- Ding, Y. *et al.* (2014) In vivo genome-wide profiling of RNA secondary structure reveals novel regulatory features. *Nature*, **505**, 696–700.
- Ding, Y. *et al.* (2015) Genome-wide profiling of *in vivo* RNA structure at single-nucleotide resolution using structure-seq. *Nat. Protoc.*, **10**, 1050–1066.
- Halvorsen, M. *et al.* (2010) Disease-associated mutations that alter the RNA structural ensemble. *PLoS Genet.*, **6**, e1001074.
- Hofacker, I.L. (2003) Vienna RNA secondary structure server. *Nucleic Acids Res.*, **31**, 3429–3431.
- Kerpedjiev, P. *et al.* (2015) Forna (force-directed RNA): simple and effective online RNA secondary structure diagrams. *Bioinformatics*, **31**, 3377–3379.
- Kwok, C.K. *et al.* (2013) Determination of *in vivo* RNA structure in low-abundance transcripts. *Nat. Commun.*, **4**, 2971.
- Lorenz, R. *et al.* (2011) ViennaRNA package 2.0. *Algorithms Mol. Biol.*, **6**, 26.

- Lorenz, R. *et al.* (2015) SHAPE directed RNA folding. *Bioinformatics*, *btv523*.
- McGinnis, J.L. and Weeks, K.M. (2014) Ribosome RNA assembly intermediates visualized in living cells. *Biochemistry*, *53*, 3237–3247.
- Mortimer, S.A. *et al.* (2014) Insights into RNA structure and function from genome-wide studies. *Nat. Rev. Genet.*, *15*, 469–479.
- Ouyang, *et al.* (2013) SeqFold: genome-scale reconstruction of RNA secondary structure integrating high-throughput sequencing data. *Genome Res.*, *23*, 377–387.
- Reuter, J.S. and Mathews, D.H. (2010) RNAstructure: software for RNA secondary structure prediction and analysis. *BMC Bioinformatics*, *11*, 129.
- Rouskin, S. *et al.* (2014) Genome-wide probing of RNA structure reveals active unfolding of mRNA structures in vivo. *Nature*, *505*, 701–705.
- Sharp, P.A. (2009) The centrality of RNA. *Cell*, *136*, 577–580.
- Spitale, R.C. *et al.* (2013) RNA SHAPE analysis in living cells. *Nat. Chem. Biol.*, *9*, 18–20.
- Spitale, R.C. *et al.* (2015) Structural imprints in vivo decode RNA regulatory mechanisms. *Nature*, *519*, 486–490.
- Talkish, J. *et al.* (2014) Mod-seq: high-throughput sequencing for chemical probing of RNA structure. *RNA*, *20*, 713–720.
- Tang, Y. *et al.* (2015) StructureFold: genome-wide RNA secondary structure mapping and reconstruction in vivo. *Bioinformatics*, *31*, 2668–2675.