



香港城市大學  
City University of Hong Kong

專業 創新 胸懷全球  
Professional · Creative  
For The World

## CityU Scholars

**Response to Letter to Editor by Zavascki A.P.**

**Urgent need for evaluating point-of-care tests as a RT-PCR-sparing strategy for the diagnosis of Covid-19 in symptomatic patients**

Chow, W. K.; Chow, C. L.

**Published in:**

Epidemiology and Infection

**Published:** 01/01/2021

**Document Version:**

Final Published version, also known as Publisher's PDF, Publisher's Final version or Version of Record

**License:**

CC BY-NC-ND

**Publication record in CityU Scholars:**

[Go to record](#)

**Published version (DOI):**

[10.1017/S0950268821000182](https://doi.org/10.1017/S0950268821000182)

**Publication details:**

Chow, W. K., & Chow, C. L. (2021). Response to *Letter to Editor* by Zavascki A.P. Urgent need for evaluating point-of-care tests as a RT-PCR-sparing strategy for the diagnosis of Covid-19 in symptomatic patients. *Epidemiology and Infection*, 149, [e33]. <https://doi.org/10.1017/S0950268821000182>

**Citing this paper**

Please note that where the full-text provided on CityU Scholars is the Post-print version (also known as Accepted Author Manuscript, Peer-reviewed or Author Final version), it may differ from the Final Published version. When citing, ensure that you check and use the publisher's definitive version for pagination and other details.

**General rights**

Copyright for the publications made accessible via the CityU Scholars portal is retained by the author(s) and/or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights. Users may not further distribute the material or use it for any profit-making activity or commercial gain.

**Publisher permission**

Permission for previously published items are in accordance with publisher's copyright policies sourced from the SHERPA RoMEO database. Links to full text versions (either Published or Post-print) are only available if corresponding publishers allow open access.

**Take down policy**

Contact [lbscholars@cityu.edu.hk](mailto:lbscholars@cityu.edu.hk) if you believe that this document breaches copyright and provide us with details. We will remove access to the work immediately and investigate your claim.

## Letter to the Editor

**Cite this article:** Chow WK, Chow CL (2021). Response to *Letter to Editor* by Zavascki A.P.: Urgent need for evaluating point-of-care tests as a RT-PCR-sparing strategy for the diagnosis of Covid-19 in symptomatic patients. *Epidemiology and Infection* **149**, e33, 1. <https://doi.org/10.1017/S0950268821000182>

Received: 12 January 2021  
Accepted: 12 January 2021

**Author for correspondence:**  
W.K. Chow,  
E-mail: [elize.yeung.polyu@gmail.com](mailto:elize.yeung.polyu@gmail.com)

# Response to *Letter to Editor* by Zavascki A.P.: Urgent need for evaluating point-of-care tests as a RT-PCR-sparing strategy for the diagnosis of Covid-19 in symptomatic patients

W.K. Chow<sup>1</sup>  and C.L. Chow<sup>2</sup>

<sup>1</sup>Department of Building Services Engineering, The Hong Kong Polytechnic University, Hong Kong, China and  
<sup>2</sup>Department of Architecture and Civil Engineering, City University of Hong Kong, Hong Kong, China

We agree entirely that the proposed pool testing mathematical strategy [1] is needed [2] in low-income countries, assisting the government in operating the public health system. The resources bottleneck can be solved via applying optimised testing policies in low-resource setting.

Further, if the mathematical strategy is well supported by disaster management personnel, quick, reliable and cheap detection testing schemes on coronavirus disease 2019 (COVID-19) with smaller number of tests can be worked out.

Another update on recent mandatory testing [3] in several suburbs with confirmed infection cases, the pooling testing strategy will be useful in handling thousands of tests quickly for picking up asymptomatic patients at a certain time. Although residents having negative results are not yet safe, and they are asked to do the test again a few days later.

## References

1. **Chow WK and Chow CL** (2021) A discussion on implementing pooling detection tests of novel coronavirus (SARS-CoV-2) for a large population. *Epidemiology and Infection*, **149**, e17. doi: 10.1017/S0950268820003155.
2. **Zavascki AP** (2021) Urgent need for evaluating point-of-care tests as a RT-PCR-sparing strategy for the diagnosis of Covid-19 in symptomatic patients. *Epidemiology and Infection*, 1–4. doi: 10.1017/S0950268821000170.
3. **Cheung E, Cheung T and Lam J** (2021) Hong Kong fourth wave: mandatory testing of residents in about 70 more buildings, as 56 new coronavirus cases logged. *South China Morning Post*, 19 January 2021. Available at <https://www.scmp.com/news/hong-kong/health-environment/article/3118267/hong-kong-fourth-wave-carrie-lam-pledges-stepped>.

© The Author(s), 2021. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.