



香港城市大學
City University of Hong Kong

專業 創新 胸懷全球
Professional · Creative
For The World

CityU Scholars

Do Auditory Mismatch Responses Differ Between Acoustic Features?

An, HyunJung; Ho Kei, Shing; Auksztulewicz, Ryszard; Schnupp, Jan W. H.

Published in:

Frontiers in Human Neuroscience

Published: 01/02/2021

Document Version:

Final Published version, also known as Publisher's PDF, Publisher's Final version or Version of Record

License:

CC BY

Publication record in CityU Scholars:

[Go to record](#)

Published version (DOI):

[10.3389/fnhum.2021.613903](https://doi.org/10.3389/fnhum.2021.613903)

Publication details:

An, H., Ho Kei, S., Auksztulewicz, R., & Schnupp, J. W. H. (2021). Do Auditory Mismatch Responses Differ Between Acoustic Features? *Frontiers in Human Neuroscience*, 15, [613903].
<https://doi.org/10.3389/fnhum.2021.613903>

Citing this paper

Please note that where the full-text provided on CityU Scholars is the Post-print version (also known as Accepted Author Manuscript, Peer-reviewed or Author Final version), it may differ from the Final Published version. When citing, ensure that you check and use the publisher's definitive version for pagination and other details.

General rights

Copyright for the publications made accessible via the CityU Scholars portal is retained by the author(s) and/or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights. Users may not further distribute the material or use it for any profit-making activity or commercial gain.

Publisher permission

Permission for previously published items are in accordance with publisher's copyright policies sourced from the SHERPA RoMEO database. Links to full text versions (either Published or Post-print) are only available if corresponding publishers allow open access.

Take down policy

Contact lbscholars@cityu.edu.hk if you believe that this document breaches copyright and provide us with details. We will remove access to the work immediately and investigate your claim.



Do Auditory Mismatch Responses Differ Between Acoustic Features?

HyunJung An¹, Shing Ho Kei¹, Ryszard Aukstulewicz^{1,2*†} and Jan W. H. Schnupp^{1*†}

¹ Department of Neuroscience, City University of Hong Kong, Kowloon, Hong Kong, ² Department of Neuroscience, Max Planck Institute for Empirical Aesthetics, Frankfurt, Germany

OPEN ACCESS

Edited by:

Martin Sommer,
University of Göttingen, Germany

Reviewed by:

Lavinia Slabu,
Université Côte d'Azur, France
Erich Schröger,
Leipzig University, Germany

*Correspondence:

Jan W. H. Schnupp
wschnupp@cityu.edu.hk
Ryszard Aukstulewicz
rikismo@gmail.com

[†]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Cognitive Neuroscience,
a section of the journal
Frontiers in Human Neuroscience

Received: 04 October 2020

Accepted: 07 January 2021

Published: 01 February 2021

Citation:

An H, Ho Kei S, Aukstulewicz R and
Schnupp JWH (2021) Do Auditory
Mismatch Responses Differ Between
Acoustic Features?
Front. Hum. Neurosci. 15:613903.
doi: 10.3389/fnhum.2021.613903

Mismatch negativity (MMN) is the electroencephalographic (EEG) waveform obtained by subtracting event-related potential (ERP) responses evoked by unexpected deviant stimuli from responses evoked by expected standard stimuli. While the MMN is thought to reflect an unexpected change in an ongoing, predictable stimulus, it is unknown whether MMN responses evoked by changes in different stimulus features have different magnitudes, latencies, and topographies. The present study aimed to investigate whether MMN responses differ depending on whether sudden stimulus change occur in pitch, duration, location or vowel identity, respectively. To calculate ERPs to standard and deviant stimuli, EEG signals were recorded in normal-hearing participants ($N = 20$; 13 males, 7 females) who listened to roving oddball sequences of artificial syllables. In the roving paradigm, any given stimulus is repeated several times to form a standard, and then suddenly replaced with a deviant stimulus which differs from the standard. Here, deviants differed from preceding standards along one of four features (pitch, duration, vowel or interaural level difference). The feature levels were individually chosen to match behavioral discrimination performance. We identified neural activity evoked by unexpected violations along all four acoustic dimensions. Evoked responses to deviant stimuli increased in amplitude relative to the responses to standard stimuli. A univariate (channel-by-channel) analysis yielded no significant differences between MMN responses following violations of different features. However, in a multivariate analysis (pooling information from multiple EEG channels), acoustic features could be decoded from the topography of mismatch responses, although at later latencies than those typical for MMN. These results support the notion that deviant feature detection may be subserved by a different process than general mismatch detection.

Keywords: electroencephalography, mismatch negativity, predictive coding, auditory processing, multivariate decoding

INTRODUCTION

Neural activity is typically suppressed in response to expected stimuli and enhanced following novel stimuli (Carbajal and Malmierca, 2018). This effect is often summarized as a mismatch response, calculated by subtracting the neural response waveform to unexpected deviant stimuli from the response to expected standard stimuli. Auditory deviance detection has been associated with a human auditory-evoked potential, the mismatch negativity, occurring at about 150–250 ms from sound change onset (Naatanen, 2007; Garrido et al., 2008). The principal neural sources of the MMN are thought to be superior temporal regions adjacent to the primary auditory cortex, as

well as frontoparietal areas (Doeller et al., 2003; Chennu et al., 2013). Initially, the MMN was interpreted as a correlate of pre-attentive encoding of physical features between standard and deviant sounds (Doeller et al., 2003). However, more recent studies have led to substantial revisions of this hypothesis, and currently, the most widely accepted explanation of the MMN is that it reflects a prediction error response.

An important theoretical question remains whether mismatch signaling has a domain-general or domain-specific (feature-dependent) implementation in the auditory processing pathway. A recent study using invasive recordings from the cortical surface (Aukstulewicz et al., 2018) demonstrated that neural mechanisms of predictions regarding stimulus contents (“what”) and timing (“when”) can be dissociated in terms of their topographies and latencies throughout the frontotemporal network, and that activity in auditory regions is sensitive to interactions between different kinds of predictions. Additionally, biophysical modeling of the measured signals has shown that predictions of contents and timing are best explained either by short-term plasticity or by classical neuromodulation, respectively, suggesting separable mechanisms for signaling different kinds of predictions. However, these dissociations might be specific to predictions of contents vs. timing, which may have fundamentally different roles in processing stimulus sequences (Friston and Buzsaki, 2016).

Interestingly, an earlier magnetoencephalography (MEG) study (Phillips et al., 2015) provided evidence for a hierarchical model, whereby violations of sensory predictions regarding different stimulus contents were associated with similar response magnitudes in auditory cortex, but different connectivity patterns at hierarchically higher levels of the frontotemporal network. This result is consistent with the classical predictive coding hypothesis in which reciprocal feedforward and feedback connections at the lower levels of the hierarchy are thought to signal prediction errors and predictions regarding simple sensory features, but hierarchically higher levels are thought to signal more complex predictions and prediction errors, integrating over multiple features (Kiebel et al., 2008). Several studies, however, reported independent processing of prediction violations along different acoustic features or sound dimensions. An earlier study (Giard et al., 1995) investigated the neural correlates of mismatch processing across three different acoustic features (frequency, intensity, and duration). Mismatch responses to each feature were source-localized by fitting equivalent current dipoles to EEG signals, and the results indicated that violations of different features can be linked to dissociable sources, suggesting the involvement of different underlying populations. Similar conclusions have been reached in another set of studies (Schroger, 1995; Paavilainen et al., 2001), which quantified the additivity of MMN to changes along different acoustic features, either in isolation or by combining two or more features. In these studies, the MMN response to violating two features could largely be reproduced by adding the MMN responses to violating two single features, suggesting that the latter are mutually independent. A more recent study has combined these two approaches (source localization and additivity analyses), demonstrating partial independence of three different timbre


dimensions (Caclin et al., 2006). The notion that mismatch responses to violations of different features are mediated by independent mechanisms is also supported by studies showing that MMN (as well as the later P3a component) typically decreases following two identical deviants presented in direct succession, but remains stable following two deviants which vary from the standard along different features (for a review, see Rosburg et al., 2018).

However, in most previous studies (Giard et al., 1995; Schroger, 1995; Paavilainen et al., 2001; Phillips et al., 2015; Rosburg et al., 2018), physical differences between deviants and standards were not behaviorally matched across different features or participants, raising the possibility that differences in mismatch-evoked activity might to some extent be explained by differences in stimulus salience (Shiramatsu and Takahashi, 2018). This was also the case in the more recent studies on MMN responses to multiple acoustic features (Phillips et al., 2015) or in previous roving paradigms (Garrido et al., 2008). Interestingly, a recent study investigating the MMN to acoustic violations along multiple independent features in the auditory cortex of anesthetized rats (An et al., 2020) revealed that the topography of MMN signals was highly diverse across not only acoustic features but also individual animals, even though several sources of inter-subject variability (e.g., electrode placement) were better controlled than in typical non-invasive studies, suggesting that the spatial resolution of non-invasive methods such as EEG or MEG might not be sufficient for mapping more subtle differences between mismatch responses to violations of different features. The few EEG studies that did use behaviorally matched deviant sounds across different features either used very small sample sizes ($N = 8$; Deouell and Bentin, 1998) or were limited to relatively specialized perceptual characteristics (e.g., different timbre features; Caclin et al., 2006). In contrast, our study used a larger sample size ($N = 20$) and manipulated relatively general sound dimensions (location, pitch, duration, and syllable identity). Our primary goal was to test whether mismatch responses to violations of different features differ in magnitude or latency, in an attempt to replicate previous studies (Deouell and Bentin, 1998). However, in addition to testing the effects of acoustic feature on the MMN time-course in a mass-univariate analysis (i.e., on an electrode-by-electrode basis), we also aimed at decoding acoustic features from differences in MMN topography in a multivariate analysis (i.e., pooling signals from multiple electrodes).

MATERIALS AND METHODS

Participants

Twenty volunteers (13 males and 7 females; mean age 23.9 years old) enrolled in the study upon written informed consent. All participants self-reported as having normal hearing and no history of neurological disorders, and all but two were right-handed. All participants but one were native Hong Kong residents, and their mother tongue was Cantonese. A musical training questionnaire indicated that 16 participants had no musical training, and the remaining participants had <4 years' experience in playing a musical instrument. Participants were



seated in a sound-attenuated and electrically shielded room in front of a computer screen. They were instructed to fixate on a fixation cross displayed on the screen during the acoustic stimulation. All experimental procedures were approved by the Human Subjects Ethics Sub-Committee of the City University of Hong Kong.

Stimuli

The present study employed a roving oddball paradigm in which auditory deviants could differ from preceding standards along one of four independent acoustic features. Specifically, we manipulated two consonant-vowel (CV) syllable stimuli, /ta/ and /ti/ (Retsa et al., 2018), along the following independent acoustic features: duration, pitch, interaural level difference (ILD) or vowel (An et al., 2020). Prior to the EEG recording, per participant, we estimated the feature interval yielding ~80% behavioral performance by employing a 1-up-3-down staircase procedure. In each staircase trial, two out of three stimuli, chosen at random, were presented at a mean level of a given feature (e.g., a 50/50 vowel mixture or a 0 dB ILD) while the third stimulus was higher or lower than the mean level by a certain interval. Participants had to indicate which stimulus was the “odd one out.” Following three consecutive hits, the interval decreased by 15%; following a mistake, the interval increased by 15%. Each participant performed 30 staircase trials for each feature (**Figure 1B**). For the roving oddball stimulus

sequences, the stimulus duration was set to 120 ms and the inter-stimulus intervals (ISIs) were fixed at 500 ms. Stimuli formed a roving oddball sequence: after 4–35 repetitions of a given stimulus (forming a standard), it was replaced with another (deviant) stimulus, randomly drawn from the set of 5 possible levels (**Figure 1A**). Roving oddball sequences corresponding to different features were administered in separate blocks, in a randomized order across participants. The total number of stimuli in each block was ~2,000, including 200 deviant stimuli and 200 corresponding (immediately preceding) standards.

Experimental Procedure

We recorded signals from 64 EEG channels in a 10–20 system using an ANT Neuro EEG Sports amplifier. EEG channels were grounded at the nasion and referenced to the Cpz electrode. Participants were seated in a quiet room and fitted with Brainwavz B100 earphones, which delivered the audio stimuli via a MOTU Ultralite MK3 USB soundcard at 44.1 kHz. EEG signals were pre-processed using the SPM12 Toolbox for MATLAB. The continuous signals were first notch-filtered between 48 and 52 Hz and band-pass filtered between 0.1 and 90 Hz (both filters: 5th order zero-phase Butterworth), and then downsampled to 300 Hz. Eye blinks were automatically detected using the Fp1 channel, and the corresponding artifacts were removed by subtracting the two principal spatiotemporal components associated with each eye blink from all EEG channels (Ille

et al., 2002). Then, data were re-referenced to the average of all channels, segmented into epochs ranging from -100 ms before to 400 ms after each stimulus onset, baseline-corrected to the average pre-stimulus voltage, and averaged across trials to obtain ERPs for deviants and standards for each of the four acoustic features.

Data Analyses

First, to establish the presence of the MMN response, we converted the EEG time-series into 3D images (2D spatial topography \times 1D time-course) and entered them into a general linear model (GLM) with two factors (random effect of mismatch: deviant vs. standard; fixed effect of participant), corresponding to a paired t -test. Statistical parametric maps were thresholded at an uncorrected $p < 0.005$, and the resulting spatiotemporal clusters of main effects were tested for statistical significance at the family-wise error corrected threshold $p_{FWE} < 0.05$, taking into account the spatiotemporal correlations and multiple comparisons across channels and time points.

In an additional control analysis, we have tested whether the mismatch responses observed in this study were modulated by adaptation effects, which have been shown to be especially prominent in the N1 range (Baldeweg et al., 2004). To this end, per standard stimulus (i.e., the last stimulus in a sequence of identical stimuli), we have calculated the number of stimuli separating it from the preceding deviant (i.e., the first stimulus in a sequence of identical stimuli). If our results were indeed confounded by adaptation, the difference between responses evoked by deviants vs. standards should be modulated by the number of stimuli preceding each deviant. To test this hypothesis, we have regressed out the number of preceding stimuli from single-trial standard-evoked responses (using two regressors: a linear regressor, coding for the actual number of preceding stimuli, and a log-transformed regressor, approximating empirically observed adaptation effects; e.g., Baldeweg et al., 2004), and subjected the residuals to the remaining univariate analysis steps (i.e., averaging the single-trial responses to obtain ERPs, and performing statistical inference while correcting for multiple comparisons across channels and time points).

Then, to test whether MMN amplitudes differed between stimulus features, ERP data were entered into a flexible-factorial GLM with one random factor (participant) and two fixed factors (mismatch: deviant vs. standard; feature: pitch, duration, ILD, and vowel), corresponding to a repeated-measures 2×4 ANOVA. Statistical significance thresholds were set as above.

Finally, to test whether mismatch responses can be used to decode the violated acoustic features, we subjected the data to a multivariate analysis. Prior to decoding, we calculated single-trial mismatch response signals by subtracting the EEG signal evoked by each standard from the signal evoked by the subsequent deviant. Data dimensionality was reduced using PCA (principal component analysis), resulting in spatial principal components (describing channel topographies) and temporal principal components (describing voltage time-series), sorted by the ratio of explained variance. Only those top components which, taken together, explained 95% of the original variance,


were retained for further analysis. In decoding acoustic features, we adopted a sliding window approach, integrating over the relative voltage changes within a 100 ms window around each time-point (Wolff et al., 2020). To this end, per channel and trial, the time segments within 100 ms of each analyzed time-point were down-sampled by binning the data over 10 ms bins, resulting in a vector of 10 average voltage values per component. Next, the data were de-meant by removing the component-specific average voltage over the entire 100 ms time window from each component and time bin. These steps ensured that the multivariate analysis approach was optimized for decoding transient activation patterns (voltage fluctuations around a zero mean) at the expense of more stationary neural processes (overall differences in mean voltage) (Wolff et al., 2020).

The binned single-trial mismatch fluctuations were then concatenated across components for subsequent leave-one-out cross-validation decoding. Per trial and time point, we calculated the Mahalanobis distance (De Maesschalck et al., 2000) (scaled by the noise covariance matrix of all components) between the vector of concatenated component fluctuations of this trial (test trial) and four other vectors, obtained from the remaining trials, and corresponding to the concatenated component fluctuations averaged across trials, separately for each of the four features. The resulting Mahalanobis distance values were averaged across trials, separately for each acoustic feature, resulting in 4×4 distance matrices. These distance matrices were summarized per time point and participant as a single decoding estimate, by subtracting the mean off-diagonal from diagonal terms (Figure 3A).

In a final analysis, since we have observed univariate mismatch responses as well as multivariate mismatch-based feature decoding at similar latencies (see Results), we have tested whether these two effects are related. To this end, we performed a correlation analysis between single-trial decoding estimates (i.e., the relative Mahalanobis distance values between EEG topography corresponding to mismatch responses following violations of the same vs. different features), and single-trial MMN amplitudes. We calculated Pearson correlation coefficients across single trials, per channel, time point, and participants. The resulting correlation coefficients were subject to statistical inference using statistical parametric mapping (one-sample t -test; significance thresholds as in the other univariate analysis, corrected for multiple comparisons across time points and channels using family-wise error).

RESULTS

Taken together, in this study, we tested whether auditory mismatch responses are modulated by violations of independent acoustic features. First, consistent with previous literature (Doeller et al., 2003; Garrido et al., 2008), we observed overall differences between the ERPs evoked by deviant stimuli vs. standard stimuli, in a range typical for MMN responses as well as at longer latencies (Figure 2A). Specifically, the univariate ERP analysis confirmed that EEG amplitudes differed significantly between deviants and standards when pooling over




all the acoustic features tested. This effect was observed over two clusters: the central EEG channels showed a significant mismatch response between 115 and 182 ms (cluster-level pFWE < 0.001 , $T_{max} = 3.94$), while posterior channels showed a significant mismatch response between 274 and 389 ms (cluster-level pFWE < 0.001 , $T_{max} = 5.46$), within the range of a P3b component. A control analysis, in which we controlled for single-trial adaptation effect to the standard tones, yielded a virtually identical pattern of results as the original analysis (two significant clusters of differences between responses to deviants vs. standards: an earlier cluster between 130 and 143 ms over central channels, cluster-level pFWE < 0.001 , $T_{max} = 15.48$, and a later cluster between 317 and 327 ms over posterior channels, cluster-level pFWE < 0.001 , $T_{max} = 17.48$).

Although the ERP time-courses differed between deviant and standard stimuli when pooling over violations of different acoustic features, a univariate (channel-by-channel) analysis revealed no significant differences in the amplitudes or time-courses of mismatch responses between independent stimulus features (**Figure 2B**). These results are consistent with a previous study (Phillips et al., 2015) which found that multiple deviant stimulus features (frequency, intensity, location, duration, and silent gap) were not associated with differences in activity in the

auditory regions, but instead were reflected in more distributed activity patterns (frontotemporal connectivity estimates).

The resulting decoding time-courses of each participant were entered into a GLM and subject to one-sample t -tests, thresholded at an uncorrected $p < 0.05$ and correcting for multiple comparisons across time points at a cluster-level pFWE < 0.05 . In this analysis, significant acoustic feature decoding was observed between 247 and 350 ms relative to tone onset (cluster-level pFWE = 0.000, $T_{max} = 2.77$) (**Figure 3B**). Thus, when pooling information from multiple EEG channels, acoustic features could be decoded from the topography of mismatch responses, although at later latencies than typical for MMN.

Since we have observed both univariate mismatch responses and multivariate mismatch-based feature decoding at late latencies (univariate: 274–389 ms; multivariate: 247–350 ms), we have performed an additional single-trial correlation analysis to test whether these two effects are related. This analysis (**Figure 3C**) has yielded no significant clusters of correlation coefficients between single-trial mismatch amplitudes and decoding estimates, while correcting for multiple comparisons across channels and time points ($T_{max} = 3.74$, all pFWE > 0.005).



DISCUSSION

In this study, since a univariate analysis of interactions between mismatch signals and acoustic features might not be sensitive enough to reveal subtle and distributed amplitude differences between conditions, we adopted a multivariate analysis aiming at decoding the violated acoustic feature from single-trial mismatch response topographies. This demonstrated that acoustic features could be decoded from the topography of mismatch responses, although at later latencies than typical for MMN (**Figure 3B**). An earlier oddball study (Leung et al., 2012) examined ERP differences to violations of four features (frequency, duration, intensity, and interaural difference). The study found that frequency deviants were associated with a significant amplitude change in the middle latency range. This result indicated that deviant feature detection may be subserved by a different process than general mismatch detection. Consistent with this notion,

another study has used magnetoencephalography to identify mid-latency effects of local prediction violations of simple stimulus features, and contrasted them with later effects of global prediction violations of stimulus patterns (Recasens et al., 2014). Taken together, these studies would suggest that, in paradigms where multiple acoustic features vary independently (such as here), a plausible pattern of results would be that independent feature predictions should be mismatched at relatively early latencies, since an integrated representation is not required. Here, however, we found feature-specificity in the late latency range, rather than in the mid-latency range. The discrepancy between our results and the previous studies might be explained by different stimulus types. While the previous studies used simple acoustic stimuli, here we used complex syllable stimuli, possibly tapping into the later latencies of language-related mismatch responses, as compared to MMN following violations of non-speech sounds.

Speech sounds have been hypothesized to be processed in separate streams which independently derive semantic information (“what” processing) and sound location (“where” processing) (Kaas and Hackett, 2000; Tian et al., 2001; Schubotz et al., 2003; Camalier et al., 2012; Kusmirek and Rauschecker, 2014). In most animal studies, the hierarchical organization of the auditory cortex has been linked to a functional distribution of stimulus processing, such that core (hierarchically lower) regions respond preferentially to simple stimuli, whereas belt and other downstream (hierarchically higher) regions respond to more complex stimuli such as band-passed noise and speech (Rauschecker et al., 1995; Recanzone et al., 2000; Rauschecker and Tian, 2004; Kusmirek and Rauschecker, 2009; Rauschecker and Scott, 2009). This is supported by evidence functional magnetic resonance imaging (fMRI) studies in humans (Binder et al., 2000) showing that earlier auditory regions (Heschl’s gyrus and surrounding fields) respond preferentially to unstructured noise stimuli, while progressively more complex stimuli such as frequency-modulated tones show more lateral response activation patterns. In that study, speech sounds showed most pronounced activations spreading ventrolaterally into the superior temporal sulcus. This result supports a hierarchical model of auditory speech processing in the human auditory cortex based on complexity and integration of temporal and spectral features. Based on this notion, the relatively long latency of neural responses compared to previous studies using pure tones might be partially explained by the fact that we used spectrally and temporally complex speech stimuli.

However, our results can also be explained in terms of a hierarchical deviance detection system based on predictive coding (Kiebel et al., 2008). On this account, neural responses supporting the lower and higher hierarchical stages communicate continuously through reciprocal pathways. When exposed to repetitive stimuli, the bottom-up (ascending) sensory inputs can be “explained away” by top-down (descending) connections mediating prediction signaling, resulting in weaker prediction error signaling back to the hierarchically higher regions. Substituting the predicted standard with unpredicted deviant results in a failure of top-down suppression by prior predictions. This leads to an increased prediction error signaling back to higher regions, providing an update for subsequent predictions. As a result, the later and more distributed activity patterns might reflect higher-order prediction errors, signaled to regions integrating multiple stimulus features and representing the entire range of stimuli likely to appear in a particular context.

REFERENCES

- An, H., Aukstulewicz, R., Kang, H., and Schnupp, J. W. H. (2020). Cortical mapping of mismatch responses to independent acoustic features. *Hear. Res.* 399:107894. doi: 10.1016/j.heares.2020.107894
- Aukstulewicz, R., Schwiedrzik, C. M., Thesen, T., Doyle, W., Devinsky, O., Nobre, A. C., et al. (2018). Not all predictions are equal: “What” and “When” predictions modulate activity in auditory cortex through different mechanisms. *J. Neurosci.* 38, 8680–8693. doi: 10.1523/JNEUROSCI.0369-18.2018
- Baldeweg, T., Klugman, A., Gruzelier, J., and Hirsch, S. R. (2004). Mismatch negativity potentials and cognitive impairment in schizophrenia. *Schizophr. Res.* 69, 203–217. doi: 10.1016/j.schres.2003.09.009
- Binder, J. R., Frost, J. A., Hammeke, T. A., Bellgowan, P. S., Springer, J. A., Kaufman, J. N., et al. (2000). Human temporal lobe activation by speech and nonspeech sounds. *Cereb. Cortex* 10, 512–528. doi: 10.1093/cercor/10.5.512
- Caclin, A., Brattico, E., Tervaniemi, M., Naatanen, R., Morlet, D., Giard, M. H., et al. (2006). Separate neural processing of timbre dimensions in auditory sensory memory. *J. Cogn. Neurosci.* 18, 1959–1972. doi: 10.1162/jocn.2006.18.12.1959

In conclusion, the present study identified functional dissociations between deviance detection and deviance feature detection. First, while mismatch responses were observed at latencies typical for the MMN as well as at longer latencies, channel-by-channel analyses revealed no robust differences between mismatch responses following violations of different acoustic features. However, we demonstrate that acoustic features could be decoded at longer latencies based on fine-grained spatiotemporal patterns of mismatch responses. This finding suggests that deviance feature detection might be mediated by later and more distributed neural responses than deviance detection itself.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Human Subjects Ethics Sub-Committee of the City University of Hong Kong. The patients/participants provided their written informed consent to participate in this study. Written informed consent was obtained from the individual(s) for the publication of any potentially identifiable images or data included in this article.

AUTHOR CONTRIBUTIONS

HA: formal analysis, writing original draft, conceptualization, and conducted experiment. SH: conducted experiment and formal analysis. RA: formal analysis, supervision, project administration, and conceptualization. JS: project administration, conceptualization, investigation, and supervision. All authors contributed to the article and approved the submitted version.

FUNDING

This work has been supported by the European Commission’s Marie Skłodowska-Curie Global Fellowship (750459 to RA), the Hong Kong General Research Fund (11100518 to RA and JS), and a grant from European Community/Hong Kong Research Grants Council Joint Research Scheme (9051402 to RA and JS).

- Camalier, C. R., D'Angelo, W. R., Sterbing-D'Angelo, S. J., de la Mothe, L. A., and Hackett, T. A. (2012). Neural latencies across auditory cortex of macaque support a dorsal stream supramodal timing advantage in primates. *Proc. Natl. Acad. Sci. U.S.A.* 109, 18168–18173. doi: 10.1073/pnas.1206387109
- Carbajal, G. V., and Malmierca, M. S. (2018). The neuronal basis of predictive coding along the auditory pathway: from the subcortical roots to cortical deviance detection. *Trends Hear.* 22:2331216518784822. doi: 10.1177/2331216518784822
- Chennu, S., Noreika, V., Gueorguiev, D., Blenkmann, A., Kochen, S., Ibanez, A., et al. (2013). Expectation and attention in hierarchical auditory prediction. *J. Neurosci.* 33, 11194–11205. doi: 10.1523/JNEUROSCI.0114-13.2013
- De Maesschalck, R., Jouan-Rimbaud, D., and Massart, D. L. (2000). The Mahalanobis distance. *Chemomet. Intellig. Lab. Syst.* 50, 1–18. doi: 10.1016/S0169-7439(99)00047-7
- Deouell, L. Y., and Bentin, S. (1998). Variable cerebral responses to equally distinct deviance in four auditory dimensions: a mismatch negativity study. *Psychophysiology* 35, 745–754. doi: 10.1111/1469-8986.3560745
- Doeller, C. F., Opitz, B., Mecklinger, A., Krick, C., Reith, W., and Schroger, E. (2003). Prefrontal cortex involvement in preattentive auditory deviance detection: neuroimaging and electrophysiological evidence. *Neuroimage* 20, 1270–1282. doi: 10.1016/S1053-8119(03)00389-6
- Friston, K., and Buzsaki, G. (2016). The functional anatomy of time: what and when in the brain. *Trends Cogn. Sci.* 20, 500–511. doi: 10.1016/j.tics.2016.05.001
- Garrido, M. I., Friston, K. J., Kiebel, S. J., Stephan, K. E., Baldeweg, T., and Kilner, J. M. (2008). The functional anatomy of the MMN: a DCM study of the roving paradigm. *Neuroimage* 42, 936–944. doi: 10.1016/j.neuroimage.2008.05.018
- Giard, M. H., Lavikainen, J., Reinikainen, K., Perrin, F., Bertrand, O., Pernier, J., et al. (1995). Separate representation of stimulus frequency, intensity, and duration in auditory sensory memory: an event-related potential and dipole-model analysis. *J. Cogn. Neurosci.* 7, 133–143. doi: 10.1162/jocn.1995.7.2.133
- Ille, N., Berg, P., and Scherg, M. (2002). Artifact correction of the ongoing EEG using spatial filters based on artifact and brain signal topographies. *J. Clin. Neurophysiol.* 19, 113–124. doi: 10.1097/00004691-200203000-00002
- Kaas, J. H., and Hackett, T. A. (2000). Subdivisions of auditory cortex and processing streams in primates. *Proc. Natl. Acad. Sci. U.S.A.* 97, 11793–11799. doi: 10.1073/pnas.97.22.11793
- Kiebel, S. J., Daunizeau, J., and Friston, K. J. (2008). A hierarchy of time-scales and the brain. *PLoS Comput. Biol.* 4:e1000209. doi: 10.1371/journal.pcbi.1000209
- Kusmirek, P., and Rauschecker, J. P. (2009). Functional specialization of medial auditory belt cortex in the alert rhesus monkey. *J. Neurophysiol.* 102, 1606–1622. doi: 10.1152/jn.00167.2009
- Kusmirek, P., and Rauschecker, J. P. (2014). Selectivity for space and time in early areas of the auditory dorsal stream in the rhesus monkey. *J. Neurophysiol.* 111, 1671–1685. doi: 10.1152/jn.00436.2013
- Leung, S., Cornella, M., Grimm, S., and Escera, C. (2012). Is fast auditory change detection feature specific? An electrophysiological study in humans. *Psychophysiology* 49, 933–942. doi: 10.1111/j.1469-8986.2012.01375.x
- Naatanen, R. (2007). The mismatch negativity - where is the big fish? *J. Psychophysiol.* 21, 133–137. doi: 10.1027/0269-8803.21.34.133
- Paavilainen, P., Valppu, S., and Naatanen, R. (2001). The additivity of the auditory feature analysis in the human brain as indexed by the mismatch negativity: 1+1 approximate to 2 but 1+1+1 < 3. *Neurosci. Lett.* 301, 179–182. doi: 10.1016/S0304-3940(01)01635-4
- Phillips, H. N., Blenkmann, A., Hughes, L. E., Bekinschtein, T. A., and Rowe, J. B. (2015). Hierarchical organization of frontotemporal networks for the prediction of stimuli across multiple dimensions. *J. Neurosci.* 35, 9255–9264. doi: 10.1523/JNEUROSCI.5095-14.2015
- Rauschecker, J. P., and Scott, S. K. (2009). Maps and streams in the auditory cortex: nonhuman primates illuminate human speech processing. *Nat. Neurosci.* 12, 718–724. doi: 10.1038/nn.2331
- Rauschecker, J. P., and Tian, B. (2004). Processing of band-passed noise in the lateral auditory belt cortex of the rhesus monkey. *J. Neurophysiol.* 91, 2578–2589. doi: 10.1152/jn.00834.2003
- Rauschecker, J. P., Tian, B., and Hauser, M. (1995). Processing of complex sounds in the macaque nonprimary auditory cortex. *Science* 268, 111–114. doi: 10.1126/science.7701330
- Recanzone, G. H., Guard, D. C., and Phan, M. L. (2000). Frequency and intensity response properties of single neurons in the auditory cortex of the behaving macaque monkey. *J. Neurophysiol.* 83, 2315–2331. doi: 10.1152/jn.2000.83.4.2315
- Recasens, M., Grimm, S., Wollbrink, A., Pantev, C., and Escera, C. (2014). Encoding of nested levels of acoustic regularity in hierarchically organized areas of the human auditory cortex. *Hum. Brain Mapp.* 35, 5701–5716. doi: 10.1002/hbm.22582
- Rettsa, C., Matusz, P. J., Schnupp, J. W. H., and Murray, M. M. (2018). What's what in auditory cortices? *Neuroimage* 176, 29–40. doi: 10.1016/j.neuroimage.2018.04.028
- Rosburg, T., Weigl, M., Thiel, R., and Mager, R. (2018). The event-related potential component P3a is diminished by identical deviance repetition, but not by non-identical repetitions. *Exp. Brain Res.* 236, 1519–1530. doi: 10.1007/s00221-018-5237-z
- Schroger, E. (1995). Processing of auditory deviants with changes in one versus two stimulus dimensions. *Psychophysiology* 32, 55–65. doi: 10.1111/j.1469-8986.1995.tb03406.x
- Schubotz, R. I., von Cramon, D. Y., and Lohmann, G. (2003). Auditory what, where, and when: a sensory somatotopy in lateral premotor cortex. *Neuroimage* 20, 173–185. doi: 10.1016/S1053-8119(03)00218-0
- Shiramatsu, T. I., and Takahashi, H. (2018). Mismatch negativity in rat auditory cortex represents the empirical salience of sounds. *Front. Neurosci.* 12:924. doi: 10.3389/fnins.2018.00924
- Tian, B., Reser, D., Durham, A., Kustov, A., and Rauschecker, J. P. (2001). Functional specialization in rhesus monkey auditory cortex. *Science* 292, 290–293. doi: 10.1126/science.1058911
- Wolff, M. J., Kandemir, G., Stokes, M. G., and Akyurek, E. G. (2020). Unimodal and bimodal access to sensory working memories by auditory and visual impulses. *J. Neurosci.* 40, 671–681. doi: 10.1523/JNEUROSCI.1194-19.2019

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 An, Ho Kei, Aukstulewicz and Schnupp. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.