



香港城市大學
City University of Hong Kong

專業 創新 胸懷全球
Professional · Creative
For The World

CityU Scholars

Undetected Circulation of African Swine Fever in Wild Boar, Asia

Vergnean, Timotheé; Guinat, Claire; Pfeiffer, Dirk U.

Published in:

Emerging Infectious Diseases

Published: 01/10/2020

Document Version:

Final Published version, also known as Publisher's PDF, Publisher's Final version or Version of Record

License:

CC BY

Publication record in CityU Scholars:

[Go to record](#)

Published version (DOI):

[10.3201/EID2610.200608](https://doi.org/10.3201/EID2610.200608)

Publication details:

Vergnean, T., Guinat, C., & Pfeiffer, D. U. (2020). Undetected Circulation of African Swine Fever in Wild Boar, Asia. *Emerging Infectious Diseases*, 26(10), 2480-2482. <https://doi.org/10.3201/EID2610.200608>

Citing this paper

Please note that where the full-text provided on CityU Scholars is the Post-print version (also known as Accepted Author Manuscript, Peer-reviewed or Author Final version), it may differ from the Final Published version. When citing, ensure that you check and use the publisher's definitive version for pagination and other details.

General rights

Copyright for the publications made accessible via the CityU Scholars portal is retained by the author(s) and/or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights. Users may not further distribute the material or use it for any profit-making activity or commercial gain.

Publisher permission

Permission for previously published items are in accordance with publisher's copyright policies sourced from the SHERPA RoMEO database. Links to full text versions (either Published or Post-print) are only available if corresponding publishers allow open access.

Take down policy

Contact lbscholars@cityu.edu.hk if you believe that this document breaches copyright and provide us with details. We will remove access to the work immediately and investigate your claim.

- bovine PrP transgenic mice. *Arch Virol.* 2003;148:677–91. <https://doi.org/10.1007/s00705-002-0958-4>
7. Bett C, Piccardo P, Cervenak J, Torres J-M, Asher DM, Gregori L. Both murine host and inoculum modulate expression of experimental variant Creutzfeldt-Jakob disease. *J Gen Virol.* 2018;99:422–33. <https://doi.org/10.1099/jgv.0.001017>
 8. Gray JG, Dudas S, Czup S. A study on the analytical sensitivity of 6 BSE tests used by the Canadian BSE reference laboratory. *PLoS One.* 2011;6:e17633. <https://doi.org/10.1371/journal.pone.0017633>
 9. Orrú CD, Groveman BR, Raymond LD, Hughson AG, Nonno R, Zou W, et al. Bank vole prion protein as an apparently universal substrate for RT-QuIC-based detection and discrimination of prion strains. *PLoS Pathog.* 2015;11:e1004983. <https://doi.org/10.1371/journal.ppat.1004983>
 10. Huang Y, Forshee RA, Keire D, Lee S, Gregori L, Asher DM, et al. Assessment of risk of variant Creutzfeldt-Jakob disease (vCJD) from use of bovine heparin. *Pharmacoepidemiol Drug Saf.* 2020;29:575–81. <https://doi.org/10.1002/pds.4982>

Address for correspondence: Cyrus Bett, US Food and Drug Administration, 10903 New Hampshire Ave, Silver Spring, MD 20993, USA; email: cyrus.bett@fda.hhs.gov

Undetected Circulation of African Swine Fever in Wild Boar, Asia

Timothée Vergne, Claire Guinat, Dirk U. Pfeiffer

Author affiliations: National Veterinary School of Toulouse, Toulouse, France (T. Vergne, C. Guinat); Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement, Toulouse (T. Vergne, C. Guinat); City University of Hong Kong, Hong Kong, China (D.U. Pfeiffer)

DOI: <https://doi.org/10.3201/eid2610.200608>

African swine fever is a growing threat to the livestock industry. We examined data indicating that in most countries in Asia, most notified events were related to farm outbreaks; meanwhile, only a few wild boar cases were reported. We hypothesize the virus circulates unnoticed in wild boar populations in Asia.

African swine fever (ASF) is one of the greatest threats to the livestock industry worldwide. Since 2007, ASF virus (ASFV) has been reported in 34

countries in Europe and Asia (1). Some strains of ASFV can be associated with case-fatality ratios of almost 100% and with economic damage caused by trade disruptions (2). The absence of a safe and effective vaccine and the evolving understanding of the epidemiologic role of wild boar have complicated efforts to control this disease (3).

Observations from the Russian Federation, Ukraine, and Romania have suggested that ASFV primarily circulates among small pig farms and spills over into commercial farms occasionally and into wild boar populations regularly (4). However, more recent reports from several countries in Europe, including the Baltic states and Belgium, suggest the virus might maintain itself within wild boar populations and occasionally spill over into domestic pig farms. This newly described epidemiologic cycle proposes direct transmission among wild boar and indirect intraspecific transmission through contaminated wild boar carcasses (4). This cycle also suggests the persistence of ASFV in wild boar populations even in the context of low wild boar density and despite high death rates caused by the disease (5).

In August 2018, ASFV was detected in China, the leading pig producer worldwide. Since then, the virus has also been reported in Mongolia, Vietnam, Cambodia, Hong Kong, North Korea, South Korea, Laos, Myanmar, the Philippines, Timor-Leste, Papua New Guinea, Indonesia, and India (1). The large-scale spread of ASFV demonstrates the challenges of controlling the disease in this region, which has a high density of domestic pigs, a high proportion of low biosecurity farms, a widespread practice of feeding pigs with food waste (6), and a complex pork trade network involving wet markets and slaughterhouses with poor hygiene (2).

Maps of predicted habitat suitability suggest that most areas of East and Southeast Asia are highly suitable for wild boar (7). Although information is limited about the spatial distribution of wild boar in Asia, studies suggest that in some regions of China, wild boar density could be similar (8) to that of eastern Poland, where ASFV has circulated in wild boar for ≥ 5 years (9). Reports of crop losses caused by wild boar in China (10) indicate that close interactions between wild boar and human activities occur in the region.

The widespread presence of ASFV in pigs in Asia implies regular environmental spillover from the pig supply chain. Therefore, it is highly likely that ASFV is already widely circulating within some wild boar populations in Asia, causing substantial wild boar death. Because life expectancy of ASFV-infected




Figure. Proportion of wild boar cases out of the total number of reported African swine fever events in the most affected countries in Europe (orange) and Asia (blue). Numbers in parentheses at right side of bars indicate the reported number of wild boar cases and the reported number of outbreaks in farms, from the date of the first reported ASF event in these countries through May 8, 2020 (1).

animals is very short, the most effective way to conduct surveillance in boar populations is to monitor reports of infected boar carcasses. Numbers of reported ASFV wild boar cases and farm outbreaks vary by nation. Except for South Korea, which reported 605 infected wild boar carcasses and 17 outbreaks in domestic pig farms from the time of the emergence of the virus in Asia until May 8, 2020, countries in Asia reported a much lower ratio of wild boar cases to farm outbreaks than their counterparts in Europe (Figure) (1). In these countries (excluding South Korea), there were only 23 reports of infected wild boar carcasses (in China and Laos only) despite 843 official ASF notifications of farm outbreaks after the virus emerged in Asia in August 2018 (Figure) (1).

The near absence of notifications of ASFV-infected wild boar cases in Asia highlights shortcomings in surveillance for ASF in wildlife, jeopardizing the success of ASF control policies in the region. We believe this lack of surveillance partly results from government division of responsibilities; in most countries, responsibility for ASF surveillance in livestock belongs to a different government department than that for monitoring wildlife populations (6). We argue that long-term success of ASF control in Asia is possible only with risk-based ASF surveillance in wild

boar populations by a multisectoral effort of wildlife and agricultural departments. Although surveillance of wild boar is a necessary component of an ASF control strategy, it must be complemented by effective ASF control measures in domestic pigs, such as improved regional coordination, increased resources for surveillance, incentives for farmers to report outbreaks, and enforcement of interventions (2). Without these measures, the region might become a major hub of ASFV infection.

About the Author

Dr. Vergne is an associate professor of veterinary public health at the National Veterinary School of Toulouse, Toulouse, France. His primary research interests include disease transmission and improving surveillance and intervention strategies for animal infectious diseases.

References

1. Food and Agriculture Organization of the United Nations. EMPRES Global Animal Disease Information System [cited 2020 May 8]. <http://empres-i.fao.org/eipws3g>
2. Dixon LK, Stahl K, Jori F, Vial L, Pfeiffer DU. African swine fever epidemiology and control. *Annu Rev Anim Biosci*. 2020;8:221–6. PubMed <https://doi.org/10.1146/annurev-animal-021419-083741>

3. Gavrier-Widén D, Ståhl K, Dixon L. No hasty solutions for African swine fever. *Science*. 2020;367:622–4. <https://doi.org/10.1126/science.aaz8590>
4. Chenais E, Ståhl K, Guberti V, Depner K. Identification of wild boar-habitat epidemiologic cycle in African swine fever epizootic. *Emerg Infect Dis*. 2018;24:810–2. <https://doi.org/10.3201/eid2404.172127>
5. Abrahantes JC, Gogin A, Richardson J, Gervelmeyer A, et al.; European Food Safety Authority. Epidemiological analyses on African swine fever in the Baltic countries and Poland. *EFSA Journal*. 2017;15:e04732. <https://doi.org/10.2903/j.efsa.2017.4732>
6. Vergne T, Chen-Fu C, Li S, Cappelle J, Edwards J, Martin V, et al. Pig empire under infectious threat: risk of African swine fever introduction into the People's Republic of China. *Vet Rec*. 2017;181:117. <https://doi.org/10.1136/vr.103950>
7. Bosch J, Iglesias I, Muñoz MJ, de la Torre A. A cartographic tool for managing African swine fever in Eurasia: mapping wild boar distribution based on the quality of available habitats. *Transbound Emerg Dis*. 2017;64:1720–33. <https://doi.org/10.1111/tbed.12559>
8. Wu S, Chen H, Cai X. Preliminary study on the population structures and reproductive habit in wild boar (*Sus scrofa*) in Dawaling Natural Reserve. *Acta Theriologica Sinica*. 2000;20:151–6.
9. Śmietanka K, Woźniakowski G, Kozak E, Niemczuk K, Frączyk M, Bocian Ł, et al. African swine fever epidemic, Poland, 2014–2015. *Emerg Infect Dis*. 2016;22:1201–7. <https://doi.org/10.3201/eid2207.151708>
10. Liu Q, Yan K, Lu Y, Li M, Yan Y. Conflict between wild boars (*Sus scrofa*) and farmers: distribution, impacts, and suggestions for management of wild boars in the Three Gorges Reservoir Area. *Journal of Mountain Science*. 2019;16:2404–16. <https://doi.org/10.1007/s11629-019-5453-4>

Address for correspondence: Timothée Vergne, UMR ENVT-INRAE 1225, National Veterinary School of Toulouse, 23 chemin des Capelles, 31300 Toulouse, France; email: timothee.vergne@envt.fr

Review of Mental Health Response to COVID-19, China

Adriana Miu, Hui Cao, Bohan Zhang,¹ Huaiyu Zhang

Author affiliations: University of Texas Southwestern Medical Center, Dallas, Texas, USA (A. Miu); Beijing Institute of Education, Beijing, China (H. Cao); Lesley University, Cambridge, Massachusetts, USA (B. Zhang); University of California San Francisco, San Francisco, California, USA (H. Zhang)

DOI: <https://doi.org/10.3201/eid2610.201113>

¹Current affiliation: Private practice, Nanjing, China.

Public mental health response to coronavirus disease is essential. After reviewing systemic and local efforts in China, we found efficient coordination and human resources. We recommend better symptom assessment, monitoring of organizations, and basic needs protection. This recommendation can inform how other countries can overcome mental health challenges during this pandemic.

The coronavirus disease (COVID-19) outbreak and quarantines have caused major distress in China (1,2). Therefore, effective public mental health response to COVID-19 is needed (3). We review systemic and local mental health efforts in China based on psychiatric emergency guidelines from the Inter-Agency Standing Committee (4). These guidelines are coordination between multiple sectors; human resources; assessment, monitoring, and evaluation; and protection and human rights standards. Our discussion will inform mental health response for the COVID-19 pandemic.

Mental health efforts in China have been coordinated and facilitated through multiple systems, including government, academic societies, universities, hospitals, and nonprofit organizations (5). Services include a countrywide 24/7 hotline, text support through apps, psychoeducation materials, and webinars (5). The government prioritized psychosocial support for COVID-19, as shown by the National Health Commission mandate requiring all mental health associations to provide psychosocial support, establish professional focus groups, and aid the provincial and city health departments (6).

Academic organizations in psychology (Chinese Psychological Society [CPS]) (Table) and psychiatry (Chinese Society of Psychiatry) provide evidence-based guidelines on psychosocial support and training (5,7). The Ministry of Education (MoE) has mandated all college counselors across the nation to volunteer for the primary Huazhong University hotline at the epicenter in Wuhan (8). At the systemic level, there is good coordination and resource allocation. The government agencies coordinate human resources, and academic associations provide professional knowledge and guidelines for frontline effort.

Coordination and resource allocation were compiled from local efforts at the Wuhan epicenter (Appendix, <https://wwwnc.cdc.gov/EID/article/26/10/20-1113-App1.pdf>). On January 23, 2020, immediately after the quarantine, Zhongnan Hospital and the Hubei Psychological Consultant Association began offering hotline services. As of April 30, more than 2,000 persons had been served. Beyond the hotline, Wuhan University and Huazhong