



香港城市大學  
City University of Hong Kong

專業 創新 胸懷全球  
Professional · Creative  
For The World

## CityU Scholars

### Potential for zoonotic transmission of *Brachyspira pilosicoli*

Hampson, David J.; Oxberry, Sophy L.; La, Tom

**Published in:**

Emerging Infectious Diseases

**Published:** 01/05/2006

**Document Version:**

Final Published version, also known as Publisher's PDF, Publisher's Final version or Version of Record

**License:**

CC BY

**Publication record in CityU Scholars:**

[Go to record](#)

**Published version (DOI):**

[10.3201/eid1205.051180](https://doi.org/10.3201/eid1205.051180)

**Publication details:**

Hampson, D. J., Oxberry, S. L., & La, T. (2006). Potential for zoonotic transmission of *Brachyspira pilosicoli*. *Emerging Infectious Diseases*, 12(5), 869-870. <https://doi.org/10.3201/eid1205.051180>

**Citing this paper**

Please note that where the full-text provided on CityU Scholars is the Post-print version (also known as Accepted Author Manuscript, Peer-reviewed or Author Final version), it may differ from the Final Published version. When citing, ensure that you check and use the publisher's definitive version for pagination and other details.

**General rights**

Copyright for the publications made accessible via the CityU Scholars portal is retained by the author(s) and/or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights. Users may not further distribute the material or use it for any profit-making activity or commercial gain.

**Publisher permission**

Permission for previously published items are in accordance with publisher's copyright policies sourced from the SHERPA RoMEO database. Links to full text versions (either Published or Post-print) are only available if corresponding publishers allow open access.

**Take down policy**

Contact [lbscholars@cityu.edu.hk](mailto:lbscholars@cityu.edu.hk) if you believe that this document breaches copyright and provide us with details. We will remove access to the work immediately and investigate your claim.

6. Koprowski H. The mouse inoculation test. In: Meslin FX, Kaplan MM, Koprowski H, editors. *Laboratory techniques in rabies*. Geneva: World Health Organization; 1996. p. 80–7.
7. Orciari LA, Niezgodna M, Hanlon CA, Shaddock JH, Sanderlin DW, Yager PA, et al. Rapid clearance of SAG-2 rabies virus from dogs after oral vaccination. *Vaccine*. 2001;19:4511–8.
8. Centers for Disease Control and Prevention. Investigation of rabies infections in organ donor and transplant recipients—Alabama, Arkansas, Oklahoma, and Texas, 2004. *MMWR Morb Mortal Wkly Rep*. 2004;53:1–3.
9. Willoughby RE Jr, Tieves KS, Hoffman GM, Ghanayem NS, Amlie-Lefond CM, Schwabe MJ, et al. Survival after treatment of rabies with induction of coma. *N Engl J Med*. 2005;352:2508–14.
10. David D, Yakobson B, Rotenberg D, Dveres N, Davidson I, Stram Y. Rabies virus detection by RT-PCR in decomposed naturally infected brains. *Vet Microbiol*. 2002;87:111–8.

Address for correspondence: Rafael Oliveira, Instituto Pasteur, Laboratory of Diagnosis, Av Paulista, 393 Cerqueira Cesar São Paulo, São Paulo 01311-001, Brazil; email: rafaeldenovais@yahoo.com.br

---

## Potential for Zoonotic Transmission of *Brachyspira pilosicoli*

**To the Editor:** Anaerobic intestinal spirochetes of the genus *Brachyspira* colonize the large intestine (1). Most *Brachyspira* species have a restricted host range, whereas *Brachyspira* (formerly *Serpulina*) *pilosicoli* colonizes a variety of animal and bird species and humans. *B. pilosicoli* is an important colonic pathogen of pigs and chickens (2). It occurs at high prevalence rates in humans in developing countries and

in male homosexuals and HIV-positive persons in industrialized countries (3). Its potential as a human pathogen was emphasized after its identification in the bloodstream of a series of debilitated persons (4).

*B. pilosicoli* isolates from humans and other species have been used experimentally to colonize chicks, piglets, and mice (5–7). While these results indicate that the *B. pilosicoli* strains used lacked host-species specificity, few data exist on whether natural zoonotic spread of *B. pilosicoli* strains occurs. In 1 study that used pulsed-field gel electrophoresis (PFGE) to type isolates from Papua New Guinea, 2 dogs were colonized with *B. pilosicoli* isolates with the same PFGE types as those from villagers. However, the higher prevalence of colonization with *B. pilosicoli* in humans than dogs suggested that the dogs were infected with human isolates, probably through consumption of human feces (8).

Multilocus enzyme electrophoresis (MLEE) has been used to study variation in *B. pilosicoli* isolates; most studies have focused on isolates from only 1 or 2 host species (8–10). Generally, *B. pilosicoli* isolates are diverse, and a lack of linkage disequilibrium in the MLEE data for human isolates suggests that the species is recombinant (8).

We used MLEE to investigate relationships between 107 *B. pilosicoli* isolates of diverse geographic and host-species origins and the *B. aalborgi* type strain (NCTC 11492<sup>T</sup>). Isolates were selected on the basis of their diverse origins and availability in the Murdoch University culture collection. They originated from feces of 34 pigs, 19 chickens, 13 ducks, 1 rhea, 25 humans, and 4 dogs; from 7 human blood samples; and from 4 water sources frequented by waterfowl. Isolates originated from Australia, Canada, France, Italy, the Netherlands, Oman, Papua New Guinea, the United Kingdom, and the United States.

The MLEE method used was as previously described (8–10); the electrophoretic mobility of 15 constitutive enzymes was analyzed. Variations in electrophoretic mobility were interpreted as representing products of different alleles at each enzyme locus. Isolates with identical enzymatic profiles at 15 loci were grouped into an electrophoretic type (ET). Genetic distance between ETs was calculated as the proportions of loci at which dissimilar alleles occurred. PHYLIP version 3.51c (Phylogeny Inference Package, University of Washington, Seattle, WA, USA) was used to analyze data and generate a dendrogram by using the unweighted pair-group method with arithmetic mean clustering fusion strategy. Genetic diversity (h) was calculated for the number of ETs as  $(1 - \sum p_i^2)/(n/n - 1)$ , where  $p_i$  is the frequency of the indicated allele and  $n$  is the number of ETs.

*B. pilosicoli* isolates were divided into 80 ETs (mean 1.35 isolates per ET) (Figure). *B. aalborgi* NTCC 11492<sup>T</sup> was distinct in ET81. The *B. pilosicoli* isolates were diverse, with an h value of 0.41. Generally, they did not cluster according to host species of origin, and isolates from a given species were distributed throughout the dendrogram. Isolates from birds were more diverse than those from humans and pigs. Eight ETs contained multiple isolates, in each case from the same host species (either chickens or pigs). In 4 cases these originated from different countries: ET47 contained 2 Australian porcine isolates and 2 from the United States; ET53 contained 2 Australian porcine isolates and Scottish porcine type strain P43/6/78<sup>T</sup>; ET54 contained 2 Australian and 2 Canadian porcine isolates; ET65 contained 1 Dutch and 1 US chicken isolate.

Although human isolates did not share an ET with isolates from other species, they were frequently closely related, differing in 1 allele. This occurred with US and Australian pig




Figure. Dendrogram showing relationships between 107 isolates of *Brachyspira pilosicoli* originating from various host species located in electrophoretic types (ETs) 1–80 and *B. aalborgi* NCTC 11492T located in ET81.

isolates in ET47 and a human isolate from Oman in ET48; an Australian pig isolate in ET61 and a UK human isolate in ET62; an isolate from an Australian HIV-positive person in ET64, and 1 Dutch and 1 US chicken isolate in ET65; and a Papua New Guinea canine isolate in ET68 and a French human blood isolate in ET69.

The distribution continuum of isolates of diverse host species and geographic origin was consistent with a lack of species specificity and suggests that *B. pilosicoli* isolates naturally have the potential to be transmitted between species. Even should there be some unexpected species-specific barrier preventing “true” animal or bird isolates from colonizing humans, animals have been colonized

by human isolates, and thus could act as a reservoir of these for subsequent retransmission to humans. The results suggest that zoonotic transfer of *B. pilosicoli* isolates likely occurs in nature, e.g., after exposure to infected animals or birds, their feces, or contaminated water.

David J. Hampson,\*

Sophy L. Oxberry,\* and Tom La\*

\*Murdoch University, Murdoch, Western Australia, Australia

#### References

1. Stanton TB. Physiology of ruminal and intestinal spirochaetes. In: Hampson DJ, Stanton TB, editors. Intestinal spirochaetes in domestic animals and humans. Wallingford (UK): CAB International; 1997. p. 7–45.

2. Hampson DJ, Duhamel GE. Porcine colonic spirochetosis/intestinal spirochetosis. In: Straw B, Zimmerman JJ, D’Allaire S, Taylor DJ, editors. Diseases of swine. 9th ed. Ames (IA): Iowa State University Press; 2006. p. 755–67.
3. Hampson DJ. Intestinal spirochaetes. In: McIver CJ, editor. A compendium of laboratory diagnostic methods for common and unusual enteric pathogens—an Australian perspective. Sydney: ASM Publications; 2005. p. 101–8.
4. Trott DJ, Jensen NS, Saint Girons I, Oxberry SL, Stanton TB, Lindquist D, Hampson DJ. Identification and characterization of *Serpulina pilosicoli* isolates from the blood of critically-ill patients. J Clin Microbiol. 1997;35:482–5.
5. Trott DJ, McLaren AJ, Hampson DJ. Pathogenicity of human and porcine intestinal spirochaetes in day-old specific pathogen free chicks: an animal model of intestinal spirochetosis. Infect Immun. 1995;63:3705–10.
6. Trott DJ, Huxtable CR, Hampson DJ. Experimental infection of newly weaned pigs with human and porcine strains of *Serpulina pilosicoli*. Infect Immun. 1996;64:4648–54.
7. Sacco RE, Trampel DW, Wannemuehler MJ. Experimental infection of C3H mice with avian, porcine, or human isolates of *Serpulina pilosicoli*. Infect Immun. 1997;65:5349–53.
8. Trott DJ, Mikosza ASJ, Combs BG, Oxberry SL, Hampson DJ. Population genetic analysis of *Serpulina pilosicoli* and its molecular epidemiology in villages in the Eastern Highlands of Papua New Guinea. Int J Syst Bacteriol. 1998;48: 659–68.
9. Lee JJ, Hampson DJ, Lymbery AJ, Harders SJ. The porcine intestinal spirochaetes: identification of new genetic groups. Vet Microbiol. 1993;34:273–85.
10. McLaren AJ, Trott DJ, Swayne DE, Oxberry SL, Hampson DJ. Genetic and phenotypic characterization of intestinal spirochetes colonizing chickens, and allocation of known pathogenic isolates to three distinct genetic groups. J Clin Microbiol. 1997;35:412–7.

Address for correspondence: David J. Hampson, School of Veterinary and Biomedical Sciences, Murdoch University, Murdoch, Western Australia 6150, Australia; email: d.hampson@murdoch.edu.au

All material published in Emerging Infectious Diseases is in the public domain and may be used and reprinted without special permission; proper citation, however, is required.